

## RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

### Tribenzoatobismuth(III)

Tumanov, Nikolay A.; Timakova, Evgenia V.; Boldyreva, Elena V.

*Published in:*

Acta crystallographica. Section E: Structure reports

*DOI:*

[10.1107/S1600536810035543](https://doi.org/10.1107/S1600536810035543)

*Publication date:*

2010

#### [Link to publication](#)

*Citation for published version (HARVARD):*

Tumanov, NA, Timakova, EV & Boldyreva, EV 2010, 'Tribenzoatobismuth(III): A new polymorph', *Acta crystallographica. Section E: Structure reports*, vol. 66, no. 10. <https://doi.org/10.1107/S1600536810035543>

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Tribenzoatobismuth(III): a new polymorph

Nikolay A. Tumanov,<sup>a\*</sup> Evgenia V. Timakova<sup>b</sup> and Elena V. Boldyreva<sup>a,b</sup>

<sup>a</sup>REC-008, Novosibirsk State University, Pirogova 2, Novosibirsk 630090, Russian Federation, and <sup>b</sup>Institute of Solid State Chemistry and Mechanochemistry SB RAS, Kutateladze 18, Novosibirsk 630128, Russian Federation

Correspondence e-mail: n.tumanov@gmail.com


Received 26 July 2010; accepted 3 September 2010

Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(\text{C}-\text{C}) = 0.013$  Å;  $R$  factor = 0.042;  $wR$  factor = 0.049; data-to-parameter ratio = 16.1.

A new polymorph ( $\beta$ ) was obtained for an active pharmaceutical ingredient, bismuth tribenzoate,  $[\text{Bi}(\text{C}_6\text{H}_5\text{CO}_2)_3]$ . The new  $\beta$ -polymorph is 1.05 times denser than the previously known polymorph [Rae *et al.* (1998). *Acta Cryst.* **B54**, 438–442]. In the  $\beta$ -polymorph, the Bi atom is linked with three benzoate anions, each of them acting as a bidentate ligand, and these assemblies with  $C_3$  point symmetry can be considered as ‘molecules’. The structure of the  $\beta$ -polymorph has no polymeric chains, in contrast to the previously known polymorph. The ‘molecules’ in the  $\beta$ -polymorph are stacked along [001], so that the phenyl rings of the neighbouring molecules are parallel to each other. Based on the pronounced difference in the crystal structures, one can suppose that two polymorphs should differ in the dissolution kinetics and bioavailability.

### Related literature

The synthesis of the complex is described by Timakova *et al.* (2010). For background to bismuth complexes, see: Mehring (2007); Kislichenko (1999); Goddard *et al.* (2003); Alcock (1972). For the previously known polymorph, see: Rae *et al.* (1998). For related structures, see: Hanawalt *et al.* (1938); Rosmann *et al.* (1995).



### Experimental

#### Crystal data

$[\text{Bi}(\text{C}_6\text{H}_5\text{O}_2)_3]$	$Z = 3$
$M_r = 572.31$	Mo $K\alpha$ radiation
Trigonal, $R\bar{3}$	$\mu = 9.77 \text{ mm}^{-1}$
$a = 19.5608$ (19) Å	$T = 295$ K
$c = 4.0967$ (5) Å	$0.07 \times 0.01 \times 0.01$ mm
$V = 1357.5$ (2) Å <sup>3</sup>	

#### Data collection

Oxford Diffraction Gemini R Ultra diffractometer	3909 measured reflections
Absorption correction: multi-scan ( <i>CrysAlis PRO</i> ; Oxford Diffraction, 2010)	1355 independent reflections
$T_{\text{min}} = 0.954$ , $T_{\text{max}} = 1.000$	1076 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.087$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$	$\Delta\rho_{\text{max}} = 1.52 \text{ e \AA}^{-3}$
$wR(F^2) = 0.049$	$\Delta\rho_{\text{min}} = -0.68 \text{ e \AA}^{-3}$
$S = 0.76$	Absolute structure: Flack (1983), 622 Friedel pairs
1355 reflections	Flack parameter: $-0.034$ (12)
84 parameters	
H-atom parameters constrained	

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010) and *enCIFer* (Allen *et al.*, 2004).

This work was supported by a grant from BRHE (RUX0-008-NO-06) and the program of the Presidium of Russian Academy of Sciences ‘Basic research for medicine’; the diffractometer was purchased using money from the Innovation Project of Rosobrazovanie #456 (2007–2009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2224).

### References

- Alcock, N. W. (1972). *Advances in Inorganic Chemistry and Radiochemistry*, edited by H. J. Emeleus & A. G. Sharpe, pp. 1–58. New York: Academic Press.
- Allen, F. H., Johnson, O., Shields, G. P., Smith, B. R. & Towler, M. (2004). *J. Appl. Cryst.* **37**, 335–338.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Goddard, P. J., Klinger, J. D., Dhal, P. K., Mandeville, H. W. III, Fitzpatrick, R. J. & Neenan, X. T. (2003). US Patent 6565895.
- Hanawalt, J. D., Rihh, H. W. & Frevel, L. K. (1938). *Ind. Eng. Chem.* **10**, 457–512.
- Kislichenko, V. S. (1999). *Provizor*, **12**, 38–40.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Mehring, M. (2007). *Coord. Chem. Rev.* **251**, 974–1006.
- Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
- Rae, A. D., Gainsford, G. J. & Kemmitt, T. (1998). *Acta Cryst.* **B54**, 438–442.
- Rosmann, D., Hiibler, K. & Schmidt, A. (1995). *Monatsh. Chem.* **126**, 1045–1050.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Timakova, E. V., Udalova, T. A. & Yuhin, Yu. M. (2010). *Chem. Sustainable Dev.* **18**, 79–86.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

**supplementary materials**

*Acta Cryst.* (2010). E66, m1248 [ doi:10.1107/S1600536810035543 ]

## Tribenzoatobismuth(III): a new polymorph

N. A. Tumanov, E. V. Timakova and E. V. Boldyreva

### Comment

Bismuth tribenzoate, as other bismuth salts, is an active pharmaceutical ingredient, which has anti-infective and analgesic effects on lymphatic tissue and mucous tunic (Kislichenko, 1999), (Goddard *et al.*, 2003). Besides, it can be used as precursor for fine-dispersed powders of the metallic bismuth and its oxides (Mehring, 2007).

While developing a new method of synthesis of the title salt (Timakova *et al.*, 2010), we have discovered, that the powder diffraction pattern of the compound obtained in our experiments **I** did not match the one calculated based on the single-crystal diffraction data for the known monoclinic polymorph of bismuth tribenzoate, which was prepared by anionic exchange of bismuth triacetate with benzoic acid (Rae *et al.*, 1998). At the same time, the powder diffraction pattern of **I** agreed well with the powder diffraction pattern reported in a much earlier publication (Hanawalt *et al.*, 1938). We have managed to select a small crystal from the batch of the synthesized compound **I** of quality suitable for a single-crystal diffraction study. The powder diffraction pattern calculated based on the results of the structure solution using this single-crystal agreed well with the diffraction pattern of the whole polycrystalline batch, thus proving, that the crystal was representative for the whole sample. We report herein the crystal structure of the  $\beta$ -polymorph of bismuth tribenzoate.

The new  $\beta$ -polymorph **I** is 1.05 times denser than the previously known polymorph **II** with the space group  $P2_1/m$  (Rae *et al.*, 1998): the density of **I** at room temperature is 2.098 g/cm<sup>3</sup>, and that of **II** at 173 K is 1.99 g/cm<sup>3</sup>. The structure of **I** is isomorphous to the structure of antimony tribenzoate (Rosmann *et al.*, 1995). In **I** each bismuth atom is linked to three benzoate anions, which act as bidentate ligands (Bi1—O1 distance is 2.254 (5)Å) and Bi1—O2 distance is 2.513 (5)Å). These assemblies with  $C_3$  point symmetry can be considered as 'molecules' (Fig. 1). The 'molecules' in **I** are stacked along [0 0 1], so that the phenyl rings of the neighbouring 'molecules' are parallel to each other (Fig. 2). The carboxylate group in a ligand is rotated at 13.3 (3)° relative to phenyl ring. The neighbouring 'molecules' can be supposed to interact noticeably with each other, as far as one can judge from the intermolecular Bi1...O1<sup>i</sup> [symmetry code: (i)  $x, y, z-1$ ] distances (3.110 (6)Å), which are shorter, than the sum of van der Waals radii of Bi and O (3.67Å; Alcock, 1972). Still, the structure of **I** has no polymeric chains, in contrast to the **II**. Phenyl ring of each benzoic anion is placed between two benzoic cations of neighbouring 'molecules' (Fig. 3). In the structure of **II** each bismuth atom is also linked to three benzoate ligands, however, O atoms act as bridges between the neighbouring Bi atoms, thus forming infinite polymeric chains along  $a$  axis. No individual molecules can be selected in **II**. Based on the pronounced difference in the crystal structures, one can suppose that two polymorphs should differ in the dissolution kinetics and bioavailability.

### Experimental

The title salt was precipitated from bismuth perchloric acid solution (prepared from 217 ml of distilled water, 483 ml of concentrated perchloric acid and 850 g of bismuth oxide, then diluted to 1:10 with water) by benzoic acid with the mole ratio of benzoate-ions to bismuth being equal to 3. The reaction was carried out at 343 K during 1 h. The obtained white powder of bismuth tribenzoate was filtered and washed once with water and dried at room temperature.

## supplementary materials

Analysis found: Bi 36.6 (7), C 43.9 (4), H 2.65 (4); calculated Bi 36.5, C 44.07, H 2.64.

The typical particle size in the powder sample was in the range 0.002–0.01 mm, but a few larger crystals could be found. The largest of the selected crystals (0.07×0.01×0.01 mm) was fixed at a Mitigen MicroMesh holder with cryoil and used for a single-crystal X-ray diffraction study. Powder X-ray diffraction (Stoe Stadi MP, Cu K $\alpha_1$ , curved germanium monochromator (111), linear PSD) has proved that the selected crystal was representative for the whole powder sample batch.

### Refinement

After the positions of Bi atoms were determined by direct methods, carbon and oxygen atoms could be located from difference Fourier maps one after another in several cycles of refinement. Hydrogen atoms of the aromatic ring were placed geometrically with C–H distance 0.93 Å with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . The highest peak at the electron density map is located at 1.23 Å distance from Bi1 atom and could be a consequence of the Fourier sum truncation.

### Figures




Fig. 1. An ellipsoid plot of the title compound, showing the atom-numbering scheme and 50% probability displacement ellipsoids. Hydrogen atoms were shown as small spheres of arbitrary radii and were not labeled for clarity. Symmetry codes: (i)  $-1-x+y, 1-x, z$ ; (ii)  $1-y, 2+x-y, z$ .




Fig. 2. The stacking of bismuth tribenzoate 'molecules' along [001] in the title compound. Intermolecular Bi1...O1 contacts shown as blue dashed lines. Hydrogen atoms were omitted for clarity.




Fig. 3. The crystal packing of the title compound viewed along the  $c$  axis. Hydrogen atoms were omitted for clarity.

### Tribenzoatobismuth(III)

#### Crystal data

[Bi(C<sub>7</sub>H<sub>5</sub>O<sub>2</sub>)<sub>3</sub>]

$M_r = 572.31$

Trigonal,  $R\bar{3}$

$D_x = 2.098 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 964 reflections

Hall symbol: R 3	$\theta = 3.6\text{--}28.0^\circ$
$a = 19.5608 (19) \text{ \AA}$	$\mu = 9.77 \text{ mm}^{-1}$
$c = 4.0967 (5) \text{ \AA}$	$T = 295 \text{ K}$
$V = 1357.5 (2) \text{ \AA}^3$	Needle, colourless
$Z = 3$	$0.07 \times 0.01 \times 0.01 \text{ mm}$
$F(000) = 816$	

### Data collection

Oxford Diffraction Gemini R Ultra diffractometer	1355 independent reflections
Radiation source: fine-focus sealed tube graphite	1076 reflections with $I > 2\sigma(I)$
Detector resolution: $10.3457 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.087$
$\omega$ scans	$\theta_{\text{max}} = 28.0^\circ, \theta_{\text{min}} = 3.6^\circ$
Absorption correction: multi-scan ( <i>Crys.Alis PRO</i> ; Oxford Diffraction, 2010)	$h = -25 \rightarrow 20$
$T_{\text{min}} = 0.954, T_{\text{max}} = 1.000$	$k = -16 \rightarrow 25$
3909 measured reflections	$l = -5 \rightarrow 5$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.042$	H-atom parameters constrained
$wR(F^2) = 0.049$	$w = 1/[\sigma^2(F_o^2) + (0.P)^2]$
$S = 0.76$	where $P = (F_o^2 + 2F_c^2)/3$
1355 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
84 parameters	$\Delta\rho_{\text{max}} = 1.52 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.68 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 622 Friedel pairs
	Flack parameter: $-0.034 (12)$

### Special details

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

## supplementary materials

---

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Bi1	0.0000	1.0000	0.0000	0.02741 (15)
O1	0.0761 (3)	0.9819 (3)	0.3632 (14)	0.0364 (15)
O2	0.0144 (3)	0.8798 (3)	0.0486 (14)	0.0401 (15)
C1	0.0647 (5)	0.9145 (5)	0.271 (2)	0.0275 (19)
C2	0.1132 (4)	0.8836 (4)	0.4092 (19)	0.0264 (19)
C7	0.1798 (5)	0.9308 (5)	0.585 (2)	0.036 (2)
H7	0.1952	0.9835	0.6222	0.043*
C6	0.2236 (5)	0.8990 (5)	0.7059 (19)	0.038 (2)
H6	0.2682	0.9307	0.8307	0.045*
C3	0.0925 (5)	0.8056 (5)	0.3449 (19)	0.041 (2)
H3	0.0483	0.7739	0.2184	0.049*
C4	0.1376 (6)	0.7756 (5)	0.469 (2)	0.048 (3)
H4	0.1233	0.7232	0.4309	0.058*
C5	0.2038 (6)	0.8234 (6)	0.649 (2)	0.043 (2)
H5	0.2349	0.8038	0.7311	0.052*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Bi1	0.02780 (19)	0.02780 (19)	0.0267 (3)	0.01390 (9)	0.000	0.000
O1	0.035 (4)	0.030 (4)	0.043 (4)	0.015 (3)	0.005 (3)	-0.001 (3)
O2	0.034 (4)	0.045 (4)	0.042 (4)	0.020 (3)	0.000 (3)	-0.013 (3)
C1	0.027 (5)	0.028 (5)	0.030 (5)	0.016 (4)	0.006 (4)	0.007 (4)
C2	0.023 (5)	0.026 (5)	0.029 (5)	0.011 (4)	0.003 (4)	0.001 (4)
C7	0.040 (6)	0.036 (5)	0.040 (6)	0.025 (5)	0.002 (4)	0.000 (4)
C6	0.035 (5)	0.040 (6)	0.040 (6)	0.020 (5)	-0.002 (4)	0.001 (5)
C3	0.055 (6)	0.046 (6)	0.035 (6)	0.035 (5)	0.002 (5)	0.005 (5)
C4	0.066 (7)	0.035 (6)	0.056 (7)	0.035 (6)	0.009 (5)	0.014 (5)
C5	0.050 (6)	0.061 (7)	0.040 (6)	0.043 (6)	0.008 (5)	0.019 (5)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Bi1—O1	2.254 (5)	C5—C6	1.349 (10)
Bi1—O2	2.513 (5)	C6—C7	1.379 (10)
Bi1—C1	2.782 (8)	C7—C2	1.364 (10)
O1—C1	1.280 (8)	C7—H7	0.9300
O2—C1	1.260 (8)	C6—H6	0.9300
C1—C2	1.472 (10)	C3—H3	0.9300
C2—C3	1.394 (10)	C4—H4	0.9300
C3—C4	1.378 (10)	C5—H5	0.9300
C4—C5	1.372 (12)		
O1—Bi1—O1 <sup>i</sup>	81.2 (2)	C4—C5—C6	119.8 (8)
O1—Bi1—O2 <sup>i</sup>	76.50 (17)	C5—C6—C7	122.0 (8)
O1—Bi1—O2	53.78 (18)	C6—C7—C2	118.8 (8)

O1 <sup>i</sup> —Bi1—O2	131.88 (18)	C7—C2—C3	119.8 (8)
O2 <sup>i</sup> —Bi1—O2	119.38 (4)	C2—C7—H7	120.6
C1—O1—Bi1	100.2 (5)	C6—C7—H7	120.6
C1—O2—Bi1	88.5 (4)	C5—C6—H6	119.0
O2—C1—O1	117.2 (7)	C7—C6—H6	119.0
O2—C1—C2	123.1 (7)	C4—C3—H3	120.0
O1—C1—C2	119.5 (8)	C2—C3—H3	120.0
C1—C2—C7	121.2 (7)	C5—C4—H4	120.2
C1—C2—C3	118.9 (7)	C3—C4—H4	120.2
C2—C3—C4	120.0 (8)	C6—C5—H5	120.1
C3—C4—C5	119.6 (8)	C4—C5—H5	120.1
O1 <sup>i</sup> —Bi1—O1—C1	165.2 (4)	O2—C1—C2—C7	163.1 (8)
O1 <sup>ii</sup> —Bi1—O1—C1	82.8 (5)	O1—C1—C2—C7	-12.8 (11)
O2 <sup>i</sup> —Bi1—O1—C1	-140.1 (5)	O2—C1—C2—C3	-13.7 (11)
O1—Bi1—O2—C1	-3.1 (4)	O1—C1—C2—C3	170.4 (7)
O2 <sup>ii</sup> —Bi1—O1—C1	102.3 (5)	C3—C2—C7—C6	-2.5 (12)
O1 <sup>i</sup> —Bi1—O2—C1	-27.3 (5)	C1—C2—C7—C6	-179.2 (7)
O1 <sup>ii</sup> —Bi1—O2—C1	-92.0 (5)	C2—C7—C6—C5	1.7 (13)
O2 <sup>i</sup> —Bi1—O2—C1	38.8 (5)	C7—C2—C3—C4	2.4 (12)
O2 <sup>ii</sup> —Bi1—O2—C1	-125.6 (4)	C1—C2—C3—C4	179.2 (8)
Bi1—O2—C1—O1	5.0 (7)	C2—C3—C4—C5	-1.5 (13)
Bi1—O2—C1—C2	-171.0 (7)	C7—C6—C5—C4	-0.9 (13)
Bi1—O1—C1—O2	-5.7 (8)	C3—C4—C5—C6	0.8 (13)
Bi1—O1—C1—C2	170.5 (6)		

Symmetry codes: (i)  $-y+1, x-y+2, z$ ; (ii)  $-x+y-1, -x+1, z$ .

Fig. 1




Fig. 2




Fig. 3

