

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Biodiversity mediates the effects of stressors but not nutrients on litter decomposition

Beaumelle, Léa; De Laender, Frederik; Eisenhauer, Nico

Published in:
eLife

DOI:
[10.7554/eLife.55659](https://doi.org/10.7554/eLife.55659)

Publication date:
2020

Document Version
Peer reviewed version

[Link to publication](#)

Citation for published version (HARVARD):

Beaumelle, L, De Laender, F & Eisenhauer, N 2020, 'Biodiversity mediates the effects of stressors but not nutrients on litter decomposition', *eLife*, vol. 9, e55659, pp. 1-40. <https://doi.org/10.7554/eLife.55659>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Biodiversity mediates the effects of stressors but not nutrients on litter decomposition

Léa Beaumelle^{1,2,§*}, Frederik De Laender³, Nico Eisenhauer^{1,2}

*For correspondence:

lea.beaumelle@gmail.com (LB)

Present address: [§]INRAE, UMR SAVE, Villenave d'Ornon, France

¹ German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany; ² Institute of Biology, Leipzig University, Deutscher Platz 5e, 04103 Leipzig, Germany; ³ Research Unit of Environmental and Evolutionary Biology, Namur Institute of Complex Systems, and Institute of Life, Earth, and the Environment, University of Namur, Rue de Bruxelles 61, 5000, Namur, Belgium

Abstract Understanding the consequences of ongoing biodiversity changes for ecosystems is a pressing challenge. Controlled biodiversity-ecosystem function experiments with random biodiversity loss scenarios have demonstrated that more diverse communities usually provide higher levels of ecosystem functioning. However, it is not clear if these results predict the ecosystem consequences of environmental changes that cause non-random alterations in biodiversity and community composition. We synthesized 69 independent studies reporting 660 observations of the impacts of two pervasive drivers of global change (chemical stressors and nutrient enrichment) on animal and microbial decomposer diversity and litter decomposition. Using meta-analysis and structural equation modelling, we show that declines in decomposer diversity and abundance explain reduced litter decomposition in response to stressors but not to nutrients. While chemical stressors generally reduced biodiversity and ecosystem functioning, detrimental effects of nutrients occurred only at high levels of nutrient inputs. Thus, more intense environmental change does not always result in stronger responses, illustrating the complexity of ecosystem consequences of biodiversity change. Overall, these findings provide strong empirical evidence for significant real-world biodiversity-ecosystem functioning relationships when human activities decrease biodiversity. This highlights that the ecosystem consequences of observed biodiversity change are nontrivial and depend on the kind of environmental change.

Introduction

Human activities cause global environmental changes with important consequences for biodiversity and the functioning of ecosystems. Understanding these consequences is crucial for better policy and conservation strategies, which will ultimately promote human well-being too (*IPBES, 2019*). A key question is to what extent changes in ecosystem functioning are mediated by changes at which dimensions of biodiversity. Extensive research has demonstrated that biodiversity is needed for the stable provenance and enhancement of ecosystem processes and functions (*Cardinale et al., 2012; Schuldt et al., 2018; Tilman et al., 2012*). However, this body of evidence is mostly based on experiments comparing ecosystem functioning in artificial communities with varying number of species. Such experiments might not capture the complex ways by which shifts in biodiversity induced by global change ultimately affect ecosystem functioning (*De Laender et al., 2016; Eisenhauer et al., 2019b*).

41 Early biodiversity-ecosystem function (BEF) experiments typically controlled for environmen-
 42 tal gradients, thus not accounting for the underlying drivers of biodiversity change (*De Laender*
 43 *et al.*, 2016; *Srivastava and Vellend*, 2005; *Wardle*, 2016). These early experiments also focused on
 44 species richness as the sole biodiversity index, and manipulated it directly and randomly. How-
 45 ever, environmental change will often elicit non-random changes in several facets of biodiversity
 46 (*Eisenhauer et al.*, 2016; *Giling et al.*, 2019; *van der Plas*, 2019) (community composition and pop-
 47 ulation densities (*Glassman et al.*, 2018; *Spaak et al.*, 2017), functional diversity (*Cadotte et al.*,
 48 2011; *Craven et al.*, 2018; *Heemsbergen et al.*, 2004), trophic diversity (*Soliveres et al.*, 2016; *Wang*
 49 *and Brose*, 2018; *Zhao et al.*, 2019), or intra-specific diversity (*Des Roches et al.*, 2018)). The se-
 50 lective effects of environmental change emerge because organisms differ in their response to
 51 environmental change. For example, larger organisms and predators are often more negatively
 52 affected than smaller organisms at lower trophic levels (*Hines et al.*, 2015; *Sheridan and Bickford*,
 53 2011; *Srivastava and Vellend*, 2005; *Voigt et al.*, 2007). Using realistic extinction scenarios, exper-
 54 iments found contrasting effects of non-random shifts in biodiversity on ecosystem functioning
 55 (e.g. *Cárdenas et al.*, 2017; *Jonsson et al.*, 2002; *Melguizo-Ruiz et al.*, 2020; *Oliveira et al.*, 2019;
 56 *Smith and Knapp*, 2003; *Zavaleta and Hulvey*, 2004). In addition, several variables that are not
 57 directly related to biodiversity control ecosystem functions (e.g. physiological rates (*Dib et al.*,
 58 2020; *Thakur et al.*, 2018) and alterations of physical and chemical conditions (*De Laender et al.*,
 59 2016; *Giling et al.*, 2019)). When environmental change affects these mechanisms, teasing out the
 60 relative importance of biodiversity-mediated effects is complicated even more. Given the number
 61 of different potential mechanisms, quantifying the extent to which shifts in biodiversity underpin
 62 the effect of environmental change on ecosystem functioning under real-world scenarios of global
 63 change is a key challenge for ecology (*De Laender et al.*, 2016; *Duffy et al.*, 2017; *Eisenhauer et al.*,
 64 2019b; *Srivastava and Vellend*, 2005; *van der Plas*, 2019; *Wardle*, 2016). Incorporating the impacts
 65 of environmental change drivers into BEF studies and meta-analyses is an important step forward
 66 to address such questions (*De Laender et al.*, 2016; *Eisenhauer et al.*, 2019b).

67 The vast majority of BEF experiments has focused on plant richness and ecosystem functions
 68 such as biomass production (*van der Plas*, 2019). However, litter decomposition has a tremendous
 69 importance in ecosystems and biogeochemical cycles (*Follstad Shah et al.*, 2017). Small changes in
 70 the rate of this process can have important consequences for the overall carbon balance. Indeed,
 71 increases in decomposition rates could have positive feedback effects on climate warming by
 72 enhancing C losses (*Kirschbaum*, 2000). The diversity of decomposers (invertebrates and micro-
 73 organisms that fragment and decompose organic matter in both aquatic and terrestrial systems)
 74 is crucial for litter decomposition (*Eisenhauer et al.*, 2012; *García-Palacios et al.*, 2013; *Gessner*
 75 *et al.*, 2010; *Handa et al.*, 2014; *Hättenschwiler et al.*, 2005) and for other ecosystem functions as
 76 well (*Eisenhauer et al.*, 2019a; *Lefcheck et al.*, 2015; *Schuldt et al.*, 2018). Despite the importance
 77 of decomposers, BEF experiments focusing on litter decomposition more often addressed the
 78 influence of plant litter diversity than of decomposers (*Gessner et al.*, 2010; *Tonin et al.*, 2018). In
 79 a meta-analysis, decomposer diversity had a greater effect on decomposition than the diversity
 80 of plant litter (*Srivastava et al.*, 2009), although also weak and neutral effects have been reported
 81 (*van der Plas*, 2019). Facilitation and complementarity through niche partitioning are primary
 82 mechanisms underlying the positive relationship between decomposer diversity and decomposition
 83 (*Gessner et al.*, 2010; *Hättenschwiler et al.*, 2005; *Tonin et al.*, 2018). Experiments conducted
 84 in natural conditions and reflecting realistic extinction scenarios are still relatively scarce, and
 85 demonstrate contrasting effects of non-random shifts in decomposer diversity on decomposition
 86 (*Cárdenas et al.*, 2017; *Jonsson et al.*, 2002; *Melguizo-Ruiz et al.*, 2020). The need to quantify
 87 environmental change effects on decomposer diversity, along with potential knock-on effects on
 88 litter decomposition, is therefore particularly pressing.

89 There is a variety of environmental change drivers, and different types of drivers may have
 90 diverse effects on biodiversity and ecosystem functions (*De Laender et al.*, 2016; *Dib et al.*, 2020).
 91 We postulate that there are two main categories of environmental change: stressors and resource

92 shifts. While stressors cannot be consumed, and act as conditions that alter growth rates (e.g., tem-
 93 perature, drought, chemical stressors), resources are by definition consumed (e.g., CO₂ or mineral
 94 nutrients), which has important implications for how they should enter theory (*Chase and Leibold,*
 95 *2003; De Laender, 2018*). Chemical stressors and nutrient enrichment are important case studies
 96 of environmental stressors and resource enrichment, because their presence is increasing rapidly
 97 (*Bernhardt et al., 2017*) and they are projected to have severe effects on biodiversity (*Mazor et al.,*
 98 *2018*). They are also of particular relevance for decomposer communities. Chemical stressors such
 99 as metals and pesticides decrease the diversity, abundance, growth and activity of decomposers
 100 across terrestrial and aquatic systems (e.g., *Hogsden and Harding, 2012; Pelosi et al., 2014; Schäfer,*
 101 *2019*). In contrast, nutrient enrichment can have positive impacts on the abundance and physio-
 102 logical rates of decomposer organisms by reducing resource limitations (*Treseder, 2008*), but at
 103 the same time decrease decomposer diversity (*Lecerf and Chauvet, 2008; Woodward et al., 2012*).
 104 Across ecosystems, stressors and nutrients can exert opposite impacts on litter decomposition
 105 rates, with decreases in response to chemical stressors but increases following nutrient enrichment
 106 (*Ferreira et al., 2015, 2016*). In addition, decomposition involves both microorganisms and inverte-
 107 brates (*Bardgett and van der Putten, 2014; Gessner et al., 2010; Hättenschwiler et al., 2005*) that
 108 may respond differently to stressors and nutrients with a higher sensitivity of invertebrates than
 109 microorganisms (*Peters et al., 2013; Siebert et al., 2019*). Although many published case studies
 110 report shifts in decomposer diversity and in rates of litter decomposition at sites impacted by
 111 stressors and nutrients, biodiversity-mediated effects have not yet been quantified across systems.

112 Here we addressed the question if the effects of stressors and nutrient enrichment on decom-
 113 poser diversity and abundance explain the response of litter decomposition to these two types
 114 of pervasive environmental change drivers (*Figure 1*). We synthesized 69 published case studies
 115 reporting the impact of stressors (metals, pesticides) and nutrients (nitrogen or phosphorous
 116 additions) on litter decomposition and on decomposer diversity (taxa richness, Shannon diversity,
 117 evenness) or abundance (density, biomass) at sites differing in stressor or nutrient levels. Our
 118 comprehensive global dataset of 660 observations encompasses studies across taxonomic groups
 119 (animal (soil micro-, meso- and macrofauna, stream macroinvertebrates) and microbial (fungi and
 120 bacteria) decomposers), ecosystems (aquatic and terrestrial), and study types (experimental and
 121 observational) (*Figure 2*). We quantified the effect size of environmental change on decomposer
 122 diversity or abundance and on litter decomposition within studies using correlation coefficients
 123 between stressor or nutrient levels and decomposer diversity, abundance, and litter decomposition.
 124 We also characterized stressor and nutrient intensities, and standardized their levels in water, soil,
 125 or sediment using environmental quality criteria issued by environmental authorities (e.g. ECHA,
 126 USEPA, UKTAG). Using meta-analysis and structural equation modelling (SEM), we first compared
 127 the overall effects of stressors and nutrients on decomposers and decomposition across systems
 128 and studies (first meta-analysis), and second, addressed to what extent changes in decomposer
 129 diversity and abundance mediate the impacts of these two contrasting drivers of environmental
 130 change on decomposition (second meta-analysis and SEM). Third, we explored the effects of three
 131 main moderators on decomposers diversity, abundance, and decomposition responses, as found
 132 in the second meta-analysis: stressor or nutrient intensity, taxonomic group (animal vs. microbes)
 133 and study type (experimental vs. observational studies).

134 We expected that chemical stressors and nutrients would have contrasting effects on de-
 135 composer diversity and abundance, and on litter decomposition across ecosystems and studies
 136 (*Figure 1*). We hypothesized that chemical stressors generally decrease decomposer diversity,
 137 abundance (*Hogsden and Harding, 2012; Petrin et al., 2008*), and litter decomposition rates (*Fer-*
 138 *reira et al., 2016; Peters et al., 2013*), and that nutrients generally decrease decomposer diversity
 139 (*Lecerf and Chauvet, 2008; Woodward et al., 2012*) but increase decomposer abundance and litter
 140 decomposition rates (based on physiological effects and decreasing resource limitations (*Bergfur*
 141 *et al., 2007; Ferreira et al., 2015; Treseder, 2008; Woodward et al., 2012*)). We further hypothesized
 142 that litter decomposition responses to environmental change depend on changes in decomposer di-




Figure 1. Schematic representation of the structural hypotheses tested in this study. Green arrows depict expected positive effects, red arrows represent negative effects. Stressors and nutrients are hypothesized to decrease decomposer diversity. The response of decomposers diversity to environmental change drivers determines the response of decomposition (Srivastava et al., 2009). Nutrients are hypothesized to increase decomposer abundance. Stressors and nutrients can affect litter decomposition independent of changes in decomposer diversity and abundance, especially through changes in physiological activity (De Laender et al., 2016, Giling et al., 2019).

143 versity and abundance, and expected an overall positive relationship independent of environmental
 144 change intensity (Srivastava et al., 2009).

145 Results

146 Description of the data and overall patterns


147 The final dataset contained 69 (case) studies from 59 publications, representing 660 observations.
 148 Data were mostly from Europe (44 ; 443 (studies; observations)) and North and South America (19;
 149 168), while Asia (2; 9) and Oceania (4, 40) were less well represented (Figure 2.A). The studies covered
 150 aquatic (55; 388) and terrestrial systems (14; 272) (Figure 2.C), and used observational (43; 336) or
 151 experimental approaches (26; 324). Studies reported abundance (66; 463) or diversity responses
 152 (48; 197) (Figure 2.B) of soil and benthic invertebrates (48; 509) and microbes (fungi and bacteria)
 153 associated with litter materials (36; 151) (Figure 2.C). Chemical stressors were mostly metals (13;
 154 257) and pesticides (12; 66) associated with industrial activities, accidental spills, and agricultural
 155 practices. Nutrient enrichment studies addressed fertilization by various N and/or P forms (26;
 156 175), and eutrophication due to agricultural runoffs (10; 59) or wastewater effluents (4; 44). There
 157 was no study reporting nutrient enrichment impacts on soil decomposer diversity in the dataset.
 158 Funnel plots and intercepts of Egger's regression showed evidence for positive publication bias in
 159 nutrient enrichment studies reporting decomposer abundance (Appendix 2-Figure 2, -Table 1). No
 160 publication bias was detected in the other datasets.

161 We found largely contrasting effects of stressors and nutrients on each of the three response
 162 variables in a first-level meta-analysis comparing the overall effects of the two drivers of environ-
 163 mental change (Figure 3, Appendix 2-Table 2). Chemical stressors overall decreased decomposer
 164 diversity, abundance and litter decomposition across studies (Figure 3). Nutrient enrichment
 165 tended to decrease decomposer diversity but to increase abundance, and decomposition, although
 166 these trends were not significant as indicated by confidence intervals of the grand mean effects
 167 overlapping with zero (Figure 3).


168 Biodiversity-mediated effects of stressors and nutrients on litter decomposition

169 The responses of decomposition and of decomposer diversity and abundance to chemical stressors
 170 were correlated: decreases in decomposition were associated with decreases in decomposer
 171 diversity and abundance (Figure 4 upper panels). We did not find such a relationship for nutrients.


A



B



C



Legend: Abundance (grey), Biomass (light grey), Diversity indices (dark grey), Taxa richness (medium grey), Aquatic (white), Terrestrial (black)

Figure 2. Description of the data used in the present meta-analysis. A: countries represented and corresponding number of observations, B: decomposer diversity and abundance metrics covered, and C: ecosystem types and decomposer taxonomic groups (animals: soil micro-, meso-, macro-fauna, stream macroinvertebrates; and microbial decomposers: fungi and bacteria) represented.




Figure 3. Grand mean effect sizes of chemical stressors and nutrient enrichment on decomposer diversity (taxa richness and diversity indices), abundance (density and biomass), and litter decomposition. Effect sizes are z-transformed correlation coefficients. Error bars show 95% confidence intervals. Numbers in parentheses indicate number of studies and observations, respectively. Symbols show the significance level for the comparison between mean effect size and zero (*** $P < 0.001$; * $P < 0.05$). For full model results, see [Appendix 2-Table 2](#).




Figure 4. Relationship between the responses of decomposition and decomposer diversity and abundance to chemical stressors and nutrient enrichment. Variables are effect sizes (z-transformed correlation coefficients) of stressors or nutrients on litter decomposition and on animal and microbial decomposer diversity (left panels) or abundance/biomass (right panels). Gray symbols are individual observations of effect sizes; Colored symbols indicate the mean effect size on biodiversity or abundance across effect sizes on litter decomposition. Darker colors represent a higher standardized level of environmental change. Lines represent meta-regressions between effect sizes for decomposition and decomposers, where solid lines are statistically significant ($P < 0.05$), dashed lines are non-significant ($P > 0.05$), and thin lines depict the regression's confidence interval. QM and P represent the model heterogeneity P -value of the meta-regressions, respectively, with sample size (number of studies; number of observations).

172 Instead, a range of positive and negative responses of decomposer diversity, abundance, and
 173 decomposition to nutrients were found, without significant associations between them (**Figure 4**
 174 lower panels). In addition, when decomposer diversity and abundance responses to nutrients were
 175 close to zero, there was a wide range of decomposition responses (intercepts from **Figure 4** lower
 176 panels).

177 According to our overarching hypothesis, the SEM indicated that the effects of stressors on litter
 178 decomposition were mediated by shifts in decomposer diversity and abundance. Including the
 179 direct paths from decomposer diversity or abundance to litter decomposition improved both the
 180 models according to mediation tests and AIC comparisons (**Figure 5**). In addition, the path coeffi-
 181 cients from diversity and abundance to the decomposition response to stressors had (standardized)
 182 values higher than 0.1 (**Figure 5**) and were statistically different from zero (**Appendix 2-Table 3**).
 183 However, in contrast to chemical stressors, the SEM did not support biodiversity-mediated effects
 184 of nutrient enrichment on litter decomposition. While the mediation test and AIC indicated that
 185 the decomposer diversity-mediated path improved the model (**Figure 5**), the path coefficient was
 186 not significantly different from 0 (**Appendix 2-Table 3**). The decomposer abundance-mediated path
 187 of nutrients was not supported by the data: an SEM without the direct path from decomposer

188 abundance to decomposition could not be rejected based on the mediation test (*Figure 5*), and
 189 including this path did not improve the model according to the AIC comparison. Besides, we found
 190 publication bias in this dataset (*Appendix 2-Figure 2, -Table 1*), and model check indicated that the
 191 residuals of the nutrients-abundance model were non-independent from the fitted values. Thus,
 192 the results from this model are reported here for comparison purposes only.

193 The magnitude of the biodiversity-mediated effects of chemical stressors on decomposition was
 194 stronger than that of the direct effects of stressor intensity on decomposition. The indirect effect of
 195 stressors on decomposition mediated by diversity (i.e. mathematical product of the standardized
 196 paths from stressor intensity to decomposer diversity and from diversity to decomposition *Figure 5*)
 197 was higher than the direct effect of stressors on decomposition, while the abundance-mediated
 198 effect of stressors was negligible (*Figure 5*). In the case of nutrient enrichment, however, decomposi-
 199 tion responses were not explained by shifts in decomposer diversity and abundance, and the direct
 200 effects of nutrient intensity dominated the total effect (*Figure 5*). Finally, between-model compar-
 201 isons (based on unstandardized path coefficients (*Grace, 2006*)) revealed that decomposer diversity
 202 was a stronger driver of decomposition response to stressors than decomposer abundance (unstan-
 203 dardized paths were 0.42 and 0.24 respectively for diversity and abundance, *Appendix 2-Table 3*).

204 Sensitivity analyses revealed that the results were robust to the inclusion of approximated
 205 standard deviations (*Appendix 3- -Table 1, -Table 2*), and extreme values of effect sizes (*Appendix 3-*
 206 *Table 3, -Table 4*). We found partially different results when using log-response ratios as effect
 207 sizes (*Appendix 3-Table 5, -Table 6*), due to lower sample sizes and emergence of extreme values in
 208 these datasets. In addition, the log-response ratio is probably sensitive to the various metrics of
 209 biodiversity, abundance, and decomposition covered by the individual studies that we included,
 210 while correlation coefficients better accommodate such discrepancies (*Koricheva et al., 2013*).

211 **Response of animal and microbial decomposers and decomposition to stressor and** 212 **nutrient intensity**

213 Despite the overall negative effects of stressors on decomposition, negative responses in decompo-
 214 sition were not associated with higher stressor intensity (*Figure 5, Figure 6*). This result held for two
 215 complementary approaches: multivariate SEM (*Figure 5*) that relied on data resampling to account
 216 for replicated values of decomposition matching several decomposer responses (e.g. for different
 217 taxa in the same litterbag), and meta-regressions (*Figure 6*) where data resampling was not neces-
 218 sary (see Methods). There was mixed support for a stressor intensity effect on decomposer diversity
 219 across the two approaches: decomposer diversity responses decreased with stressor intensity
 220 according to the SEM (*Figure 5*), but this trend was not significant according to the second level
 221 meta-analysis (*Figure 6*). Similar slopes were obtained both with the SEM relying on data resampling
 222 (the slope of the relationship was -0.10 ± 0.04 , *Appendix 2-Table 2*) and with the meta-regression
 223 (the slope was -0.05 ± 0.03). The differences between the two approaches can be explained by
 224 the different data included. Decomposer abundance responses were not associated to stressor
 225 intensity in both the SEM and meta-regression approaches (*Figure 5, Figure 6*). We found different
 226 patterns for nutrient enrichment, where decomposition responses decreased with nutrient intensity
 227 (*Figure 5, Figure 6*), from positive effects at low intensity to negative effects at higher intensity (*Fig-*
 228 *ure 6*). A similar pattern was observed for decomposer diversity, where responses decreased with
 229 nutrient intensity from positive to neutral to negative responses at high nutrient levels (*Figure 6*).
 230 Nutrient intensity, however, did not explain the responses of decomposer abundance (*Figure 5,*
 231 *Figure 6*), and both positive and negative responses were found at high nutrient levels.

232 The meta-analysis further revealed clear discrepancies between the response of animal and
 233 microbial (fungi and bacteria) decomposers to stressors and nutrients. Animal decomposers
 234 responded more strongly to chemical stressors than microbial decomposers. The mean effects
 235 of chemical stressors on animal decomposer diversity and abundance were more negative than
 236 that on microbial decomposers, confirmed by Wald type tests of the second-level meta-analyses
 237 (*Figure 7* upper panels, *Appendix 2-Table 4*). Animal decomposers overall decreased in diversity but




Figure 5. Decomposer diversity and abundance explained litter decomposition response to chemical stressors but not to nutrient enrichment. Structural equation models investigating decomposer diversity- or abundance-mediated effects of chemical stressors and nutrient enrichment on litter decomposition across 69 studies. Arrows represent relationships between stressor or nutrient intensity levels, and effect sizes of stressors or nutrients on litter decomposition and on decomposer diversity (taxa richness, Shannon diversity, or evenness: left panels) or abundance and biomass (right panels). Values along the arrows are standardized path coefficients. Green, red, and gray arrows indicate positive, negative, and non-significant relationships, respectively. Curved arrows depict the indirect effects of stressors or nutrients on decomposition as mediated by diversity or abundance. Arrow widths are scaled relative to the magnitude of standardized path coefficients. C statistic, P -value ($P < 0.05$ indicate poor model fit), and sample sizes (number of studies; number of observations). Results of mediation tests: comparison with models omitting the path from diversity or abundance to decomposition ($\Delta AIC < -2$ indicates that reduced models were not consistent with the data).




Figure 6. Decomposer and decomposition responses to the intensity levels of chemical stressors and nutrient enrichment. Values are effect sizes (z-transformed correlation coefficients). Stressor or nutrient intensity represents the standardized level of environmental change in the treatment with the highest level (values <0: observed level below quality criteria considered to be safe for the environment; values > 0: observed level above quality criteria). Point size is proportional to the inverse of the variance in effect size. Lines are the slopes and 95% confidence intervals from bivariate meta-regressions, with associated QM statistics, *P*-value and sample size (number of studies; number of observations).




Figure 7. Moderator effects on decomposer diversity and abundance and on decomposition responses to chemical stressors and nutrient enrichment. Responses of decomposer diversity (taxa richness and diversity indices) and abundance (densities and biomass) to stressors and nutrients according to the taxonomic group (animals and microbes) and study type (Expe. = experimental; Obs. = observational studies). Values are mean effect sizes (z-transformed correlation coefficients) and 95% confidence intervals derived from meta-analytic models. Sample sizes are reported for each moderator: (number of studies; number of observations).

238 increased in abundance in response to nutrient enrichment (*Figure 7*, lower panels). On the other
 239 hand, the mean effects of nutrients on microbial decomposer diversity and abundance had lower
 240 magnitudes compared to animals (*Appendix 2-Table 4*), with confidence intervals overlapping with
 241 zero (*Figure 7* lower left panel). Finally, there was no clear difference between observational and
 242 experimental studies (*Figure 7*, *Appendix 2-Table 4*), and between biodiversity responses in terms
 243 of taxa richness or of diversity indices (*Appendix 2-Table 4*).

244 Discussion

245 The present synthesis brings new insights into how changes in decomposer biodiversity induced
 246 by two pervasive drivers of environmental change ultimately affect decomposition. We find con-
 247 comitant changes in biodiversity and decomposition under the influence of chemical stressors
 248 but not nutrient enrichment, highlighting that real-world patterns relating shifts in biodiversity
 249 and ecosystem functioning depend on the type of environmental change. In fact, we observed
 250 significant correlations between effects on biodiversity and ecosystem function in a scenario where
 251 chemical stressors caused a significant decline in biodiversity. In contrast, in cases where nutri-
 252 ent enrichment caused variable responses in biodiversity, relationships between biodiversity and
 253 ecosystem function responses were weaker. It remains an understudied but important question if
 254 results of controlled BEF experiments are applicable to non-random changes in biodiversity caused
 255 by human activities (e.g., *De Laender et al., 2016; Duffy et al., 2017; Eisenhauer et al., 2019b; Sri-*

256 *vastava and Vellend, 2005; van der Plas, 2019; Wardle, 2016*). The present results provide strong
 257 empirical evidence for significant real-world BEF relationships when human activities decrease
 258 biodiversity.

259 **Biodiversity-mediated effects of chemical stressors on decomposition**

260 Chemical stressors caused consistent reductions in decomposer diversity and abundance as well as
 261 in litter decomposition rates, in line with several previous case studies (*Beketov et al., 2013; Malaj*
 262 *et al., 2014*) and meta-analyses (*Ferreira et al., 2016; Peters et al., 2013*). Adding to the previous
 263 knowledge, the present meta-analysis shows that changes in decomposer diversity and abundance
 264 explained the decomposition response to stressors, providing evidence for the expectation that
 265 shifts in biodiversity mediate the impact of chemical stressors on decomposition. We acknowledge
 266 that despite the SEM analysis, the approach conducted here remains correlative. However, our
 267 study builds on a body of experimental and observational evidence that already demonstrated that
 268 more diverse and abundant decomposer communities support higher decomposition rates, albeit
 269 not under the influence of environmental change (e.g. *García-Palacios et al., 2013; Handa et al.,*
 270 *2014*).

271 We especially complement a previous meta-analysis showing the importance of decomposer
 272 diversity for decomposition across experiments manipulating the richness of invertebrate and
 273 microbial decomposer communities (*Srivastava et al., 2009*). We extend on this and show that
 274 non-random biodiversity losses induced by stressors are closely associated with decreases in
 275 decomposition across a wide range of studies. A recent review pointed out that in naturally-
 276 assembled terrestrial communities, studies more often found neutral and to a lesser extent positive
 277 relationships between decomposer diversity and decomposition (*van der Plas, 2019*). In that
 278 review, communities were not influenced by environmental change drivers, and the vote counting
 279 approach used is sensitive to the statistical power of individual studies and could have increased
 280 the probability of finding non-significant relationships (*Koricheva et al., 2013*). In line with our
 281 findings, an experiment mimicking the sequence in which freshwater invertebrate decomposers
 282 are lost after disturbances showed that decreasing non-randomly the number of species decreased
 283 decomposition rates (*Jonsson et al., 2002*).

284 Biodiversity-ecosystem function experiments manipulating biodiversity directly are key to un-
 285 derstand the mechanisms involved in this relationship (*Eisenhauer et al., 2016*), especially because
 286 they control for the effects of environmental heterogeneity or abundance. However, in real-world
 287 scenarios, environmental change drivers affect both biodiversity and abundance simultaneously. As
 288 demonstrated here, this is especially the case for stressors that decrease decomposer diversity and
 289 abundance (*Hogsden and Harding, 2012*). The abundance or biomass of different decomposers
 290 is of critical importance for decomposition (e.g. *Bergfur et al., 2007; Ebeling et al., 2014; Manning*
 291 *and Cutler, 2018*). Even at constant richness and community composition, strong decreases in
 292 abundance can have important impacts on ecosystem functioning (*Spaak et al., 2017*; but see
 293 *Dainese et al., 2019*). It is beyond the scope of the present meta-analysis to disentangle the effects
 294 of biodiversity from the effects of abundance, and we found that both contributed to explain shifts
 295 in decomposition in separate analyses. It is interesting to note that the few cases where negative
 296 effect sizes of stressors on biodiversity were associated with positive effect sizes on decomposition
 297 were also cases where decomposer abundance was positively associated with stressors (*Figure 4*).
 298 Although we cannot specifically test this with the present data, it seems that in those particular
 299 cases, increases in decomposer abundance counteracted the negative effects of decreases in
 300 decomposer diversity (*Lucisne et al., 2015*). Those results could therefore be in line with the mass-
 301 ratio hypothesis (*Grime, 1998; Smith and Knapp, 2003*). Indeed, an exclusion experiment showed
 302 that dominant, small, detritivores can compensate reductions in litter decomposition caused by
 303 the removal of large detritivores (*Cárdenas et al., 2017*). These concomitant shifts in both diversity
 304 and abundance further have important implications for our estimates of diversity responses, as
 305 studies mostly reported richness to estimate decomposer diversity, but rarely corrected for the

306 sampling effort (*Gotelli and Colwell, 2001*). This means that lower abundances rather than a lower
 307 number of species *per se* might have directly caused some of the negative effects on biodiversity
 308 reported here (*Chase and Knight, 2013*). This common caveat in meta-analysis approaches that
 309 rely on how individual studies report biodiversity, also applies to the present study, and reinforces
 310 the importance of reporting raw data in future studies of the impacts of chemical stressors on
 311 biodiversity and ecosystem functioning.

312 The effects of changes in decomposer diversity and abundance on decomposition found in
 313 the present study might also have channeled changes in community and food-web structure
 314 not captured by our biodiversity metrics. Changes in keystone species (*Hättenschwiler et al.,*
 315 *2005*), functional diversity (*Cadotte et al., 2011; Dangles et al., 2012; Heemsbergen et al., 2004*),
 316 vertical diversity (*Gessner et al., 2010; Melguizo-Ruiz et al., 2020; Wang and Brose, 2018; Zhao et al.,*
 317 *2019*), or dominance patterns (*Dangles and Malmqvist, 2004*) might have shifted concomitantly to
 318 taxonomic diversity and abundance. Moreover, these different components of diversity might act
 319 at different timings of decomposition (*Oliveira et al., 2019*). Unfortunately, studies rarely reported
 320 such measurements together with decomposition. For example in our dataset, only 7 studies
 321 reported evenness. Future studies need to explore shifts in decomposer community composition in
 322 more detail to better understand what particular aspect of biodiversity is responsible for changes
 323 in decomposition rates (*Giling et al., 2019; Hättenschwiler et al., 2005*). In particular, few of the
 324 included studies reported comparable functional groups allowing to address the effect of functional
 325 diversity across the multiple systems and taxonomic groups addressed by the present analysis.
 326 Future synthesis work could specifically address the effect of functional diversity, by focusing on a
 327 given system type. Indeed, there is ample evidence that shifts in functional diversity are crucial for
 328 decomposition (*Heemsbergen et al., 2004*), and that facilitative interactions occur primarily between
 329 decomposers of contrasting body size (*Dangles et al., 2012; Tonin et al., 2018*). This is especially the
 330 case for interactions between animal and microbial decomposers, where fragmentation of litter by
 331 detritivores facilitates access for microbial decomposers (*Eisenhauer et al., 2010; Hättenschwiler*
 332 *et al., 2005; Yang et al., 2012*).

333 Here, we found that invertebrates were more affected by chemical stressors than microbes,
 334 across aquatic and terrestrial ecosystems. Invertebrate decomposers are particularly sensitive to
 335 the impacts of metals and pesticides (*Hogsden and Harding, 2012; Pelosi et al., 2014; Peters et al.,*
 336 *2013; Schäfer, 2019*). Microbial decomposers are known to be sensitive to metals (*Giller et al.,*
 337 *2009*) and pesticides as well (*DeLorenzo et al., 2001*). Nevertheless, our result is consistent with the
 338 general expectation that larger organisms are more sensitive to environmental change due to longer
 339 generation time, higher energetic demands and lower population densities (*Hines et al., 2015;*
 340 *Sheridan and Bickford, 2011; Woodward et al., 2005; Yvon-Durocher et al., 2011*). These different
 341 sensitivities between groups of decomposers could imply that the biodiversity-mediated effects of
 342 stressors on decomposition are more strongly linked to shifts in invertebrates than microbes, as
 343 reported in a previous review (*Peters et al., 2013*). However, in another meta-analysis focusing on
 344 microbial-driven decomposition rates, changes in fungal biomass and richness explained shifts in
 345 decomposition under the impacts of chemical stressors, but also of nutrient enrichment (*Lecerf*
 346 *and Chauvet, 2008*).

347 **Nutrient-induced changes in decomposition were not related to shifts in decom-** 348 **poser diversity**

349 The impacts of nutrient enrichment on litter decomposition and decomposer diversity were different
 350 from those caused by stressors, confirming our expectations. These different biodiversity and
 351 function responses led to different emergent relationships between decomposer diversity and
 352 decomposition compared to stressors. We found that nutrients had a variety of effects ranging from
 353 positive to negative depending on the taxonomic group (*Figure 7*) and nutrient intensity (*Figure 6*),
 354 and resulting in neutral overall mean effects (*Figure 3*). Previous syntheses also found positive
 355 (*Ferreira et al., 2015*) as well as inconsistent (*Knorr et al., 2005*) responses of decomposition rates

356 to nutrient enrichment in streams. The relatively small mean effect of nutrient enrichment on
357 decomposition in the present meta-analysis could be explained by the use of correlation as an effect
358 size, which does not capture potentially non-monotonic responses of decomposition to nutrients
359 (*Woodward et al., 2012*). However, we noted that most of the studies included in the present
360 meta-analysis did not individually span nutrient gradients sufficiently large to capture this potential
361 non-monotonous response. Taken together, the studies show positive effects on decomposition
362 at low nutrient intensities that shifted towards neutral to negative effects at higher intensities
363 (*Figure 6*), which is consistent with previous findings (*Ferreira et al., 2015; Woodward et al., 2012*).
364 Low nutrient intensities might have enhanced microbial activity and biomass by alleviating resource
365 limitation, resulting in enhanced decomposition. At higher intensities, however, negative impacts
366 on invertebrates might have decreased decomposition rates (*Peters et al., 2013; Woodward et al.,*
367 *2012*).

368 These nutrient intensity patterns contrasted with the results for chemical stressors. The overall
369 negative effects of stressors (*Figure 1*) on decomposition were not explained by stressor intensity
370 levels (*Figure 6*), and there was mixed support for a stressor intensity effect on decomposer diversity
371 based on two complementary data analysis approaches (SEM based on data resampling (*Figure 5*)
372 vs. second level meta-analysis *Figure 6*). Thus, negative responses to chemical stressors happened
373 across the range of stressor intensity. Such contrasting patterns between stressor and nutrient
374 intensity effects may reflect the greater number of stressor types (different metals, pesticides,
375 mixtures) covered by individual studies compared to the limited number of nutrients. In addition,
376 due to the higher variability of stressor types, we relied on more variable sources to standardize
377 stressor levels compared to nutrients in the diversity dataset (Methods, *Appendix 1-Table 1*). With
378 the data at hand, it was not possible to test the influence of the environmental quality criteria
379 used to standardize stressor and nutrient levels, because such an effect would be confounded with
380 stressor or nutrient types. The datasets were all dominated by environmental quality criteria based
381 on similar methodologies (for 75 to 100% of observations, see Methods). However, future studies
382 focusing on stressor intensity effects across ecosystems would greatly benefit from coordinated
383 efforts to derive quality criteria encompassing the vast and rapidly increasing number of chemical
384 stressors (*Wang et al., 2020*).

385 Contrary to our expectation, nutrient-induced shifts in decomposer diversity and abundance
386 were not associated with shifts in decomposition rates across studies. We found that increasing
387 nutrient intensity decreased the effects on decomposition and on decomposer diversity, but not
388 on decomposer abundance. Statistically controlling for the effect of nutrient intensity with SEM
389 indicated no residual association between shifts in decomposer diversity or abundance and in
390 decomposition rates, i.e. a non-significant BEF relationship. Changes in microbial abundance
391 in response to nitrogen deposition explained the responses of different ecosystem functions in
392 terrestrial systems in previous meta-analyses (*García-Palacios et al., 2015; Treseder, 2008*). Here
393 we show that this pattern cannot be generalized across aquatic and terrestrial systems and across
394 animal and microbial decomposers. Contrary to stressors, when the diversity and abundance of
395 animal and microbial decomposers were not affected by nutrients, we observed large positive
396 and negative shifts in decomposition (intercepts of *Figure 4*), that were explained by nutrient
397 intensity (*Figure 4*: negative effects on decomposition at invariant biodiversity are associated
398 with high intensities and positive effects with lower intensities). Together, these results show
399 that nutrient-induced shifts in decomposer diversity were not as strong drivers of decomposition
400 changes as stressor-induced biodiversity shifts. These differences may be partly due to the different
401 mechanisms underlying the effects of stressors and nutrients. Based on previous studies, we
402 speculate that our results are due to the complex responses of animal and microbial decomposers
403 at different nutrient intensities (*Ferreira et al., 2015; Lecerf and Chauvet, 2008; Treseder, 2008;*
404 *Woodward et al., 2012*).

405 Animal decomposers showed a stronger response to nutrients than microbes. Invertebrate
406 decomposers overall decreased in diversity, but they increased in abundance under nutrient

407 enrichment. These results could reflect a loss of sensitive taxa to the benefit of tolerant taxa that
 408 were able to use additional resources and would then increase in density (*Bergfur et al., 2007*).
 409 Overall, microbial decomposers responded little to nutrient enrichment, probably reflecting a
 410 mixture of positive and negative effects that nutrients can have on microbial growth (*Lecerf and*
 411 *Chauvet, 2008; Treseder, 2008*), as well as on different microbial taxa. Indeed, nutrients can alleviate
 412 resource limitations at low intensities, but can also exert toxic effects at high intensities. The initial
 413 levels of nutrients thus condition subsequent responses in decomposers and decomposition to
 414 nutrient enrichment (*Ferreira et al., 2015; Knorr et al., 2005*). Furthermore, at high intensities,
 415 nutrients can be associated with other chemical stressors (e.g. pesticides in agricultural runoffs)
 416 (*Ferreira et al., 2015; Woodward et al., 2012*). The influence of interactive effects of stressors and
 417 nutrients was impossible to quantify with the data at hand, given that only a few experiments
 418 assessed the effects of both drivers independently, but many observational studies may have
 419 been confounded by such joint effects. Chemical stressors and nutrients are often co-occurring in
 420 e.g. agricultural landscapes, and the consequences of such combinations are still poorly understood.
 421 Furthermore, stressor and nutrient effects might be modulated by climatic and other environmental
 422 conditions, and studies on interaction effects are scarce (*Rillig et al., 2019; Thakur et al., 2018*).
 423 Finally, although our comparison of stressors versus resources allowed us to test a clear concept,
 424 any kind of grouping in ecological studies may mask some of the variation within the categories
 425 and future studies may be interested in different categories. As data availability improves, future
 426 work could include different environmental change drivers. This would also allow to test additional
 427 groupings of drivers and ecological concepts unifying stressors and resources (*De Laender, 2018;*
 428 *Harley et al., 2017*).

429 Conclusions

430 In conclusion, this study brings new insights into the real-world patterns relating ecosystem function
 431 to non-random changes in biodiversity induced by environmental change. We found that the conse-
 432 quences of changes in biodiversity for ecosystem functioning depend on the type of environmental
 433 change. Real-world scenarios do not necessarily involve concomitant changes in both biodiversity
 434 and function across terrestrial and aquatic systems. We further found that with the environmental
 435 quality criteria used in risk assessment, there were already significant positive and negative effects
 436 on decomposers and decomposition (*Figure 6*), highlighting the need to better incorporate biodi-
 437 versity and ecosystem function into ecological risk assessment programs (*De Laender and Janssen,*
 438 *2013*). Finally, we report overall negative effects of chemical stressors on biodiversity and ecosys-
 439 tem functioning across terrestrial and aquatic ecosystems that reinforce recent calls to consider
 440 chemical stressors as important global change drivers and address their impacts on biodiversity
 441 and ecosystems (*Bernhardt et al., 2017; Mazor et al., 2018; Steffen et al., 2015*). Positive real-world
 442 BEF relationships may be particularly significant in cases where environmental changes decrease
 443 biodiversity, such as in the case of chemical stressors. Such information are crucial if we are to
 444 design policy and conservation strategies able to reconcile human development with biodiversity
 445 conservation.

446 Methods

447 Data collection

448 We searched the Web of Science for studies that addressed the impact of environmental drivers and
 449 recorded decomposer community responses and litter decomposition rates. The search strategy
 450 is fully reported in Supplementary Methods (*Appendix 1*). The search retrieved 2536 references.
 451 Abstracts and titles were screened to identify a final set of 61 records that met our inclusion criteria
 452 (PRISMA plot, *Appendix 1-Figure 1*, and list of included references (*Appendix 4*)). To be included in
 453 the meta-analysis, studies had to:

- 454 • Report litter decomposition (rates, mass loss, proportion of mass remaining) and the diversity,

- 455 abundance, or biomass of decomposers at sites differing in chemical stressor or nutrient
 456 levels.
- 457 • Focus on naturally-assembled communities subjected to the impact of chemical stressors or
 458 nutrient enrichment. Studies that manipulated decomposer diversity directly were not consid-
 459 ered to only focus on non-random biodiversity change scenarios. We included mesocosm
 460 studies only when they used field-sampled communities and left time for the community
 461 to reach an equilibrium in mesocosms in order to reflect real-world conditions as much as
 462 possible.
 - 463 • Report the response of animal (benthic macroinvertebrates, or soil micro, meso or macro-
 464 fauna) or microbial decomposers (bacteria or fungi from decomposing leaves or in surround-
 465 ing water or soil samples).
 - 466 • Report decomposer abundance (density or biomass), or decomposer diversity (taxa richness,
 467 Shannon diversity, evenness).

468 When a reference reported different environmental change drivers or geographical areas with
 469 a specific reference site for each case, we considered these as individual (case) studies (*García-
 470 Palacios et al., 2015*). We extracted means or sums, standard deviations, and sample sizes of litter
 471 decomposition, decomposer diversity, and abundance (outcomes) in non-impacted vs. impacted
 472 sites (control-treatment studies), or at each site when gradients of chemical stressors or nutrients
 473 were investigated (gradient studies). When response variables were reported at different time
 474 points, we kept only the last time point to capture long-term responses. For studies reporting
 475 decomposition, decomposer abundance or diversity for several litter types (e.g. different litter
 476 species), several groups of organisms (e.g. functional feeding groups for macroinvertebrates),
 477 and several diversity metrics (e.g. Shannon indices and taxon richness), we created separate
 478 observations within case studies. We also extracted chemical stressor or nutrient levels at those
 479 sites (water, soil, or sediment concentrations of chemical stressors or nutrients, or application rate
 480 of pesticides or fertilizers). The study type (experimental vs. observational), taxonomic group (animal
 481 decomposers or microbial decomposers) and metric of diversity (taxa richness or diversity indices
 482 (Shannon diversity and evenness)) were also recorded. We used the online software Webplotdigitizer
 483 to extract data from figures (*Rohatgi, 2018*). We converted standard errors and confidence intervals
 484 into standard deviations using the equations in *Lajeunesse (2013)*. When reported as mass loss,
 485 litter decomposition data were transformed into k rates using the exponential decay equation used
 486 in *Ferreira et al. (2015)*.

487 **Effect size calculation**

488 We used z-transformed correlation coefficients as effect sizes in order to cope with the heterogeneity
 489 of data and study types (*Koricheva et al., 2013*). For control-treatment studies, we first calculated
 490 Hedge's d, and then transformed Hedge's d into correlation coefficients (*Lajeunesse, 2013*). For
 491 gradient studies (4 or more treatment levels), we calculated correlation coefficients between the
 492 mean values of abundance, diversity, or decomposition rate and the corresponding chemical
 493 stressor or nutrient concentrations. When means, standard deviations, or sample sizes were
 494 missing, we contacted the authors to retrieve the data. When the information could not be
 495 retrieved, standard deviations were approximated from the data, using the linear relationship
 496 between mean values and standard deviations across our datasets (*Lajeunesse, 2013*).

497 **Standardization of chemical stressors and nutrient enrichment intensities**

498 Given the variability in the different stressors and nutrients combinations in the studies, stressor and
 499 nutrient levels were standardized into a common environmental change driver intensity ($ECD_{intensity}$)
 500 as follows:

$$501 \quad ECD_{intensity} = \log([Compound_i]_{treatment} / [Compound_i]_{criteria})$$

502 where $[Compound_i]_{criteria}$ were environmental quality criteria set by European or US environ-
 503 mental authorities for the chemical stressor or nutrient considered (*Appendix 1-Table 1*), and

504 [$Compound_i$]_{treatment} were the concentrations of the chemical stressor or nutrient at the treatment
 505 or impacted sites. When multiple stressors or nutrients were reported, we used the standardized
 506 intensity of the stressor or nutrient corresponding to the highest standardized intensity for the rest
 507 of the analyses.

508 We used consistent sources for the environmental quality criteria as much as possible. For
 509 chemicals, we relied primarily on quality criteria from the European Chemical Agency (ECHA)
 510 and United States Environmental Protection Agency (USEPA) that use standardized procedures
 511 across aquatic and terrestrial realms based on ecotoxicological data. For nutrients, we relied
 512 mostly on European Water Framework Directive (WFD) benchmarks. Using various sources for
 513 those quality criteria was inevitable due to the high number of chemicals and the various way the
 514 authors reported stressor or nutrient levels in individual studies. When we could not find quality
 515 criteria for the stressors or nutrients considered in the studies in our main sources, we relied
 516 on the authors' statements and expert knowledge regarding their stressor or nutrient levels (e.g.
 517 recommended application rates of pesticides, citation for ecotoxicological data, or synthesis studies,
 518 **(Appendix 1-Table 1)**). Despite this, the final datasets were all dominated by similar sources for
 519 standardizing stressor and nutrient intensity levels: thresholds from ECHA or USEPA for 80 and
 520 90% of observations in the stressor-diversity and stressor-abundance datasets, respectively, and
 521 for nutrients, thresholds from WFD for 100 and 75% of observations in the nutrient-diversity and
 522 nutrient-abundance datasets, respectively.

523 **Overall effects of chemical stressors and nutrient enrichment: first-level meta-** 524 **analysis**

525 We first tested the differences between the effects of chemical stressors and nutrient enrichment
 526 on decomposer diversity, abundance and litter decomposition responses by quantifying the grand
 527 mean effect sizes on the three response variables (first level meta-analysis). Three separate meta-
 528 analyses were conducted, one for each response variable, and included the type of driver (stressors
 529 or nutrients) as a categorical moderator, and a random effect of the case study. We used a weighted
 530 meta-analysis giving more weight to effect sizes derived from studies with larger sample sizes.
 531 Weights were the inverse of the variance in z-transformed correlation coefficients (**Viechtbauer,**
 532 **2010**). Publication bias was evaluated using funnel plots with environmental change driver type as
 533 covariate. The intercepts from Egger's regressions (standardized effect size vs. precision = 1/SE)
 534 were inspected for significant deviation from zero that would indicate publication bias (**Koricheva**
 535 **et al., 2013**). Residual plots were used to detect strong deviation from normality and outliers.
 536 We estimated the grand mean effect sizes and compared the effect of chemical stressors and of
 537 nutrients using Wald-type chi-square tests. The `rma.mv()` function of the R package `metafor` was
 538 used (**R Core Team, 2018; Viechtbauer, 2010**).

539 **Relationship between biodiversity and decomposition: Structural equation mod-** 540 **elling**

541 An SEM was fitted to estimate the relationship between decomposer diversity or abundance and
 542 litter decomposition responses to environmental change drivers while controlling for the joint
 543 influence of stressor or nutrient intensity and categorical covariates. We used piecewise SEM
 544 (**Lefcheck, 2016**) estimating two linear mixed effect models, one for decomposition (z_{LD}) and one
 545 for decomposer diversity or abundance responses (z_B), with a random effect of the case study
 546 on the intercepts. These two sub-models embedded in the piecewise SEM were the second-level
 547 meta-analyses in our hierarchical approach. The random effect structure, weighting approach and
 548 variance structure were coded with the R package `nlme` (**Pinheiro et al., 2018**) in a way that fully
 549 reproduced the meta-analysis approach of weighting and of known residual variance (**Viechtbauer,**
 550 **2016**):

$$551 \quad z_{LD} \sim z_B + ECD_{intensity} + study\ type, \ random = \sim 1 | Case\ study / ID$$

$$552 \quad z_B \sim ECD_{intensity} + study\ type + taxonomic\ group (+diversity\ metric), \ random = \sim 1 | Case\ study / ID$$

553 This SEM was tested separately for each of four datasets: Stressors – Biodiversity; Stressors
 554 – Abundance; Nutrients – Biodiversity and Nutrients – Abundance datasets. The influence of the
 555 diversity metric (diversity indices versus taxa richness) was tested in the Biodiversity datasets only.
 556 We initially considered more complex model structures, but were unable to use them for analysis
 557 due to data limitations (in particular the effect of the ecosystem type and of interactions between
 558 our covariates).

559 Outliers, relationships between covariates, and non-linear patterns between continuous covari-
 560 ates were explored graphically. Studies often reported different decomposer diversity or abundance
 561 values for the same litter decomposition (e.g. when several taxonomic or functional groups were
 562 reported in the same litterbag). This variability could have affected the model estimates. We
 563 thus used data resampling to account for duplicated effect sizes on litter decomposition in the
 564 analyses. A stratified resampling was conducted, where for each duplicated value of effect size on
 565 decomposition, one randomly selected effect size on biodiversity was kept at each out of 1,000
 566 iterations. The models were fitted for each data resampling iteration, and we averaged model
 567 estimates and statistics across iterations and used the means as final values (path coefficients and
 568 standard error of the path and intercepts, Chi-square statistics and AICs).

569 Goodness-of-fit of the SEMs was assessed using directed separation tests based on the Fisher's
 570 C statistic. We used mediation tests to explore the significance of the path between decomposer
 571 diversity or abundance and litter decomposition based on the Fisher's C statistic of SEM that did not
 572 include the biodiversity-mediated path (*Lefcheck, 2016; Shipley, 2009*). We calculated the *P*-value
 573 associated with the mean Fisher's C statistic across data resampling iterations (*P*-value < 0.05
 574 indicated poor model fit). The AICs of models with and without the biodiversity-mediated paths
 575 were further compared using averaged AICs across data resampling iterations. We considered
 576 the biodiversity (or abundance) path to be consistent with the data when the SEM without the
 577 biodiversity-path had *P*-value < 0.05 (poor fit) and was not associated with a better AIC value
 578 (i.e. lower than 2 units) than the SEM including the biodiversity path. Residuals from the two sub-
 579 models of each SEM were graphically evaluated for strong departure to normality and relationship
 580 with the fitted values (*Duffy et al., 2015*). For these analyses, we averaged the residuals across
 581 data resampling iterations for each observation. We finally compared the relative magnitude of
 582 the biodiversity-mediated path versus the direct path from stressor or nutrient intensity to litter
 583 decomposition based on the mathematical product of the standardized path coefficients (*Grace,*
 584 *2006*).

585 **Moderator analyses: Second-level meta-analyses**

586 In order to quantify the influence of the categorical (study type, taxonomic group and diversity met-
 587 rics) and continuous (environmental change intensity) moderators on the three response variables,
 588 we further analyzed the results of the second-level meta-analyses (i.e. the sub-models embedded
 589 in the SEMs). The data resampling used in the SEM was no longer necessary, because there were
 590 no repeated values of decomposition matching different decomposer diversity or abundance mea-
 591 surements in this univariate approach. We quantified the effects of the different moderators based
 592 on the Wald-type chi-square tests derived with the R package metafor (*Viechtbauer, 2010*).

593 **Sensitivity analyses**

594 We finally tested the robustness of the results to the approximation of standard deviations, the
 595 presence of extreme values, and the metric of effect size used. The analyses were re-run with
 596 datasets that did not include the effect sizes for which we approximated standard deviations, for
 597 datasets that did not include extreme values of effect sizes (values beyond the whiskers of boxplots
 598 i.e. below quantile 1 minus 1.5 times the interquartile range or above quantile 3 plus 1.5 times the
 599 interquartile range). Finally, we calculated log-response ratios instead of correlation coefficients as
 600 effect sizes and re-run the analyses.

Acknowledgements

L.B. was funded by the Synthesis Centre (sDiv) of the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German Research Foundation (FZT 118). We are grateful to Helen R. Phillips, Benjamin Rosenbaum, Adam T. Clark and Katharina Gerstner for statistical advice and to Simone Cesarz for creating the images for Fig. 1.

Authors contribution

All authors conceived the project; L.B. collected the data, performed the analyses and wrote the manuscript; all authors discussed the results and contributed to the manuscript text.

References

- Bardgett RD**, van der Putten WH. Belowground Biodiversity and Ecosystem Functioning. *Nature*. 2014 Nov; 515(7528):505–511. doi: 10.1038/nature13855.
- Beketov MA**, Kefford BJ, Schäfer RB, Liess M. Pesticides Reduce Regional Biodiversity of Stream Invertebrates. *Proceedings of the National Academy of Sciences*. 2013 Jul; 110(27):11039–11043. doi: 10.1073/pnas.1305618110.
- Bergfur J**, Johnson RK, Sandin L, Goedkoop W, Nygren K. Effects of Nutrient Enrichment on Boreal Streams: Invertebrates, Fungi and Leaf-Litter Breakdown. *Freshwater Biology*. 2007 Aug; 52(8):1618–1633. doi: 10.1111/j.1365-2427.2007.01770.x.
- Bernhardt ES**, Rosi EJ, Gessner MO. Synthetic Chemicals as Agents of Global Change. *Frontiers in Ecology and the Environment*. 2017 Mar; 15(2):84–90. doi: 10.1002/fee.1450.
- Cadotte MW**, Carscadden K, Mirotnick N. Beyond Species: Functional Diversity and the Maintenance of Ecological Processes and Services. *Journal of Applied Ecology*. 2011 Oct; 48(5):1079–1087. doi: 10.1111/j.1365-2664.2011.02048.x, wOS:000295095100003.
- Cárdenas RE**, Donoso DA, Argoti A, Dangles O. Functional Consequences of Realistic Extinction Scenarios in Amazonian Soil Food Webs. *Ecosphere*. 2017; 8(2):e01692. doi: 10.1002/ecs2.1692, _eprint: https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1002/ecs2.1692.
- Cardinale BJ**, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, Narwani A, Mace GM, Tilman D, Wardle DA, Kinzig AP, Daily GC, Loreau M, Grace JB, Larigauderie A, Srivastava DS, Naeem S. Biodiversity Loss and Its Impact on Humanity. *Nature*. 2012 Jun; 486(7401):59–67. doi: 10.1038/nature11148.
- Chase JM**, Knight TM. Scale-Dependent Effect Sizes of Ecological Drivers on Biodiversity: Why Standardised Sampling Is Not Enough. *Ecology Letters*. 2013 May; 16:17–26. doi: 10.1111/ele.12112.
- Chase JM**, Leibold MA. *Ecological Niches: Linking Classical and Contemporary Approaches*. University of Chicago Press; 2003.
- Craven D**, Eisenhauer N, Pearse WD, Hautier Y, Isbell F, Roscher C, Bahn M, Beierkuhnlein C, Bönisch G, Buchmann N, Byun C, Catford JA, Cerabolini BEL, Cornelissen JHC, Craine JM, Luca ED, Ebeling A, Griffin JN, Hector A, Hines J, et al. Multiple Facets of Biodiversity Drive the Diversity–Stability Relationship. *Nature Ecology & Evolution*. 2018 Oct; 2(10):1579–1587. doi: 10.1038/s41559-018-0647-7.
- Dainese M**, Martin EA, Aizen MA, Albrecht M, Bartomeus I, Bommarco R, Carvalheiro LG, Chaplin-Kramer R, Gagic V, Garibaldi LA, Ghazoul J, Grab H, Jonsson M, Karp DS, Kennedy CM, Kleijn D, Kremen C, Landis DA, Letourneau DK, Marini L, et al. A Global Synthesis Reveals Biodiversity-Mediated Benefits for Crop Production. *Science Advances*. 2019 Oct; 5(10):eaax0121. doi: 10.1126/sciadv.aax0121.
- Dangles O**, Carpio C, Woodward G. Size-Dependent Species Removal Impairs Ecosystem Functioning in a Large-Scale Tropical Field Experiment. *Ecology*. 2012; 93(12):2615–2625. doi: 10.1890/12-0510.1, _eprint: https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/12-0510.1.
- Dangles O**, Malmqvist B. Species Richness–Decomposition Relationships Depend on Species Dominance. *Ecology Letters*. 2004 May; 7(5):395–402. doi: 10.1111/j.1461-0248.2004.00591.x.
- De Laender F**. Community- and Ecosystem-Level Effects of Multiple Environmental Change Drivers: Beyond Null Model Testing. *Global Change Biology*. 2018 Nov; 24(11):5021–5030. doi: 10.1111/gcb.14382, wOS:000447760300003.

- 649 **De Laender F**, Janssen CR. Brief Communication: The Ecosystem Perspective in Ecotoxicology as a Way Forward
 650 for the Ecological Risk Assessment of Chemicals: How Ecosystem Ecotoxicology Can Inform Risk Assessment.
 651 *Integrated Environmental Assessment and Management*. 2013 Jul; 9(3):e34–e38. doi: [10.1002/ieam.1428](https://doi.org/10.1002/ieam.1428).
- 652 **De Laender F**, Rohr JR, Ashauer R, Baird DJ, Berger U, Eisenhauer N, Grimm V, Hommen U, Maltby L, Meliàn
 653 CJ, Pomati F, Roessink I, Radchuk V, Van den Brink PJ. Reintroducing Environmental Change Drivers in
 654 Biodiversity-Ecosystem Functioning Research. *Trends in Ecology & Evolution*. 2016 Dec; 31(12):905–915. doi:
 655 [10.1016/j.tree.2016.09.007](https://doi.org/10.1016/j.tree.2016.09.007).
- 656 **DeLorenzo ME**, Scott GI, Ross PE. Toxicity of Pesticides to Aquatic Microorganisms: A Review. *En-*
 657 *vironmental Toxicology and Chemistry*. 2001; 20(1):84–98. doi: [10.1002/etc.5620200108](https://doi.org/10.1002/etc.5620200108),
 658 *_eprint:* <https://setac.onlinelibrary.wiley.com/doi/pdf/10.1002/etc.5620200108>.
- 659 **Des Roches S**, Post DM, Turley NE, Bailey JK, Hendry AP, Kinnison MT, Schweitzer JA, Palkovacs EP. The Ecological
 660 Importance of Intraspecific Variation. *Nature Ecology & Evolution*. 2018 Jan; 2(1):57. doi: [10.1038/s41559-017-](https://doi.org/10.1038/s41559-017-0402-5)
 661 [0402-5](https://doi.org/10.1038/s41559-017-0402-5).
- 662 **Dib V**, Pires APF, Nova CC, Bozelli RL, Farjalla VF. Biodiversity-Mediated Effects on Ecosystem Functioning Depend
 663 on the Type and Intensity of Environmental Disturbances. *Oikos*. 2020; 129(3):433–443. doi: [10.1111/oik.06768](https://doi.org/10.1111/oik.06768),
 664 *_eprint:* <https://onlinelibrary.wiley.com/doi/pdf/10.1111/oik.06768>.
- 665 **Duffy JE**, Godwin CM, Cardinale BJ. Biodiversity Effects in the Wild Are Common and as Strong as Key Drivers of
 666 Productivity. *Nature*. 2017 Sep; 549(7671):261–264. doi: [10.1038/nature23886](https://doi.org/10.1038/nature23886).
- 667 **Duffy JE**, Reynolds PL, Boström C, Coyer JA, Cusson M, Donadi S, Douglass JG, Eklöf JS, Engelen AH, Eriksson BK,
 668 Fredriksen S, Gamfeldt L, Gustafsson C, Hoarau G, Hori M, Hovel K, Iken K, Lefcheck JS, Moksnes PO, Nakaoka
 669 M, et al. Biodiversity Mediates Top-down Control in Eelgrass Ecosystems: A Global Comparative-Experimental
 670 Approach. *Ecology Letters*. 2015; 18(7):696–705. doi: [10.1111/ele.12448](https://doi.org/10.1111/ele.12448).
- 671 **Ebeling A**, Meyer ST, Abbas M, Eisenhauer N, Hillebrand H, Lange M, Scherber C, Vogel A, Weigelt A, Weisser
 672 WW. Plant Diversity Impacts Decomposition and Herbivory via Changes in Aboveground Arthropods. *PLoS*
 673 *One*. 2014 Sep; 9(9):e106529. doi: [10.1371/journal.pone.0106529](https://doi.org/10.1371/journal.pone.0106529).
- 674 **Eisenhauer N**, Barnes AD, Cesarz S, Craven D, Ferlian O, Gottschall F, Hines J, Sendek A, Siebert J, Thakur MP,
 675 Türke M. Biodiversity-Ecosystem Function Experiments Reveal the Mechanisms Underlying the Consequences
 676 of Biodiversity Change in Real World Ecosystems. *Journal of Vegetation Science*. 2016 Sep; 27(5):1061–1070.
 677 doi: [10.1111/jvs.12435](https://doi.org/10.1111/jvs.12435).
- 678 **Eisenhauer N**, Bonn A, Guerra CA. Recognizing the Quiet Extinction of Invertebrates. *Nature Communications*.
 679 2019 Jan; 10(1):50. doi: [10.1038/s41467-018-07916-1](https://doi.org/10.1038/s41467-018-07916-1).
- 680 **Eisenhauer N**, Hörsch V, Moeser J, Scheu S. Synergistic Effects of Microbial and Animal Decomposers
 681 on Plant and Herbivore Performance. *Basic and Applied Ecology*. 2010 Feb; 11(1):23–34. doi:
 682 [10.1016/j.baae.2009.11.001](https://doi.org/10.1016/j.baae.2009.11.001).
- 683 **Eisenhauer N**, Reich PB, Isbell F. Decomposer Diversity and Identity Influence Plant Diversity Effects on
 684 Ecosystem Functioning. *Ecology*. 2012 Oct; 93(10):2227–2240. doi: [10.1890/11-2266.1](https://doi.org/10.1890/11-2266.1).
- 685 **Eisenhauer N**, Schielzeth H, Barnes AD, Barry K, Bonn A, Brose U, Bruehlheide H, Buchmann N, Buscot F, Ebeling
 686 A, Ferlian O, Freschet GT, Giling DP, Hättenschwiler S, Hillebrand H, Hines J, Isbell F, Koller-France E, König-Ries
 687 B, de Kroon H, et al. A Multitrophic Perspective on Biodiversity-Ecosystem Functioning Research. In: *Advances*
 688 *in Ecological Research* Academic Press; 2019.
- 689 **Ferreira V**, Castagnyrol B, Koricheva J, Gulis V, Chauvet E, Graça MAS. A Meta-Analysis of the Effects of
 690 Nutrient Enrichment on Litter Decomposition in Streams. *Biological Reviews*. 2015 Aug; 90(3):669–688. doi:
 691 [10.1111/brv.12125](https://doi.org/10.1111/brv.12125).
- 692 **Ferreira V**, Koricheva J, Duarte S, Niyogi DK, Guérol F. Effects of Anthropogenic Heavy Metal Contamination
 693 on Litter Decomposition in Streams - A Meta-Analysis. *Environmental Pollution*. 2016; 210:261–270. doi:
 694 [10.1016/j.envpol.2015.12.060](https://doi.org/10.1016/j.envpol.2015.12.060).
- 695 **Follstad Shah JJ**, Kominoski JS, Ardón M, Dodds WK, Gessner MO, Griffiths NA, Hawkins CP, Johnson SL, Lecerf
 696 A, LeRoy CJ, Manning DWP, Rosemond AD, Sinsabaugh RL, Swan CM, Webster JR, Zeglin LH. Global Synthesis
 697 of the Temperature Sensitivity of Leaf Litter Breakdown in Streams and Rivers. *Global Change Biology*. 2017
 698 Aug; 23(8):3064–3075. doi: [10.1111/gcb.13609](https://doi.org/10.1111/gcb.13609).

- 699 **García-Palacios P**, Maestre FT, Kattge J, Wall DH. Climate and Litter Quality Differently Modulate the Effects
700 of Soil Fauna on Litter Decomposition across Biomes. *Ecology Letters*. 2013 Aug; 16(8):1045–1053. doi:
701 [10.1111/ele.12137](https://doi.org/10.1111/ele.12137).
- 702 **García-Palacios P**, Vandegehuchte ML, Shaw EA, Dam M, Post KH, Ramirez KS, Sylvain ZA, de Tomasel CM, Wall
703 DH. Are There Links between Responses of Soil Microbes and Ecosystem Functioning to Elevated CO₂, N
704 Deposition and Warming? A Global Perspective. *Global Change Biology*. 2015 Apr; 21(4):1590–1600. doi:
705 [10.1111/gcb.12788](https://doi.org/10.1111/gcb.12788).
- 706 **Gessner MO**, Swan CM, Dang CK, McKie BG, Bardgett RD, Wall DH, Hättenschwiler S. Diversity Meets Decompo-
707 sition. *Trends in Ecology & Evolution*. 2010 Jun; 25(6):372–380. doi: [10.1016/j.tree.2010.01.010](https://doi.org/10.1016/j.tree.2010.01.010).
- 708 **Gilling DP**, Beaumelle L, Phillips HRP, Cesarz S, Eisenhauer N, Ferlian O, Gottschall F, Guerra C, Hines J, Sendek A,
709 Siebert J, Thakur MP, Barnes AD. A Niche for Ecosystem Multifunctionality in Global Change Research. *Global*
710 *Change Biology*. 2019; 25(3):763–774. doi: [10.1111/gcb.14528](https://doi.org/10.1111/gcb.14528).
- 711 **Giller KE**, Witter E, McGrath SP. Heavy Metals and Soil Microbes. *Soil Biology and Biochemistry*. 2009 Oct;
712 41(10):2031–2037. doi: [10.1016/j.soilbio.2009.04.026](https://doi.org/10.1016/j.soilbio.2009.04.026).
- 713 **Glassman SI**, Weihe C, Li J, Albright MBN, Looby CI, Martiny AC, Treseder KK, Allison SD, Martiny JBH. Decom-
714 position Responses to Climate Depend on Microbial Community Composition. *Proceedings of the National*
715 *Academy of Sciences*. 2018 Nov; 115(47):11994–11999. doi: [10.1073/pnas.1811269115](https://doi.org/10.1073/pnas.1811269115).
- 716 **Gotelli NJ**, Colwell RK. Quantifying Biodiversity: Procedures and Pitfalls in the Measurement and Comparison
717 of Species Richness. *Ecology Letters*. 2001 Jul; 4(4):379–391. doi: [10.1046/j.1461-0248.2001.00230.x](https://doi.org/10.1046/j.1461-0248.2001.00230.x).
- 718 **Grace JB**. *Structural Equation Modelling and Natural Systems*. Cambridge University Press; 2006.
- 719 **Grime JP**. Benefits of Plant Diversity to Ecosystems: Immediate, Filter and Founder Effects.
720 *Journal of Ecology*. 1998; 86(6):902–910. doi: [10.1046/j.1365-2745.1998.00306.x](https://doi.org/10.1046/j.1365-2745.1998.00306.x), _eprint:
721 <https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1046/j.1365-2745.1998.00306.x>.
- 722 **Handa IT**, Aerts R, Berendse F, Berg MP, Bruder A, Butenschoen O, Chauvet E, Gessner MO, Jabiol J, Makkonen
723 M, McKie BG, Malmqvist B, Peeters ETHM, Scheu S, Schmid B, van Ruijven J, Vos VCA, Hättenschwiler S.
724 Consequences of Biodiversity Loss for Litter Decomposition across Biomes. *Nature*. 2014 May; 509(7499):218–
725 221. doi: [10.1038/nature13247](https://doi.org/10.1038/nature13247).
- 726 **Harley CDG**, Connell SD, Doubleday ZA, Kelaher B, Russell BD, Sarà G, Helmuth B. Conceptualizing Ecosys-
727 tem Tipping Points within a Physiological Framework. *Ecology and Evolution*. 2017; 7(15):6035–6045. doi:
728 [10.1002/ece3.3164](https://doi.org/10.1002/ece3.3164), _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/ece3.3164>.
- 729 **Hättenschwiler S**, Tiunov AV, Scheu S. Biodiversity and Litter Decomposition in Terrestrial Ecosys-
730 tems. *Annual Review of Ecology, Evolution, and Systematics*. 2005; 36(1):191–218. doi: [10.1146/an-
731 nurev.ecolsys.36.112904.151932](https://doi.org/10.1146/annurev.ecolsys.36.112904.151932).
- 732 **Heemsbergen DA**, Berg MP, Loreau M, van Hal JR, Faber JH, Verhoef HA. Biodiversity Effects on Soil Pro-
733 cesses Explained by Interspecific Functional Dissimilarity. *Science*. 2004 Nov; 306(5698):1019–1020. doi:
734 [10.1126/science.1101865](https://doi.org/10.1126/science.1101865).
- 735 **Hines J**, Eisenhauer N, Drake BG. Inter-Annual Changes in Detritus-Based Food Chains Can Enhance Plant
736 Growth Response to Elevated Atmospheric CO₂. *Global Change Biology*. 2015; 21(12):4642–4650. doi:
737 [10.1111/gcb.12965](https://doi.org/10.1111/gcb.12965).
- 738 **Hogsden KL**, Harding JS. Consequences of Acid Mine Drainage for the Structure and Function of Benthic Stream
739 Communities: A Review. *Freshwater Science*. 2012 Mar; 31(1):108–120. doi: [10.1899/11-091.1](https://doi.org/10.1899/11-091.1).
- 740 **IPBES**. Summary for Policymakers of the Global Assessment Report on Biodiversity and Ecosystem Services of
741 the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. IPBES secretariat,
742 Bonn, Germany. 2019; p. 44.
- 743 **Jonsson M**, Dangles O, Malmqvist B, Guérol F. Simulating Species Loss Following Perturbation: Assessing
744 the Effects on Process Rates. *Proceedings of the Royal Society of London B: Biological Sciences*. 2002;
745 269(1495):1047–1052.
- 746 **Kirschbaum MUF**. Will Changes in Soil Organic Carbon Act as a Positive or Negative Feedback on Global
747 Warming? *Biogeochemistry*. 2000 Jan; 48(1):21–51. doi: [10.1023/A:1006238902976](https://doi.org/10.1023/A:1006238902976).

- 748 **Knorr M**, Frey SD, Curtis PS. Nitrogen Additions And Litter Decomposition: A Meta-Analysis. *Ecology*. 2005 Dec;
749 86(12):3252–3257. doi: 10.1890/05-0150.
- 750 **Koricheva J**, Gurevitch J, Mengersen K. *Handbook of Meta-Analysis in Ecology and Evolution*. Princeton
751 University Press; 2013.
- 752 **Lajeunesse M**. Recovering Missing or Partial Data from Studies: A Survey of Conversions and Imputations for
753 Meta-Analysis. In: *Handbook of Meta-Analysis in Ecology and Evolution* Princeton University Press; 2013.
- 754 **Lecerf A**, Chauvet E. Diversity and Functions of Leaf-Decaying Fungi in Human-Altered Streams. *Freshwater
755 Biology*. 2008 Aug; 53(8):1658–1672. doi: 10.1111/j.1365-2427.2008.01986.x.
- 756 **Lefcheck JS**. PIECEWISESEM : Piecewise Structural Equation Modelling in R for Ecology, Evolution, and Systematics.
757 *Methods in Ecology and Evolution*. 2016 May; 7(5):573–579. doi: 10.1111/2041-210X.12512.
- 758 **Lefcheck JS**, Byrnes JEK, Isbell F, Gamfeldt L, Griffin JN, Eisenhauer N, Hensel MJS, Hector A, Cardinale BJ,
759 Duffy JE. Biodiversity Enhances Ecosystem Multifunctionality across Trophic Levels and Habitats. *Nature
760 Communications*. 2015 Apr; 6:6936. doi: 10.1038/ncomms7936, wOS:000353704400005.
- 761 **Lucisine P**, Lecerf A, Danger M, Felten V, Aran D, Auclerc A, Gross EM, Huot H, Morel JL, Muller S, Nahmani
762 J, Maunoury-Danger F. Litter Chemistry Prevails over Litter Consumers in Mediating Effects of Past Steel
763 Industry Activities on Leaf Litter Decomposition. *Science of The Total Environment*. 2015 Dec; 537(Supplement
764 C):213–224. doi: 10.1016/j.scitotenv.2015.07.112.
- 765 **Malaj E**, Peter C, Grote M, Kühne R, Mondy CP, Usseglio-Polatera P, Brack W, Schäfer RB. Organic Chemicals
766 Jeopardize the Health of Freshwater Ecosystems on the Continental Scale. *Proceedings of the National
767 Academy of Sciences*. 2014; 111(26):9549–9554.
- 768 **Manning P**, Cutler GC. Ecosystem Functioning Is More Strongly Impaired by Reducing Dung Beetle Abun-
769 dance than by Reducing Species Richness. *Agriculture Ecosystems & Environment*. 2018; 264:9–14. doi:
770 10.1016/j.agee.2018.05.002, wOS:000437392100002.
- 771 **Mazor T**, Doropoulos C, Schwarzmüller F, Gladish DW, Kumaran N, Merkel K, Di Marco M, Gagic V. Global
772 Mismatch of Policy and Research on Drivers of Biodiversity Loss. *Nature Ecology & Evolution*. 2018 Jul;
773 2(7):1071–1074. doi: 10.1038/s41559-018-0563-x.
- 774 **Melguizo-Ruiz N**, Jiménez-Navarro G, De Mas E, Pato J, Scheu S, Austin AT, Wise DH, Moya-Laraño J. Field
775 Exclusion of Large Soil Predators Impacts Lower Trophic Levels and Decreases Leaf-litter Decomposition in
776 Dry Forests. *Journal of Animal Ecology*. 2020 Feb; 89(2):334–346. doi: 10.1111/1365-2656.13101.
- 777 **Oliveira RAC**, Marques R, Marques MCM. Plant Diversity and Local Environmental Conditions Indirectly
778 Affect Litter Decomposition in a Tropical Forest. *Applied Soil Ecology*. 2019 Feb; 134:45–53. doi:
779 10.1016/j.apsoil.2018.09.016.
- 780 **Pelosi C**, Barot S, Capowiez Y, Hedde M, Vandenbulcke F. Pesticides and Earthworms. A Review. *Agronomy for
781 Sustainable Development*. 2014 Jan; 34(1):199–228. doi: 10.1007/s13593-013-0151-z.
- 782 **Peters K**, Bundschuh M, Schäfer RB. Review on the Effects of Toxicants on Freshwater Ecosystem Functions.
783 *Environmental Pollution*. 2013 Sep; 180(Supplement C):324–329. doi: 10.1016/j.envpol.2013.05.025.
- 784 **Petrin Z**, Englund G, Malmqvist B. Contrasting Effects of Anthropogenic and Natural Acidity in Streams: A Meta-
785 Analysis. *Proceedings of the Royal Society of London Series B: Biological Sciences*. 2008 May; 275(1639):1143–
786 1148. doi: 10.1098/rspb.2008.0023.
- 787 **Pinheiro J**, Bates D, DebRoy S, Sarkar D, R Core Team. *Nlme: Linear and Nonlinear Mixed Effects Models*; 2018.
- 788 **R Core Team**. *R: A Language and Environment for Statistical Computing*. Vienna, Austria; 2018.
- 789 **Rillig MC**, Ryo M, Lehmann A, Aguilar-Trigueros CA, Buchert S, Wulf A, Iwasaki A, Roy J, Yang G. The Role of
790 Multiple Global Change Factors in Driving Soil Functions and Microbial Biodiversity. *Science*. 2019 Nov;
791 366(6467):886–890. doi: 10.1126/science.aay2832.
- 792 **Rohatgi A**, *WebPlotDigitizer*; 2018. <https://automeris.io/WebPlotDigitizer>.
- 793 **Schäfer RB**. Responses of Freshwater Macroinvertebrates to Pesticides: Insights from Field Studies. *Current
794 Opinion in Environmental Science & Health*. 2019; 11:1–7. doi: 10.1016/j.coesh.2019.06.001.

- 795 **Schuldt A**, Assmann T, Brezzi M, Buscot F, Eichenberg D, Gutknecht J, Härdtle W, He JS, Klein AM, Kühn P, Liu X,
796 Ma K, Niklaus PA, Pietsch KA, Purahong W, Scherer-Lorenzen M, Schmid B, Scholten T, Staab M, Tang Z, et al.
797 Biodiversity across Trophic Levels Drives Multifunctionality in Highly Diverse Forests. *Nature Communications*.
798 2018 Jul; 9(1):2989. doi: 10.1038/s41467-018-05421-z.
- 799 **Sheridan JA**, Bickford D. Shrinking Body Size as an Ecological Response to Climate Change. *Nature Climate*
800 *Change*. 2011 Nov; 1(8):401–406. doi: 10.1038/nclimate1259.
- 801 **Shipley B**. Confirmatory Path Analysis in a Generalized Multilevel Context. *Ecology*. 2009; 90(2):363–368.
- 802 **Siebert J**, Thakur MP, Reitz T, Schädler M, Schulz E, Yin R, Weigelt A, Eisenhauer N. Extensive Grassland-
803 Use Sustains High Levels of Soil Biological Activity, but Does Not Alleviate Detrimental Climate Change
804 Effects. In: Bohan DA, Dumbrell AJ, editors. *Advances in Ecological Research*, vol. 60; 2019.p. 25–58. doi:
805 [10.1016/bs.aecr.2019.02.002](https://doi.org/10.1016/bs.aecr.2019.02.002).
- 806 **Smith MD**, Knapp AK. Dominant Species Maintain Ecosystem Function with Non-Random Species Loss. *Ecology*
807 *Letters*. 2003; 6(6):509–517.
- 808 **Soliveres S**, van der Plas F, Manning P, Prati D, Gossner MM, Renner SC, Alt F, Arndt H, Baumgartner V, Binkenstein
809 J, Birkhofer K, Blaser S, Blüthgen N, Boch S, Böhm S, Börschig C, Buscot F, Diekötter T, Heinze J, Hölzel N,
810 et al. Biodiversity at Multiple Trophic Levels Is Needed for Ecosystem Multifunctionality. *Nature*. 2016 Aug;
811 536(7617):456–459. doi: 10.1038/nature19092.
- 812 **Spaak JW**, Baert JM, Baird DJ, Eisenhauer N, Maltby L, Pomati F, Radchuk V, Rohr JR, Van den Brink PJ, De Laender
813 F. Shifts of Community Composition and Population Density Substantially Affect Ecosystem Function despite
814 Invariant Richness. *Ecology Letters*. 2017 Oct; 20(10):1315–1324. doi: [10.1111/ele.12828](https://doi.org/10.1111/ele.12828).
- 815 **Srivastava DS**, Cardinale BJ, Downing AL, Duffy JE, Jouseau C, Sankaran M, Wright JP. Diversity Has Stronger
816 Top-down than Bottom-up Effects on Decomposition. *Ecology*. 2009; 90(4):1073–1083. doi: [10.1890/08-0439.1](https://doi.org/10.1890/08-0439.1).
- 817 **Srivastava DS**, Vellend M. Biodiversity-Ecosystem Function Research: Is It Relevant to Conservation?
818 *Annual Review of Ecology, Evolution, and Systematics*. 2005 Dec; 36(1):267–294. doi: [10.1146/an-](https://doi.org/10.1146/annurev.ecolsys.36.102003.152636)
819 [nurev.ecolsys.36.102003.152636](https://doi.org/10.1146/annurev.ecolsys.36.102003.152636).
- 820 **Steffen W**, Richardson K, Rockström J, Cornell SE, Fetzer I, Bennett EM, Biggs R, Carpenter SR, de Vries W,
821 de Wit CA, Folke C, Gerten D, Heinke J, Mace GM, Persson LM, Ramanathan V, Reyers B, Sörlin S. Planetary
822 Boundaries: Guiding Human Development on a Changing Planet. *Science*. 2015 Feb; 347(6223). doi:
823 [10.1126/science.1259855](https://doi.org/10.1126/science.1259855).
- 824 **Thakur MP**, Reich PB, Hobbie SE, Stefanski A, Rich R, Rice KE, Eddy WC, Eisenhauer N. Reduced Feeding Activity
825 of Soil Detritivores under Warmer and Drier Conditions. *Nature Climate Change*. 2018 Jan; 8(1):75–78. doi:
826 [10.1038/s41558-017-0032-6](https://doi.org/10.1038/s41558-017-0032-6).
- 827 **Tilman D**, Reich PB, Isbell F. Biodiversity Impacts Ecosystem Productivity as Much as Resources, Disturbance,
828 or Herbivory. *Proceedings of the National Academy of Sciences*. 2012 Jun; 109(26):10394–10397. doi:
829 [10.1073/pnas.1208240109](https://doi.org/10.1073/pnas.1208240109).
- 830 **Tonin AM**, Pozo J, Monroy S, Basaguren A, Pérez J, Gonçalves JF, Pearson R, Cardinale BJ, Boyero L. Interactions
831 between Large and Small Detritivores Influence How Biodiversity Impacts Litter Decomposition. *Journal of*
832 *Animal Ecology*. 2018 Sep; 87(5):1465–1474. doi: [10.1111/1365-2656.12876](https://doi.org/10.1111/1365-2656.12876).
- 833 **Treseder KK**. Nitrogen Additions and Microbial Biomass: A Meta-Analysis of Ecosystem Studies. *Ecology Letters*.
834 2008; 11(10):1111–1120. doi: [10.1111/j.1461-0248.2008.01230.x](https://doi.org/10.1111/j.1461-0248.2008.01230.x).
- 835 **van der Plas F**. Biodiversity and Ecosystem Functioning in Naturally Assembled Communities. *Biological*
836 *Reviews*. 2019 Aug; 94(4):1220–1245. doi: [10.1111/brv.12499](https://doi.org/10.1111/brv.12499).
- 837 **Viechtbauer W**. Conducting Meta-Analyses in R with the Metafor Package. *Journal of Statistical Software*. 2010
838 Aug; 36(3):1–48.
- 839 **Viechtbauer W**, A Comparison of the Rma() and the Lm(), Lme(), and Lmer() Functions [The Metafor Package];
840 2016. http://www.metafor-project.org/doku.php/tips:rma_vs_lm_lme_lmer.
- 841 **Voigt W**, Perner J, Hefin Jones T. Using Functional Groups to Investigate Community Response to Environmental
842 Changes: Two Grassland Case Studies. *Global Change Biology*. 2007 Aug; 13(8):1710–1721. doi: [10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2007.01398.x)
843 [2486.2007.01398.x](https://doi.org/10.1111/j.1365-2486.2007.01398.x).

- 844 **Wang S**, Brose U. Biodiversity and Ecosystem Functioning in Food Webs: The Vertical Di-
845 versity Hypothesis. *Ecology Letters*. 2018; 21(1):9–20. doi: [10.1111/ele.12865](https://onlinelibrary.wiley.com/doi/pdf/10.1111/ele.12865), _eprint:
846 <https://onlinelibrary.wiley.com/doi/pdf/10.1111/ele.12865>.
- 847 **Wang Z**, Walker GW, Muir DCG, Nagatani-Yoshida K. Toward a Global Understanding of Chemical Pollution:
848 A First Comprehensive Analysis of National and Regional Chemical Inventories. *Environmental Science &*
849 *Technology*. 2020 Jan; doi: [10.1021/acs.est.9b06379](https://doi.org/10.1021/acs.est.9b06379).
- 850 **Wardle DA**. Do Experiments Exploring Plant Diversity-Ecosystem Functioning Relationships Inform How Bio-
851 diversity Loss Impacts Natural Ecosystems? *Journal of Vegetation Science*. 2016 May; 27(3):646–653. doi:
852 [10.1111/jvs.12399](https://doi.org/10.1111/jvs.12399).
- 853 **Woodward G**, Ebenman B, Emmerson M, Montoya JM, Olesen JM, Valido A, Warren PH. Body Size in Ecological
854 Networks. *Trends in Ecology & Evolution*. 2005 Jul; 20(7):402–409. doi: [10.1016/j.tree.2005.04.005](https://doi.org/10.1016/j.tree.2005.04.005).
- 855 **Woodward G**, Gessner MO, Giller PS, Gulis V, Hladyz S, Lecerf A, Malmqvist B, McKie BG, Tiegs SD, Cariss H,
856 Dobson M, Eloisegi A, Ferreira V, Graça MAS, Fleituch T, Lacoursière JO, Nistorescu M, Pozo J, Risnoveanu G,
857 Schindler M, et al. Continental-Scale Effects of Nutrient Pollution on Stream Ecosystem Functioning. *Science*.
858 2012 Jun; 336(6087):1438–1440. doi: [10.1126/science.1219534](https://doi.org/10.1126/science.1219534).
- 859 **Yang X**, Yang Z, Warren MW, Chen J. Mechanical Fragmentation Enhances the Contribution of Collembola to Leaf
860 Litter Decomposition. *European Journal of Soil Biology*. 2012 Nov; 53:23–31. doi: [10.1016/j.ejsobi.2012.07.006](https://doi.org/10.1016/j.ejsobi.2012.07.006).
- 861 **Yvon-Durocher G**, Montoya JM, Trimmer M, Woodward G. Warming Alters the Size Spectrum and Shifts the
862 Distribution of Biomass in Freshwater Ecosystems. *Global Change Biology*. 2011; 17(4):1681–1694. doi:
863 [10.1111/j.1365-2486.2010.02321.x](https://doi.org/10.1111/j.1365-2486.2010.02321.x).
- 864 **Zavaleta ES**, Hulvey KB. Realistic Species Losses Disproportionately Reduce Grassland Resistance to Biological
865 Invaders. *Science*. 2004 Nov; 306(5699):1175–1177. doi: [10.1126/science.1102643](https://doi.org/10.1126/science.1102643).
- 866 **Zhao Q**, den Brink PJV, Carpentier C, Wang YXG, Rodríguez-Sánchez P, Xu C, Vollbrecht S, Gillissen F, Vollebregt
867 M, Wang S, Laender FD. Horizontal and Vertical Diversity Jointly Shape Food Web Stability against Small and
868 Large Perturbations. *Ecology Letters*. 2019; 22(7):1152–1162. doi: [10.1111/ele.13282](https://doi.org/10.1111/ele.13282).

869 **Appendix 1**870 **Supplementary Methods**871 **Search strategy**

872 We collected data from published papers reporting the effect of various global change
 873 drivers on both decomposition rates and decomposer communities. The search strategy first
 874 involved the selection of a relevant search term combination. We compared different search
 875 term combinations based on the number of studies retrieved, their potential relevance
 876 (based on screening the titles in the search), and on maximizing the retrieval of pre-identified
 877 papers that fully matched the inclusion criteria. We used search terms of previous meta-
 878 analyses and literature reviews (Garcia Palacios et al., 2016; Covich et al., 2004; Srivastava et
 879 al., 2009). The following search terms were used to identify studies looking at the impact of
 880 various global change drivers on both decomposition rates and decomposer communities.
 881 The search was done on ISI Web Of Science on November 17th 2017 and retrieved 2,536
 882 records.

883 TS=(“global change” OR “environmental change” OR disturbance* OR stress* OR “climat*
 884 change” OR drought OR temperature* OR warming OR heat* OR precipitation* OR rain*
 885 OR flood* OR irrigation OR moisture OR watering OR fire OR “carbon dioxide” OR CO2 OR
 886 acidification OR “nitrogen deposition” OR “nutrient deposition” OR “atmospheric deposition”
 887 OR *eutroph* OR fertil* OR “nutrient* enrichment” OR “nutrient pollut*” OR “land-use” OR
 888 “landuse” OR “agricultural intensi*” OR desertif* OR pollut* OR pesticide* OR metal* OR
 889 “over-exploit*” OR overexploit* OR toxi* OR contamin* OR over-fish* OR invasi* OR alien OR
 890 “habitat loss” OR “habitat fragment*” OR “habitat degrad*” OR “habitat destruct*”)

891 AND

892 TS = ((decomposition OR processing OR breakdown OR decay OR “mass loss”) AND (litter
 893 OR leaf OR leaves OR bark OR wood))

894 AND

895 TS= (“species richness” OR richness OR “number of species” OR “number of taxa”
 896 OR “species diversity” OR “taxonomic diversity” OR biodiversity OR Shannon* OR even-
 897 ness OR “community composition” OR “community structure” OR “functional diversity”
 898 OR “trait diversity” OR “functional traits” OR “functional group richness” OR “trait-based”)
 899 AND (decomposer* OR detritivore* OR *invertebrate* OR microb* OR microorganism* OR
 900 bacteri* OR fung* OR archaea OR shredder OR *invertebrate* OR hyphomycete* OR “leaf-
 901 shredding” OR “leaf-eat*” OR “leaf-consum*” “leaf-feed*” OR “litter-feed*” OR “litter-eat*”
 902 OR “litter-shredding” OR protozoa* OR protist* OR springtail OR collembol* OR mite* OR
 903 acari* OR enchytraeid* OR nematod* OR rotifer* OR isopod* OR earthworm* OR termite*
 904 OR microarthropod* OR macroarthropod* OR microfauna OR mesofauna OR macrofauna)


905 Abstracts were individually screened using the online software Abstrackr (<https://abstrackr.cebm.brown.edu>) to identify references matching our inclusion criteria. At the screening
 906 step, tags were given to classify studies according to the type of drivers. This step resulted
 907 in 384 articles potentially relevant for the meta-analysis, 2,152 abstracts did not match
 908 the inclusion criteria (mostly because they were not looking at both decomposition rates
 909 and decomposer communities responses to global change, or because they manipulated
 910 decomposer communities directly).

911 We refined the scope of the analysis to focus on two contrasting types of drivers for
 which we had opposing hypotheses relative to their effects: chemical stressors and nutrient
 enrichment. These two drivers had a high number of studies, were represented by aquatic
 and terrestrial studies, and had similar designs (gradients or control vs. treatment with
 increased concentrations in chemical stressors or nutrients). With the refined scope, 112

912
913
914
915
916
917
918
919
920
921
922
923

studies were potentially relevant, and 272 studies were excluded based on the tags defined at the screening step.

Figure 1 reports the PRISMA diagram describing the different steps to assemble our datasets. After full text screening of the 112 potentially relevant papers, 61 papers verified our inclusion criteria and reported data that we could extract for the meta-analysis. For the SEM analysis, 2 papers were further excluded because some data needed for the models were missing (typically the levels of nutrients or stressors).



924
925
926
927
928
929
930
931

Appendix 1 Figure 1. PRISMA plot describing the data collection steps of the meta-analysis. SEM = structural equation modelling.

Environmental quality standards

Appendix 1 Table 1. Environmental quality criteria for stressors and nutrients. Quality criteria were used to standardized the intensity levels of the different chemical stressors across studies included in the meta-analysis.

System	Chemical or Nutrient	Unit1	Unit2	Quality Criteria	citation
aquatic	fungicide: pyrimethanil	µg/l	-	0.69	Abelho, M., Martins, T. F., Shinn, C., Moreira-Santos, M. & Ribeiro, R. Effects of the fungicide pyrimethanil on biofilm and organic matter processing in outdoor lentic mesocosms. <i>Ecotoxicology</i> 25, 121–131 (2016). https://echa.europa.eu/documents/10162/41e9d7a-a-4559-f904-9cb5-0a0d5f0d6445
aquatic	fungicide: tebuconazole	µg/l	-	0.10	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.316
aquatic	As	µg/l	-	13.00	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.316

aquatic	Al	µg/l	-	87.00	https://www.govinfo.gov/content/pkg/FR-2018-12-21/pdf/2018-27745.pdf
aquatic	Cu	µg/l	-	10.10	https://echa.europa.eu/brief-profile/-/briefprofile/100.124.825
aquatic	Zn	µg/l	-	20.60	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.341
aquatic	Fe	µg/l	-	1000.00	https://www.epa.gov/wqc/national-recommended-water-quality-criteria-aquatic-life-criteria-table
aquatic	Mn	µg/l	-	1000.00	https://www.epa.gov/wqc/national-recommended-water-quality-criteria-aquatic-life-criteria-table
aquatic	Hg	µg/l	-	0.06	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.278
aquatic	Cd	µg/l	-	0.19	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.320
aquatic	insecticide: chlorpyrifos	µg/l	-	0.08	https://www.epa.gov/wqc/national-recommended-water-quality-criteria-aquatic-life-criteria-table
aquatic	phenanthrene	µg/l	-	51.40	Wu, J.-Y. et al. Development of water quality criteria for phenanthrene and comparison of the sensitivity between native and non-native species. <i>Environmental Pollution</i> 196, 141–146 (2015).
aquatic	Zn	mg/kg	-	117.80	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.341
aquatic	Cd	mg/kg	-	1.80	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.320
aquatic	Hg	mg/kg	-	9.30	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.278
aquatic	Pb	mg/kg	-	186.00	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.273
terrestrial	Cu	mg/kg	-	106.35	https://echa.europa.eu/brief-profile/-/briefprofile/100.124.825
terrestrial	Zn	mg/kg	-	35.60	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.341
terrestrial	Ni	mg/kg	-	29.90	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.283
terrestrial	Mn	mg/kg	-	3.40	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.277
terrestrial	Hg	µg/kg	-	22.00	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.278
terrestrial	Pb	mg/kg	-	212.00	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.273
terrestrial	Cd	mg/kg	-	0.90	https://echa.europa.eu/brief-profile/-/briefprofile/100.028.320
terrestrial	insecticide: chlorpyrifos	kg/ha	-	1.25	Iwai, C. B. & Noller, B. Ecotoxicological assessment of diffuse pollution using biomonitoring tool for sustainable land use in Thailand. <i>Journal of Environmental Sciences</i> 22, 858–863 (2010).

terrestrial	insecticide: endosulfan	kg/ha	-	1.25	Iwai, C. B. & Noller, B. Ecotoxicological assessment of diffuse pollution using biomonitoring tool for sustainable land use in Thailand. <i>Journal of Environmental Sciences</i> 22, 858–863 (2010).
terrestrial	herbicide: atrazine	kg/ha	-	1.88	Iwai, C. B. & Noller, B. Ecotoxicological assessment of diffuse pollution using biomonitoring tool for sustainable land use in Thailand. <i>Journal of Environmental Sciences</i> 22, 858–863 (2010).
terrestrial	insecticide: carbofuran	kg/ha	-	31.25	Iwai, C. B. & Noller, B. Ecotoxicological assessment of diffuse pollution using biomonitoring tool for sustainable land use in Thailand. <i>Journal of Environmental Sciences</i> 22, 858–863 (2010).
aquatic	pesticide mixture	arbitrary	-	1.00	Talk, A. et al. Effects of multiple but low pesticide loads on aquatic fungal communities colonizing leaf litter. <i>Journal of Environmental Sciences</i> 46, 116–125 (2016).
terrestrial	herbicide: glyphosate	kg/ha	-	4.32	European Food Safety Authority (EFSA). Conclusion on the peer review of the pesticide risk assessment of the active substance glyphosate. <i>EFSA Journal</i> 13, (2015).
terrestrial	herbicide: simazine	kg/ha	-	0.10	https://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/public/?event=activesubstance.detail&language=EN&selectedID=1853
aquatic	pesticide mixture	sum or max of TU (toxic units)	-	-3.50	Schäfer, et al., 2007. Effects of pesticides on community structure and ecosystem functions in agricultural streams of three biogeographical regions in Europe. <i>Science of The Total Environment</i> 382, 272–285.
aquatic	DIN	mg/l	N	3.05	Ministère de l'Environnement, de l'Énergie et de la Mer. Guide technique Relatif à l'évaluation de l'état des eaux de surface continentales (cours d'eau, canaux, plans d'eau). (2016).
aquatic	NH4+	mg/l	NH4	0.10	Ministère de l'Environnement, de l'Énergie et de la Mer. Guide technique Relatif à l'évaluation de l'état des eaux de surface continentales (cours d'eau, canaux, plans d'eau). (2016).
aquatic	NO3	mg/l	NO3	10.00	Ministère de l'Environnement, de l'Énergie et de la Mer. Guide technique Relatif à l'évaluation de l'état des eaux de surface continentales (cours d'eau, canaux, plans d'eau). (2016).

aquatic	NO2	mg/l	NO2	0.10	<p>Ministère de l'Environnement, de l'Énergie et de la Mer. Guide technique Relatif à l'évaluation de l'état des eaux de surface continentales (cours d'eau, canaux, plans d'eau). (2016).</p> <p>US EPA, O. Water Quality Criteria. US EPA (2013). Available at: https://www.epa.gov/wqc. (Accessed: 7th January 2019)</p> <p>Guide technique Relatif à l'évaluation de l'état des eaux de surface continentales (cours d'eau, canaux, plans d'eau). (Ministère de l'Environnement, de l'Énergie et de la Mer, 2016).</p> <p>Guide technique Relatif à l'évaluation de l'état des eaux de surface continentales (cours d'eau, canaux, plans d'eau). (Ministère de l'Environnement, de l'Énergie et de la Mer, 2016).</p> <p>Pardo et al. 2011 (Ecological Applications); derived critical loads (i.e. level of deposition below which no detrimental ecological effect occurs over the long term according to current knowledge) from empirical data for various (plant) species and ecosystems</p> <p>Amery, F. & Schoumans, O. F. Agricultural phosphorus legislation in Europe. (2014).</p>
aquatic	Total_N	mg/l	N	0.67	
aquatic	SRP	mg/l	PO43	0.10	
aquatic	Total_P	mg/l	P	0.05	
terrestrial	N deposition	kg/ha/yr	N	20.00	
terrestrial	P fertilization	kg/ha/yr	P	35.00	


932 **Appendix 2**

933

Meta-analysis

934

Publication bias



935


936

937

938

940

Appendix 2 Figure 1. Assessment of publication bias. Stressors: Funnel plots of each response variables (decomposer diversity, abundance and decomposition) in the two datasets (stressors - diversity and stressors - abundance). Meta-analytic models included the effect of stressor intensity (standardized levels) as a covariate.



941

942

943

944

945

Appendix 2 Figure 2. Assessment of publication bias. Nutrients: Funnel plots of each response variables (decomposer diversity, abundance and decomposition) in the two datasets (stressors - diversity and stressors - abundance). Meta-analytic models included the effect of nutrient intensity (standardized levels) as a covariate.

947
948
949
950

Appendix 2 Table 1. Assessment of Publication bias. Results from Egger's regressions showing the intercepts, standard error (SE) and *P*-value of regressions between standard normal deviate of each response variable (effect sizes) and the inverse of their standard errors. Models also included stressor or nutrient intensity as a covariate.

Dataset	Variable	Publication bias <i>P</i>	Publication bias	Intercept	SE
Stressors - Biodiv	Biodiversity	0.10	no	-1.36	0.83
Stressors - Biodiv	Decomposition	0.58	no	-1.07	1.94
Stressors - Abdc	Abundance	0.14	no	-1.49	1.02
Stressors - Abdc	Decomposition	0.68	no	-0.67	1.60
Nutrients - Biodiv	Biodiversity	0.37	no	0.76	0.86
Nutrients - Biodiv	Decomposition	0.19	no	3.35	2.55
Nutrients - Abdc	Abundance	0.08	no	1.21	0.70
Nutrients - Abdc	Decomposition	<.001	pub. bias	5.31	1.45

951

Meta-analysis - First level: overall mean effects

952

Appendix 2 Table 2. First level meta-analysis comparing the effects of chemical stressors and nutrient enrichment. Results of Wald-type chi-square tests comparing the grand mean effect sizes of the three response variables (decomposer diversity, abundance and litter decomposition) between chemical stressors and nutrient enrichment.

953

954

955

Response	QM	df	n	P-value
Diversity	25.647174	2	174	<0.001
Abundance	7.916468	2	424	0.019
Litter decomposition	17.611818	2	165	<0.001

956

SEM analysis

957

Appendix 2 Table 3. Summary table of structural equation modelling (SEM) analysis.

958

959

960

Unstandardized path coefficients from SEMs for the four datasets: Stressors - Biodiversity (Biodiv), Stressors - Abundance (Abdc), Nutrients - Biodiversity and Nutrients, Abundance. SEMs also incorporated categorical predictors (study type, taxonomic group and diversity metric, see Methods).

Dataset	Response	Predictor	Estimate	SE	Crit.Value	df	P-value
Stressors - Biodiv	Decomposition	Diversity	0.42	0.17	2.50	19	0.022
Stressors - Biodiv	Decomposition	Stressor intensity	-0.02	0.04	-0.47	19	0.643
Stressors - Biodiv	Diversity	Stressor intensity	-0.10	0.04	-2.44	18	0.025
Stressors - Abdc	Decomposition	Abundance	0.24	0.08	2.97	25	0.007
Stressors - Abdc	Decomposition	Stressor intensity	-0.01	0.03	-0.41	25	0.683
Stressors - Abdc	Abundance	Stressor intensity	0.00	0.05	0.03	25	0.977
Nutrients - Biodiv	Decomposition	Diversity	0.01	0.11	0.06	20	0.951
Nutrients - Biodiv	Decomposition	Nutrient intensity	-0.08	0.06	-1.21	20	0.239
Nutrients - Biodiv	Diversity	Nutrient intensity	-0.25	0.07	-3.51	19	0.002
Nutrients - Abdc	Decomposition	Abundance	0.08	0.10	0.76	44	0.451
Nutrients - Abdc	Decomposition	Nutrient intensity	-0.12	0.05	-2.16	44	0.037
Nutrients - Abdc	Abundance	Nutrient intensity	-0.06	0.06	-1.00	44	0.321

961

Meta-analysis - Second-level: categorical moderators

962

963

964

965

Appendix 2 Table 4. Main effects of categorical predictors on decomposer diversity, abundance and decomposition in the four datasets: Stressors - Biodiversity (Biodiv), Stressors - Abundance (Abdc), Nutrients - Biodiversity and Nutrients, Abundance. Results are QM statistics and associated P-values of the second-level meta-analyses.

Dataset	Response	Predictor	QM	P-value
Stressors - Biodiv	Diversity	Taxonomic group	4.80	0.028
Stressors - Abdc	Abundance	Taxonomic group	10.10	0.001
Nutrients - Biodiv	Diversity	Taxonomic group	12.77	<0.001
Nutrients - Abdc	Abundance	Taxonomic group	4.53	0.033
Stressors - Biodiv	Diversity	Study type	1.89	0.169
Stressors - Abdc	Abundance	Study type	0.92	0.338
Nutrients - Biodiv	Diversity	Study type	0.24	0.625
Nutrients - Abdc	Abundance	Study type	0.98	0.323
Stressors - Biodiv	Diversity	Diversity metric	1.67	0.196
Nutrients - Biodiv	Diversity	Diversity metric	2.35	0.125
Stressors - Biodiv	Decomposition	Study type	0.16	0.693
Stressors - Abdc	Decomposition	Study type	1.85	0.174
Nutrients - Biodiv	Decomposition	Study type	2.69	0.101
Nutrients - Abdc	Decomposition	Study type	0.18	0.674

966 **Appendix 3**967 **Sensitivity analyses**968 **Influence of approximating standard deviations**

969 When studies did not report standard deviations associated with the mean decomposer
 970 diversity or abundance or the mean decomposition rates, we used linear approximations to
 971 estimate the variance based on our data (see Methods). We tested the influence of those
 972 approximations on the final results by running the structural equation modelling (SEM)
 973 analysis without those effect sizes for which standard deviations were approximated. Overall
 974 the same patterns were found showing that approximating missing standard deviations had
 975 limited effects on the final SEM results.

976 **Appendix 3 Table 1. Results of mediation tests from structural equation modelling (SEM)**
 977 **analysis based on data without approximated standard deviations.** C statistic and associated
 978 *P*-value for SEM without the path from biodiversity or abundance to decomposition for the four
 979 datasets: Stressors - Diversity, Stressors - Abundance, Nutrients - Diversity and Nutrients - Abundance.
 980 Δ -AIC is the difference in AIC score between models with and without biodiversity- or
 981 abundance-mediated effects.

Dataset	C statistic	df	<i>P</i> -value	Δ -AIC	no.studies	n
Stressors, Biodiv	12.42	6	0.053	-8.32	16	58
Stressors, Abdc	10.15	4	0.038	-6.82	23	216
Nutrient, Biodiv	13.33	6	0.038	-1.46	21	67
Nutrient, Abdc	3.82	4	0.432	-0.12	32	127

982
983
984

Appendix 3 Table 2. Summary table of structural equation modelling (SEM) analysis based on data without approximated standard deviations. Standardized (Std.Est.) and unstandardized (Estimate) path coefficients from SEMs for the four datasets.

Dataset	Response	Predictor	Std.Est.	Estimate	SE	Crit.Value	df	P-value
Stressors - Biodiv	Decomposition	Diversity	0.52	0.50	0.16	3.16	12	0.008
Stressors - Biodiv	Decomposition	Stressor intensity	-0.26	-0.05	0.03	-1.54	12	0.148
Stressors - Biodiv	Diversity	Stressor intensity	-0.39	-0.08	0.04	-1.89	11	0.085
Stressors - Abdc	Decomposition	Abundance	0.40	0.27	0.09	2.91	19	0.009
Stressors - Abdc	Decomposition	Stressor intensity	-0.11	-0.02	0.03	-0.77	19	0.450
Stressors - Abdc	Abundance	Stressor intensity	0.08	0.03	0.06	0.46	19	0.649
Nutrients - Biodiv	Decomposition	Diversity	-0.04	-0.04	0.12	-0.35	10	0.732
Nutrients - Biodiv	Decomposition	Nutrient intensity	-0.31	-0.14	0.09	-1.52	10	0.161
Nutrients - Biodiv	Diversity	Nutrient intensity	-0.49	-0.23	0.10	-2.39	9	0.040
Nutrients - Abdc	Decomposition	Abundance	0.05	0.04	0.13	0.33	29	0.742
Nutrients - Abdc	Decomposition	Nutrient intensity	-0.26	-0.12	0.06	-1.91	29	0.066
Nutrients - Abdc	Abundance	Nutrient intensity	-0.20	-0.10	0.07	-1.40	29	0.173

985
986
987
988
989
990
991
992
993
994

Influence of extreme values

We re-run our SEMs with datasets excluding extreme values of effect sizes. Extreme values were defined as values exceeding the whiskers of boxplots. Overall we found similar patterns showing that extreme effect sizes had limited effects on the final SEM results.

Appendix 3 Table 3. Results of mediation tests from structural equation modelling (SEM) analysis based on data excluding extreme values of effect sizes. C statistic and associated *P*-value for SEM without the path from biodiversity or abundance to decomposition for the four datasets: Stressors - Diversity, Stressors - Abundance, Nutrients - Diversity and Nutrients - Abundance. Δ -AIC is the difference in AIC score between models with and without biodiversity- or abundance-mediated effects.

Dataset	C statistic	df	<i>P</i> -value	Δ -AIC	no.studies	n
Stressors, Biodiv	10.18	6	0.117	-6.71	22	94
Stressors, Abdc	7.39	4	0.117	-4.23	27	254
Nutrient, Biodiv	14.80	6	0.022	-4.85	26	93
Nutrient, Abdc	2.74	4	0.603	0.15	35	159

995
996
997

Appendix 3 Table 4. Summary table of structural equation modelling (SEM) analysis based on data excluding extreme values of effect sizes. Standardized (Std.Est.) and unstandardized (Estimate) path coefficients from SEMs for the four datasets.

Dataset	Response	Predictor	Std.Est.	Estimate	SE	Crit.Value	df	P-value
Stressors - Biodiv	Decomposition	Diversity	0.41	0.40	0.18	2.20	18	0.041
Stressors - Biodiv	Decomposition	Stressor intensity	-0.04	-0.01	0.04	-0.24	18	0.814
Stressors - Biodiv	Diversity	Stressor intensity	-0.44	-0.10	0.04	-2.75	17	0.014
Stressors - Abdc	Decomposition	Abundance	0.30	0.24	0.11	2.24	23	0.035
Stressors - Abdc	Decomposition	Stressor intensity	0.05	0.01	0.03	0.35	23	0.731
Stressors - Abdc	Abundance	Stressor intensity	0.00	0.00	0.04	-0.02	23	0.980
Nutrients - Biodiv	Decomposition	Diversity	0.00	0.00	0.11	0.02	19	0.986
Nutrients - Biodiv	Decomposition	Nutrient intensity	-0.18	-0.08	0.06	-1.30	19	0.210
Nutrients - Biodiv	Diversity	Nutrient intensity	-0.53	-0.24	0.07	-3.36	18	0.003
Nutrients - Abdc	Decomposition	Abundance	0.00	0.00	0.09	0.04	37	0.968
Nutrients - Abdc	Decomposition	Nutrient intensity	-0.38	-0.13	0.04	-3.26	37	0.002
Nutrients - Abdc	Abundance	Nutrient intensity	-0.24	-0.09	0.05	-1.73	37	0.092

Influence of the effect size metric

We tested the influence of the metric of effect size selected on the results of the SEMs. Log-response ratios were calculated instead of correlation coefficients and the models were re-run based on those data. The results were partially different from the original analysis. For nutrients, similar patterns were found, however for stressors there was limited support for the biodiversity- and abundance-mediated effects on decomposition responses. We noted extreme values of log-response ratios that may have explained such patterns. Besides, the log-response ratio has a different interpretation compared to correlation coefficients. Log-response ratios are sensitive to the different metrics of diversity and abundance, taxa groups, litter types etc. used across studies included in this meta-analysis. Therefore this result reinforced our choice of correlation coefficients as relevant effect sizes in the present meta-analysis.

Appendix 3 Table 5. Results of mediation tests from structural equation modelling (SEM) analysis based on data using log-response ratio as an effect size. C statistic and associated *P*-value for SEM without the path from biodiversity or abundance to decomposition for the four datasets: Stressors - Diversity, Stressors - Abundance, Nutrients - Diversity and Nutrients - Abundance. Δ -AIC is the difference in AIC score between models with and without biodiversity- or abundance-mediated effects.

Dataset	C statistic	df	<i>P</i> -value	Δ -AIC	no.studies	n
Stressors, Biodiv	4.11	6	0.662	-0.02	22	70
Stressors, Abdc	5.59	4	0.232	-2.22	37	150
Nutrient, Biodiv	8.03	6	0.236	-2.08	14	78
Nutrient, Abdc	3.41	4	0.492	-0.44	21	307

Appendix 3 Table 6. Summary table of structural equation modelling (SEM) analysis based on data using log-response ratio as an effect size. Standardized (Std.Est.) and unstandardized (Estimate) path coefficients from SEMs for the four datasets.

Dataset	Response	Predictor	Std.Est	Estimate	SE	Crit.Value	df	<i>P</i> .value
Stressors - Biodiv	Decomposition	Diversity	0.18	0.12	0.15	0.80	15	0.437
Stressors - Biodiv	Decomposition	Stressor intensity	-0.24	-0.05	0.04	-1.47	15	0.163
Stressors - Biodiv	Diversity	Stressor intensity	-0.35	-0.12	0.03	-4.17	15	0.001
Stressors - Abdc	Decomposition	Abundance	0.14	0.04	0.05	0.86	28	0.396
Stressors - Abdc	Decomposition	Stressor intensity	0.09	0.02	0.04	0.55	28	0.586
Stressors - Abdc	Abundance	Stressor intensity	-0.14	-0.11	0.11	-1.03	28	0.312
Nutrients - Biodiv	Decomposition	Diversity	0.29	0.19	0.10	1.80	14	0.094
Nutrients - Biodiv	Decomposition	Nutrient intensity	-0.15	-0.07	0.08	-0.96	14	0.352
Nutrients - Biodiv	Diversity	Nutrient intensity	-0.20	-0.16	0.07	-2.11	14	0.054
Nutrients - Abdc	Decomposition	Abundance	0.06	0.04	0.06	0.59	42	0.559
Nutrients - Abdc	Decomposition	Nutrient intensity	-0.36	-0.16	0.05	-3.08	42	0.004
Nutrients - Abdc	Abundance	Nutrient intensity	-0.01	0.00	0.08	-0.08	42	0.935

1019 **Appendix 4**1020 **References included in the meta-analysis**1021 **References**

- 1022 **Abelho M**, Martins TF, Shinn C, Moreira-Santos M, Ribeiro R. Effects of the Fungicide Pyrimethanil
1023 on Biofilm and Organic Matter Processing in Outdoor Lentic Mesocosms. *Ecotoxicology*. 2016 Jan;
1024 25(1):121–131. doi: 10.1007/s10646-015-1574-x.
- 1025 **Allison SD**, Gartner TB, Mack MC, McGuire K, Treseder K. Nitrogen Alters Carbon Dynamics during
1026 Early Succession in Boreal Forest. *Soil Biology and Biochemistry*. 2010 Jul; 42(7):1157–1164. doi:
1027 10.1016/j.soilbio.2010.03.026.
- 1028 **Artigas J**, Majerholc J, Foulquier A, Margoum C, Volat B, Neyra M, Pesce S. Effects of the Fungicide
1029 Tebuconazole on Microbial Capacities for Litter Breakdown in Streams. *Aquatic Toxicology*. 2012 Oct;
1030 122-123(Supplement C):197–205. doi: 10.1016/j.aquatox.2012.06.011.
- 1031 **Bergfur J**, Johnson RK, Sandin L, Goedkoop W, Nygren K. Effects of Nutrient Enrichment on Boreal
1032 Streams: Invertebrates, Fungi and Leaf-Litter Breakdown. *Freshwater Biology*. 2007 Aug; 52(8):1618–
1033 1633. doi: 10.1111/j.1365-2427.2007.01770.x.
- 1034 **Bott TL**, Jackson JK, McTammany ME, Newbold JD, Rier ST, Sweeney BW, Battle JM. Abandoned Coal
1035 Mine Drainage and Its Remediation: Impacts on Stream Ecosystem Structure and Function. *Ecological*
1036 *Applications*. 2012; 22(8):2144–2163.
- 1037 **Brosed M**, Lamothe S, Chauvet E. Litter Breakdown for Ecosystem Integrity Assessment Also Applies to
1038 Streams Affected by Pesticides. *Hydrobiologia*. 2016 Jun; 773(1):87–102. doi: 10.1007/s10750-016-
1039 2681-2.
- 1040 **Cabrini R**, Canobbio S, Sartori L, Fornaroli R, Mezzanotte V. Leaf Packs in Impaired Streams: The
1041 Influence of Leaf Type and Environmental Gradients on Breakdown Rate and Invertebrate Assemblage
1042 Composition. *Water, Air, & Soil Pollution*. 2013 Oct; 224(10). doi: 10.1007/s11270-013-1697-8.
- 1043 **Carlisle DM**, Clements WH. Leaf Litter Breakdown, Microbial Respiration and Shredder Production in
1044 Metal-Polluted Streams: Litter Breakdown in Metal-Polluted Streams. *Freshwater Biology*. 2005 Jan;
1045 50(2):380–390. doi: 10.1111/j.1365-2427.2004.01323.x.
- 1046 **Chaffin JL**, Valett HM, Webster JR, Schreiber ME. Influence of Elevated As on Leaf Breakdown in an
1047 Appalachian Headwater Stream. *Journal of the North American Benthological Society*. 2005 Sep;
1048 24(3):553–568. doi: 10.1899/04-058.1.
- 1049 **Chung N**, Suberkropp K. Influence of Shredder Feeding and Nutrients on Fungal Activity and Community
1050 Structure in Headwater Streams. *Fundamental and Applied Limnology / Archiv für Hydrobiologie*.
1051 2008 Oct; 173(1):35–46. doi: 10.1127/1863-9135/2008/0173-0035.
- 1052 **Creamer RE**, Rimmer DL, Black HJ. Do Elevated Soil Concentrations of Metals Affect the Diversity and
1053 Activity of Soil Invertebrates in the Long-Term? *Soil Use and Management*. 2008 Mar; 24(1):37–46.
1054 doi: 10.1111/j.1475-2743.2007.00131.x.
- 1055 **Duarte S**, Pascoal C, Cássio F. Effects of Zinc on Leaf Decomposition by Fungi in Streams: Studies in
1056 Microcosms. *Microbial Ecology*. 2004 Nov; 48(3):366–374. doi: 10.1007/s00248-003-2032-5.
- 1057 **Duarte S**, Pascoal C, Garabetian F, Cassio F, Charcosset JY. Microbial Decomposer Communities Are
1058 Mainly Structured by Trophic Status in Circumneutral and Alkaline Streams. *Applied and Environmen-
1059 tal Microbiology*. 2009 Oct; 75(19):6211–6221. doi: 10.1128/AEM.00971-09.
- 1060 **Duarte S**, Pascoal C, Cássio F. High Diversity of Fungi May Mitigate the Impact of Pollution on Plant
1061 Litter Decomposition in Streams. *Microbial Ecology*. 2008 Nov; 56(4):688–695. doi: 10.1007/s00248-
1062 008-9388-5.
- 1063 **Duarte S**, Pascoal C, Cássio F. Functional Stability of Stream-Dwelling Microbial Decomposers Exposed
1064 to Copper and Zinc Stress. *Freshwater Biology*. 2009 Aug; 54(8):1683–1691. doi: 10.1111/j.1365-
1065 2427.2009.02217.x.

- 1066 **Dunck B**, Lima-Fernandes E, Cássio F, Cunha A, Rodrigues L, Pascoal C. Responses of Primary Production,
1067 Leaf Litter Decomposition and Associated Communities to Stream Eutrophication. *Environmental*
1068 *Pollution*. 2015 Jul; 202(Supplement C):32–40. doi: [10.1016/j.envpol.2015.03.014](https://doi.org/10.1016/j.envpol.2015.03.014).
- 1069 **Englert D**, Zubrod JP, Schulz R, Bundschuh M. Effects of Municipal Wastewater on Aquatic Ecosystem
1070 Structure and Function in the Receiving Stream. *Science of The Total Environment*. 2013 Jun; 454-
1071 455(Supplement C):401–410. doi: [10.1016/j.scitotenv.2013.03.025](https://doi.org/10.1016/j.scitotenv.2013.03.025).
- 1072 **Estevez E**, Rodríguez-Castillo T, Álvarez-Cabria M, Peñas FJ, González-Ferreras AM, Lezcano M, Barquín J.
1073 Analysis of Structural and Functional Indicators for Assessing the Health State of Mountain Streams.
1074 *Ecological Indicators*. 2017 Jan; 72(Supplement C):553–564. doi: [10.1016/j.ecolind.2016.08.052](https://doi.org/10.1016/j.ecolind.2016.08.052).
- 1075 **Fanin N**, Hättenschwiler S, Schimann H, Fromin N. Interactive Effects of C, N and P Fertilization on Soil
1076 Microbial Community Structure and Function in an Amazonian Rain Forest. *Functional Ecology*. 2015
1077 Jan; 29(1):140–150. doi: [10.1111/1365-2435.12329](https://doi.org/10.1111/1365-2435.12329).
- 1078 **Fernandes I**, Duarte S, Cássio F, Pascoal C. Mixtures of Zinc and Phosphate Affect Leaf Litter Decompo-
1079 sition by Aquatic Fungi in Streams. *Science of The Total Environment*. 2009 Jul; 407(14):4283–4288.
1080 doi: [10.1016/j.scitotenv.2009.04.007](https://doi.org/10.1016/j.scitotenv.2009.04.007).
- 1081 **Fernández D**, Voss K, Bundschuh M, Zubrod JP, Schäfer RB. Effects of Fungicides on Decomposer
1082 Communities and Litter Decomposition in Vineyard Streams. *Science of The Total Environment*. 2015
1083 Nov; 533(Supplement C):40–48. doi: [10.1016/j.scitotenv.2015.06.090](https://doi.org/10.1016/j.scitotenv.2015.06.090).
- 1084 **Ferreira V**, Gulis V, Graça MAS. Whole-Stream Nitrate Addition Affects Litter Decomposition and
1085 Associated Fungi but Not Invertebrates. *Oecologia*. 2006 Oct; 149(4):718–729. doi: [10.1007/s00442-](https://doi.org/10.1007/s00442-006-0478-0)
1086 [006-0478-0](https://doi.org/10.1007/s00442-006-0478-0).
- 1087 **Gan H**, Zak DR, Hunter MD. Chronic Nitrogen Deposition Alters the Structure and Function of Detrital
1088 Food Webs in a Northern Hardwood Ecosystem. *Ecological Applications*. 2013 Sep; 23(6):1311–1321.
1089 doi: [10.1890/12-1895.1](https://doi.org/10.1890/12-1895.1).
- 1090 **Gulis V**, Ferreira V, Graca MAS. Stimulation of Leaf Litter Decomposition and Associated Fungi and
1091 Invertebrates by Moderate Eutrophication: Implications for Stream Assessment. *Freshwater Biology*.
1092 2006 Sep; 51(9):1655–1669. doi: [10.1111/j.1365-2427.2006.01615.x](https://doi.org/10.1111/j.1365-2427.2006.01615.x).
- 1093 **Gulis V**, Suberkropp K. Leaf Litter Decomposition and Microbial Activity in Nutrient-Enriched and
1094 Unaltered Reaches of a Headwater Stream. *Freshwater biology*. 2003; 48(1):123–134.
- 1095 **Hagen EM**, Webster JR, Benfield EF. Are Leaf Breakdown Rates a Useful Measure of Stream Integrity
1096 along an Agricultural Landuse Gradient? *Journal of the North American Benthological Society*. 2006
1097 Jun; 25(2):330–343. doi: [10.1899/0887-3593\(2006\)25\[330:ALBRAU\]2.0.CO;2](https://doi.org/10.1899/0887-3593(2006)25[330:ALBRAU]2.0.CO;2).
- 1098 **Hobbelen PHF**, van den Brink PJ, Hobbelen JF, van Gestel CAM. Effects of Heavy Metals on the Structure
1099 and Functioning of Detritivore Communities in a Contaminated Floodplain Area. *Soil Biology and*
1100 *Biochemistry*. 2006 Jul; 38(7):1596–1607. doi: [10.1016/j.soilbio.2005.11.010](https://doi.org/10.1016/j.soilbio.2005.11.010).
- 1101 **Hobbie SE**, Eddy WC, Buyarski CR, Adair EC, Ogdahl ML, Weisenhorn P. Response of Decomposing Litter
1102 and Its Microbial Community to Multiple Forms of Nitrogen Enrichment. *Ecological Monographs*.
1103 2012; 82(3):389–405.
- 1104 **Hogsden KL**, Harding JS. Leaf Breakdown, Detrital Resources, and Food Webs in Streams Affected by
1105 Mine Drainage. *Hydrobiologia*. 2013 Oct; 716(1):59–73. doi: [10.1007/s10750-013-1544-3](https://doi.org/10.1007/s10750-013-1544-3).
- 1106 **Hopkins JM**, Marcarelli AM, Bechtold HA. Ecosystem Structure and Function Are Complementary
1107 Measures of Water Quality in a Polluted, Spring-Influenced River. *Water, Air, & Soil Pollution*. 2011
1108 Jan; 214(1-4):409–421. doi: [10.1007/s11270-010-0432-y](https://doi.org/10.1007/s11270-010-0432-y).
- 1109 **Iwai CB**, Noller B. Ecotoxicological Assessment of Diffuse Pollution Using Biomonitoring Tool for
1110 Sustainable Land Use in Thailand. *Journal of Environmental Sciences*. 2010 Jun; 22(6):858–863. doi:
1111 [10.1016/S1001-0742\(09\)60189-3](https://doi.org/10.1016/S1001-0742(09)60189-3).

- 1112 **Johnson KS**, Thompson PC, Gromen L, Bowman J. Use of Leaf Litter Breakdown and Macroinvertebrates
1113 to Evaluate Gradient of Recovery in an Acid Mine Impacted Stream Remediated with an Active Alkaline
1114 Doser. *Environmental Monitoring and Assessment*. 2014 Jul; 186(7):4111–4127. doi: 10.1007/s10661-
1115 014-3684-y.
- 1116 **Lecerf A**, Usseglio-Polatera P, Charcosset JY, Lambrigot D, Bracht B, Chauvet E. Assessment of Functional
1117 Integrity of Eutrophic Streams Using Litter Breakdown and Benthic Macroinvertebrates. *Archiv für*
1118 *Hydrobiologie*. 2006 Jan; 165(1):105–126. doi: 10.1127/0003-9136/2006/0165-0105.
- 1119 **Lima-Fernandes E**, Fernandes I, Pereira A, Geraldes P, Cássio F, Pascoal C. Eutrophication Modulates
1120 Plant-Litter Diversity Effects on Litter Decomposition in Streams. *Freshwater Science*. 2015 Mar;
1121 34(1):31–41. doi: 10.1086/679223.
- 1122 **Liu P**, Huang J, Sun OJ, Han X. Litter Decomposition and Nutrient Release as Affected by Soil Nitrogen
1123 Availability and Litter Quality in a Semiarid Grassland Ecosystem. *Oecologia*. 2010 Mar; 162(3):771–780.
1124 doi: 10.1007/s00442-009-1506-7.
- 1125 **Lopes ML**, Rodrigues AM, Quintino V. Can the Leaf-Bag Technique Detect Benthic Macrofauna Responses
1126 to Sediment Contamination by Metals and Metalloids in Estuaries? *Marine Pollution Bulletin*. 2017
1127 Nov; 124(1):171–180. doi: 10.1016/j.marpolbul.2017.07.027.
- 1128 **Lucisine P**, Lecerf A, Danger M, Felten V, Aran D, Auclerc A, Gross EM, Huot H, Morel JL, Muller S,
1129 Nahmani J, Maunoury-Danger F. Litter Chemistry Prevails over Litter Consumers in Mediating Effects
1130 of Past Steel Industry Activities on Leaf Litter Decomposition. *Science of The Total Environment*. 2015
1131 Dec; 537(Supplement C):213–224. doi: 10.1016/j.scitotenv.2015.07.112.
- 1132 **Matthaei CD**, Piggott JJ, Townsend CR. Multiple Stressors in Agricultural Streams: Interactions among
1133 Sediment Addition, Nutrient Enrichment and Water Abstraction: Sediment, Nutrients & Water Abstrac-
1134 tion. *Journal of Applied Ecology*. 2010 Apr; 47(3):639–649. doi: 10.1111/j.1365-2664.2010.01809.x.
- 1135 **Mederiros AO**, Callisto M, Graça MAS, Ferreira F Veronica, Rosa C, Franca J, Eller A, Rezende R,
1136 Gonçalves Jr JF. Microbial Colonisation and Litter Decomposition in a Cerrado Stream Are Limited by
1137 Low Dissolved Nutrient Concentrations. *Limnetica*. 2015; 34(2):283–292.
- 1138 **Menéndez M**. Response of Early *Ruppia Cirrhosa* Litter Breakdown to Nutrient Addition in a Coastal
1139 Lagoon Affected by Agricultural Runoff. *Estuarine, Coastal and Shelf Science*. 2009 May; 82(4):608–614.
1140 doi: 10.1016/j.ecss.2009.02.029.
- 1141 **Mesquita A**, Pascoal C, Cássio F. Assessing Effects of Eutrophication in Streams Based on Breakdown
1142 of Eucalypt Leaves. *Fundamental and Applied Limnology / Archiv für Hydrobiologie*. 2007 Mar;
1143 168(3):221–230. doi: 10.1127/1863-9135/2007/0168-0221.
- 1144 **Milcu A**, Heim A, Ellis RJ, Scheu S, Manning P. Identification of General Patterns of Nutrient and Labile
1145 Carbon Control on Soil Carbon Dynamics Across a Successional Gradient. *Ecosystems*. 2011 Aug;
1146 14(5):710–719. doi: 10.1007/s10021-011-9440-z.
- 1147 **Moreirinha C**, Duarte S, Pascoal C, Cássio F. Effects of Cadmium and Phenanthrene Mixtures on Aquatic
1148 Fungi and Microbially Mediated Leaf Litter Decomposition. *Archives of Environmental Contamination*
1149 *and Toxicology*. 2011 Aug; 61(2):211–219. doi: 10.1007/s00244-010-9610-6.
- 1150 **Münze R**, Orlinskiy P, Gunold R, Paschke A, Kaske O, Beketov MA, Hundt M, Bauer C, Schüürmann
1151 G, Möder M, Liess M. Pesticide Impact on Aquatic Invertebrates Identified with Chemcatcher®
1152 Passive Samplers and the SPEARpesticides Index. *Science of The Total Environment*. 2015 Dec;
1153 537(Supplement C):69–80. doi: 10.1016/j.scitotenv.2015.07.012.
- 1154 **Niyogi DK**, Cheatham CA, Thomson WH, Christiansen JM. Litter Breakdown and Fungal Diversity in a
1155 Stream Affected by Mine Drainage. *Fundamental and Applied Limnology / Archiv für Hydrobiologie*.
1156 2009 Jun; 175(1):39–48. doi: 10.1127/1863-9135/2009/0175-0039.
- 1157 **Niyogi DK**, Lewis WM, McKnight DM. Litter Breakdown in Mountain Streams Affected by Mine Drainage:
1158 Biotic Mediation of Abiotic Controls. *Ecological Applications*. 2001; 11(2):506–516. doi: 10.1890/1051-
1159 0761(2001)011[0506:LBIMSA]2.0.CO;2.

- 1160 **Ouédraogo E**, Mando A, Brussaard L. Soil Macrofaunal-Mediated Organic Resource Disap-
1161 pearance in Semi-Arid West Africa. *Applied Soil Ecology*. 2004 Nov; 27(3):259–267. doi:
1162 [10.1016/j.apsoil.2004.03.003](https://doi.org/10.1016/j.apsoil.2004.03.003).
- 1163 **Pascoal C**, Pinho M, Cássio F, Gomes P. Assessing Structural and Functional Ecosystem Condition Using
1164 Leaf Breakdown: Studies on a Polluted River. *Freshwater Biology*. 2003; 48(11):2033–2044.
- 1165 **Pearson RG**, Connolly NM. Nutrient Enhancement, Food Quality and Community Dynamics in a Tropical
1166 Rainforest Stream. *Freshwater Biology*. 2000; 43(1):31–42.
- 1167 **Pérez J**, Basaguren A, Descals E, Larrañaga A, Pozo J. Leaf-Litter Processing in Headwater Streams of
1168 Northern Iberian Peninsula: Moderate Levels of Eutrophication Do Not Explain Breakdown Rates.
1169 *Hydrobiologia*. 2013 Nov; 718(1):41–57. doi: [10.1007/s10750-013-1610-x](https://doi.org/10.1007/s10750-013-1610-x).
- 1170 **Perkiömäki J**, Kiikkilä O, Moilanen M, Issakainen J, Tervahauta A, Fritze H. Cadmium-Containing Wood
1171 Ash in a Pine Forest: Effects on Humus Microflora and Cadmium Concentrations in Mushrooms,
1172 Berries, and Needles. *Canadian Journal of Forest Research*. 2003; 33(12):2443–2451. doi: [10.1139/x03-
1173 169](https://doi.org/10.1139/x03-169).
- 1174 **Piscart C**, Genoel R, Doledec S, Chauvet E, Marmonier P. Effects of Intense Agricultural Practices on
1175 Heterotrophic Processes in Streams. *Environmental Pollution*. 2009 Mar; 157(3):1011–1018. doi:
1176 [10.1016/j.envpol.2008.10.010](https://doi.org/10.1016/j.envpol.2008.10.010).
- 1177 **Quintão JMB**, Rezende RS, Júnior JFG. Microbial Effects in Leaf Breakdown in Tropical Reservoirs of
1178 Different Trophic Status. *Freshwater Science*. 2013 Jul; 32(3):933–950. doi: [10.1899/12-112.1](https://doi.org/10.1899/12-112.1).
- 1179 **Schäfer RB**, Caquet T, Siimes K, Mueller R, Lagadic L, Liess M. Effects of Pesticides on Community Struc-
1180 ture and Ecosystem Functions in Agricultural Streams of Three Biogeographical Regions in Europe.
1181 *Science of The Total Environment*. 2007 Sep; 382(2):272–285. doi: [10.1016/j.scitotenv.2007.04.040](https://doi.org/10.1016/j.scitotenv.2007.04.040).
- 1182 **Schultheis AS**, Sanchez M, Hendricks AC. Structural and Functional Responses of Stream Insects to
1183 Copper Pollution. *Hydrobiologia*. 1997; 346(1):85–93.
- 1184 **Shaftel RS**, King RS, Back JA. Breakdown Rates, Nutrient Concentrations, and Macroinvertebrate
1185 Colonization of Bluejoint Grass Litter in Headwater Streams of the Kenai Peninsula, Alaska. *Journal of
1186 the North American Benthological Society*. 2011 Jun; 30(2):386–398. doi: [10.1899/10-086.1](https://doi.org/10.1899/10-086.1).
- 1187 **Talk A**, Kublik S, Uksa M, Engel M, Berghahn R, Welzl G, Schloter M, Mohr S. Effects of Multiple but
1188 Low Pesticide Loads on Aquatic Fungal Communities Colonizing Leaf Litter. *Journal of Environmental
1189 Sciences*. 2016 Aug; 46(Supplement C):116–125. doi: [10.1016/j.jes.2015.11.028](https://doi.org/10.1016/j.jes.2015.11.028).
- 1190 **Thompson MSA**, Bankier C, Bell T, Dumbrell AJ, Gray C, Ledger ME, Lehmann K, McKew BA, Sayer
1191 CD, Shelley F, Trimmer M, Warren SL, Woodward G. Gene-to-Ecosystem Impacts of a Catastrophic
1192 Pesticide Spill: Testing a Multilevel Bioassessment Approach in a River Ecosystem. *Freshwater Biology*.
1193 2016 Dec; 61(12):2037–2050. doi: [10.1111/fwb.12676](https://doi.org/10.1111/fwb.12676).
- 1194 **Wardle DA**, Yeates GW, Bonner KI, Nicholson KS, Watson RN. Impacts of Ground Vegetation Management
1195 Strategies in a Kiwifruit Orchard on the Composition and Functioning of the Soil Biota. *Soil Biology
1196 and Biochemistry*. 2001 Jun; 33(7):893–905. doi: [10.1016/S0038-0717\(00\)00235-2](https://doi.org/10.1016/S0038-0717(00)00235-2).
- 1197 **Woodcock TS**, Huryn AD. Leaf Litter Processing and Invertebrate Assemblages along a Pollution
1198 Gradient in a Maine (USA) Headwater Stream. *Environmental Pollution*. 2005 Apr; 134(3):363–375.
1199 doi: [10.1016/j.envpol.2004.09.020](https://doi.org/10.1016/j.envpol.2004.09.020).
- 1200 **Woodward G**, Gessner MO, Giller PS, Gulis V, Hladyz S, Lecerf A, Malmqvist B, McKie BG, Tiegs SD,
1201 Cariss H, Dobson M, Elosegi A, Ferreira V, Graça MAS, Fleituch T, Lacoursière JO, Nistorescu M, Pozo J,
1202 Risnoveanu G, Schindler M, et al. Continental-Scale Effects of Nutrient Pollution on Stream Ecosystem
1203 Functioning. *Science*. 2012 Jun; 336(6087):1438–1440. doi: [10.1126/science.1219534](https://doi.org/10.1126/science.1219534).