

## RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

### Hypothesis testing for tail dependence parameters on the boundary of the parameter space

Kiriliouk, Anna

*Published in:*  
Econometrics and Statistics

*DOI:*  
[10.1016/j.ecosta.2019.06.001](https://doi.org/10.1016/j.ecosta.2019.06.001)

*Publication date:*  
2020

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication](#)

*Citation for pulished version (HARVARD):*  
Kiriliouk, A 2020, 'Hypothesis testing for tail dependence parameters on the boundary of the parameter space', *Econometrics and Statistics*, vol. 16, pp. 121-135. <https://doi.org/10.1016/j.ecosta.2019.06.001>


#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Hypothesis testing for tail dependence parameters on the boundary of the parameter space

Anna Kiriliouk

Université de Namur, Faculté des sciences économiques, sociales et de gestion, Rempart de la vierge 8, Namur B-5000, Belgium

## ARTICLE INFO

### Article history:

Received 25 January 2018

Revised 19 June 2019

Accepted 24 June 2019

Available online 29 July 2019

### Keywords:

Brown–Resnick model

Hypothesis testing

Max-linear model

Multivariate extremes

Stable tail dependence function

Tail dependence

## ABSTRACT

Modelling multivariate tail dependence is one of the key challenges in extreme-value theory. Multivariate extremes are usually characterized using parametric models, some of which have simpler submodels at the boundary of their parameter space. Hypothesis tests are proposed for tail dependence parameters that, under the null hypothesis, are on the boundary of the alternative hypothesis. The asymptotic distribution of the weighted least squares estimator is given when the true parameter vector is on the boundary of the parameter space, and two test statistics are proposed. The performance of these test statistics is evaluated for the Brown–Resnick model and the max-linear model. In particular, simulations show that it is possible to recover the optimal number of factors for a max-linear model. Finally, the methods are applied to characterize the dependence structure of two major stock market indices, the DAX and the CAC40.

© 2019 EcoSta Econometrics and Statistics. Published by Elsevier B.V. All rights reserved.

## 1. Introduction

Extreme-value theory is the branch of statistics concerned with the characterization of extreme events. These occur in a large variety of fields, such as hydrology, meteorology, finance and insurance, but also in settings like human life span or athletic records. For some examples, see Einmahl and Magnus (2008), Towler et al. (2010), Chavez-Demoulin et al. (2016), Thomas et al. (2016) and Rootzén and Zholud (2017). In the univariate case, the limiting distribution of suitably normalized block maxima or threshold exceedances can be characterized entirely using the generalized extreme-value (Fisher and Tippett, 1928; Gnedenko, 1943) and the generalized Pareto distribution (Balkema and De Haan, 1974; Pickands et al., 1975) respectively. However, many extreme events are inherently multivariate, and an important challenge is to model the tail dependence between two or more random variables of interest. If dependence disappears as the variables take on more and more extreme values, we say that they are asymptotically independent. Testing for asymptotic independence and modelling asymptotically independent data has been done in Ledford and Tawn (1996), Draisma et al. (2004), Hüsler and Li (2009), Wadsworth et al. (2016) and Guillou et al. (2018), among others. In the following, we consider the framework of asymptotic dependence.

The family of limiting distributions for multivariate maxima or threshold exceedances is infinite-dimensional, so that realistic and computationally feasible parametric models need to be proposed. Examples include Gumbel (1960), Smith (1990), Tawn (1990) and Kabluchko et al. (2009). Many different approaches of estimating tail dependence parameters exist. Fitting a multivariate extreme-value distribution to componentwise block maxima is done in Padoan et al. (2010), Davison et al. (2012), Castruccio et al. (2016) and Dombry et al. (2017), along others. Multivariate threshold exceedances, which are the

E-mail address: [anna.kiriliouk@unamur.be](mailto:anna.kiriliouk@unamur.be)

focus of this work, can be tackled using the stable tail dependence function (Drees and Huang, 1998) or the multivariate generalized Pareto distribution (Rootzén and Tajvidi, 2006). Common estimation methods include maximum likelihood (Wadsworth and Tawn, 2014; Kiriliouk et al., 2018a; de Fondeville and Davison, 2018) and minimum distance estimation (Einmahl et al., 2012; 2016; 2018).

Some popular parametric models have simpler submodels at the boundary of the parameter space. The main goal of this paper is to propose hypothesis tests for tail dependence parameters that, under the null hypothesis, are on the boundary of the alternative hypothesis. A first example is the Brown–Resnick model (Kablichko et al., 2009), popular in spatial extremes, which reduces to the simpler Smith model (Smith, 1990) when the shape parameter of the former attains its upper bound. A second example is the max-linear model, where each component of a vector can be interpreted as the maximum shock among a set of independent factors. Recent work on max-linear models is numerous. In Gissibl and Klüppelberg (2018) and Gissibl et al. (2018), a recursive max-linear model on a directed acyclic graph is considered. In Cui and Zhang (2018) and Zhao and Zhang (2018), max-linear models are used to model a latent factor structure for financial returns. Finally, the Marshall–Olkin model (Embrechts et al., 2003; Segers, 2012) is a submodel that is frequently used in practice, see for instance Burtschell et al., (2009), Su and Furman, (2017) or Brigo et al., (2018).

Because of their non-differentiability, max-linear models cannot be estimated by standard likelihood methods and hence we use the weighted least squares estimator of Einmahl et al. (2018), which is centred around the stable tail dependence function. The hypothesis testing framework described in that paper is valid for parameters in the interior of the parameter space, and hence it cannot be used to test whether one or more factor loadings are equal to zero. The hypothesis tests proposed in this paper can be applied to decide how many factors are necessary to model a certain dataset or to test if a more specific model (eg, Marshall–Olkin) is sufficient, bypassing the limitation that the number of factors has to be chosen upfront.

The paper is organized as follows. Section 2 presents background on multivariate extremes and introduces the weighted least squares estimator. Section 3 establishes asymptotic normality for the weighted least squares estimator when the true parameter vector lies on the boundary of the parameter space. Using results from Andrews (1999), Andrews (2001) and Andrews (2002), a deviance- and a Wald-type test statistic are proposed, whose asymptotic distributions are easily computable when the dimension of the parameter vector on the boundary is moderate. In Section 4.2, we perform simulations to evaluate the performance of the test statistics for the Brown–Resnick and the max-linear models. Finally, Section 5 illustrates our methods on two major stock market indices, the DAX and the CAC40. Technical definitions and proofs are deferred to the appendix.

## 2. Background

### 2.1. Multivariate extreme-value theory

Let  $\mathbf{X}_i = (X_{i1}, \dots, X_{id})$ ,  $i \in \{1, \dots, n\}$ , be random vectors in  $\mathbb{R}^d$  with cumulative distribution function  $F$  and marginal cumulative distribution functions  $F_1, \dots, F_d$ . Let  $\mathbf{M}_n := (M_{n1}, \dots, M_{nd})$  with  $M_{nj} := \max(X_{1j}, \dots, X_{nj})$  for  $j = 1, \dots, d$ . We say that  $F$  is in the *max-domain of attraction* of an *extreme-value distribution*  $G$  if there exist sequences of normalizing constants  $\mathbf{a}_n = (a_{n1}, \dots, a_{nd}) > 0$  and  $\mathbf{b}_n = (b_{n1}, \dots, b_{nd}) \in \mathbb{R}^d$  such that

$$\mathbb{P}\left[\frac{\mathbf{M}_n - \mathbf{b}_n}{\mathbf{a}_n} \leq \mathbf{x}\right] = F^n(\mathbf{a}_n \mathbf{x} + \mathbf{b}_n) \xrightarrow{d} G(\mathbf{x}), \quad \text{as } n \rightarrow \infty. \quad (1)$$

The margins,  $G_1, \dots, G_d$ , of  $G$  are *univariate extreme-value distributions*,

$$G_j(x_j) = \exp \left\{ - \left( 1 + \gamma_j \frac{x_j - \mu_j}{\sigma_j} \right)_+^{-1/\gamma_j} \right\}, \quad \sigma_j > 0, \gamma_j, \mu_j \in \mathbb{R},$$

where  $x_+ := \max(x, 0)$ . The function  $G$  is determined by

$$G(\mathbf{x}) = \exp \{ -\ell(-\log G_1(x_1), \dots, -\log G_d(x_d)) \},$$

where  $\ell: [0, \infty)^d \rightarrow [0, \infty)$  is called the *(stable) tail dependence function*,

$$\ell(\mathbf{x}) := \lim_{t \downarrow 0} t^{-1} \mathbb{P}[1 - F_1(X_{11}) \leq tx_1 \text{ or } \dots \text{ or } 1 - F_d(X_{1d}) \leq tx_d]. \quad (2)$$

The cumulative distribution function  $F$  is in the max-domain of attraction of a  $d$ -variate extreme value distribution  $G$  if and only if the limit in (2) exists and the marginal distributions in (1) converge to univariate extreme-value distributions. In what follows, we only assume existence of (2), which concerns the dependence structure of  $F$ , but not the marginal distributions  $F_1, \dots, F_d$ .

Because the class of stable tail dependence functions is infinite-dimensional, one usually considers parametric models for  $\ell$ . Henceforth we assume that  $\ell$  belongs to a parametric family  $\{\ell(\cdot; \boldsymbol{\theta}); \boldsymbol{\theta} \in \Theta\}$  with  $\Theta \subset \mathbb{R}^p$ . Some examples can be found below, see also de Haan and Ferreira (2006), Falk et al. (2010), Segers (2012) and references therein.

**Example 2.1.** The  $d$ -dimensional *logistic model* (Gumbel, 1960) has stable tail dependence function

$$\ell(\mathbf{x}; \boldsymbol{\theta}) = (x_1^{1/\theta} + \dots + x_d^{1/\theta})^\theta, \quad \boldsymbol{\theta} \in (0, 1].$$

If  $\theta = 1$ , the variables are (asymptotically) independent, while  $\theta \downarrow 0$  corresponds to the situation of complete dependence.

**Example 2.2.** The  $d$ -dimensional *Brown–Resnick model* defined on spatial locations  $\mathbf{s}_1, \dots, \mathbf{s}_d \in \mathbb{R}^2$  has stable tail dependence function

$$\ell(\mathbf{x}; \theta) = \sum_{j=1}^d x_j \Phi_{d-1}(\eta^{(j)}(1/\mathbf{x}); \Upsilon^{(j)}),$$

where  $\eta^{(j)}(\mathbf{x}) = (\eta_1^{(j)}(x_1, x_j), \dots, \eta_{j-1}^{(j)}(x_{j-1}, x_j), \eta_{j+1}^{(j)}(x_{j+1}, x_j), \dots, \eta_d^{(j)}(x_d, x_j)) \in \mathbb{R}^{d-1}$ ,

$$\eta_l^{(j)}(x_l, x_j) = \sqrt{\frac{\gamma(\mathbf{s}_j - \mathbf{s}_l)}{2}} + \frac{\log(x_l/x_j)}{\sqrt{2\gamma(\mathbf{s}_j - \mathbf{s}_l)}} \in \mathbb{R},$$

$\gamma(\mathbf{s}) = (||\mathbf{s}||/\rho)^\alpha$  and  $\Upsilon^{(l)} \in \mathbb{R}^{(d-1) \times (d-1)}$  is the correlation matrix with entries

$$\Upsilon_{lk}^{(j)} = \frac{\gamma(\mathbf{s}_j - \mathbf{s}_l) + \gamma(\mathbf{s}_j - \mathbf{s}_k) - \gamma(\mathbf{s}_l - \mathbf{s}_k)}{2\sqrt{\gamma(\mathbf{s}_j - \mathbf{s}_l)\gamma(\mathbf{s}_j - \mathbf{s}_k)}}, \quad l, k = 1, \dots, d; l, k \neq j,$$

(Kabluchko et al., 2009; Huser and Davison, 2013). The parameter vector is  $\theta = (\rho, \alpha) \in (0, \infty) \times (0, 2]$ . The *Smith model* (Smith, 1989) is obtained when  $\alpha = 2$ .

**Example 2.3.** The *max-linear model* with  $r$  factors has stable tail dependence function

$$\ell(\mathbf{x}; \theta) = \sum_{t=1}^r \max_{j=1, \dots, d} b_{jt} x_j, \quad \mathbf{x} \in [0, \infty)^d, \quad (3)$$

where the factor loadings  $b_{jt}$  are non-negative constants such that  $\sum_{t=1}^r b_{jt} = 1$  for every  $j \in \{1, \dots, d\}$  and all column sums of the  $d \times r$  matrix  $B := (b_{jt})_{j,t}$  are positive (Einmahl et al., 2012). Since the rows of  $B$  sum up to one, the parameter matrix has only  $d \times (r-1)$  free elements. Rearranging the columns of  $B$  will not change the value of the stable tail dependence function. For identification purposes, we define the parameter vector  $\theta$  by stacking the columns of  $B$  in decreasing order of their sums, leaving out the column with the lowest sum. An example of a random vector  $\mathbf{Z} = (Z_1, \dots, Z_d)$  that has stable tail dependence function (3) is

$$Z_j = \max_{t=1, \dots, r} b_{jt} S_t, \quad \text{for } j \in \{1, \dots, d\},$$

where  $S_1, \dots, S_r$  are independent unit Fréchet variables,  $\mathbb{P}[S_j \leq x] = \exp(-1/x)$  for  $x > 0$  and  $j \in \{1, \dots, d\}$ .

**Example 2.4.** Let  $I_1, \dots, I_d$  denote random (possibly dependent) Bernoulli random variables with  $p_j = \mathbb{P}[I_j = 1] \in (0, 1]$  for  $j \in \{1, \dots, d\}$ . Let, for  $J \subset \{1, \dots, d\}$ ,  $p(J) = \mathbb{P}[\{j = 1, \dots, d : I_j = 1\} = J]$ , so that  $(p(J))_{J \subset \{1, \dots, d\}}$  is a probability distribution. The *multivariate Marshall–Olkin model* (Ebrechts et al., 2003; Segers, 2012) has stable tail dependence function

$$\ell(\mathbf{x}; \theta) = \sum_{\emptyset \neq J \subset \{1, \dots, d\}} p(J) \max_{j \in J} \left( \frac{x_j}{p_j} \right).$$

In this model, any subset of components of the vector is assigned a shock that influences all components of that subset. An interpretation in terms of credit risk modelling can be found in Ebrechts et al. (2003).

The Marshall–Olkin model is obtained as a special case of the max-linear model (3) by setting  $b_{jt} = \{p(J_t)/p_j\} \mathbb{1}(j \in J_t)$ , where  $J_t \in \mathcal{P}(\{1, \dots, d\})$ ,  $J_t \neq \emptyset$ , and  $\mathcal{P}(A)$  denotes the power set of  $A$ . The Marshall–Olkin model is well-known in  $d = 2$ , where

$$B = \begin{pmatrix} b_{11} & 0 & 1 - b_{11} \\ 0 & b_{22} & 1 - b_{22} \end{pmatrix} = \begin{pmatrix} \frac{P[I_1 = 1, I_2 = 0]}{P[I_1 = 1]} & 0 & \frac{P[I_1 = 1, I_2 = 1]}{P[I_1 = 1]} \\ 0 & \frac{P[I_1 = 0, I_2 = 1]}{P[I_2 = 1]} & \frac{P[I_1 = 1, I_2 = 1]}{P[I_2 = 1]} \end{pmatrix}.$$

## 2.2. Estimation of the stable tail dependence function

### 2.2.1. Nonparametric estimation of the stable tail dependence function

Let  $k = k_n \in (0, n]$  be such that  $k \rightarrow \infty$  and  $k/n \rightarrow 0$  as  $n \rightarrow \infty$ . A straightforward nonparametric estimator of  $\ell$  is obtained by replacing  $\mathbb{P}$  and  $F_1, \dots, F_d$  in (2) by the (modified) empirical distribution functions and replacing  $t$  by  $k/n$ , yielding

$$\tilde{\ell}_{n,k}(x) := \frac{1}{k} \sum_{i=1}^n \mathbb{1}\{R_{i1} > n + 1/2 - kx_1 \text{ or } \dots \text{ or } R_{id} > n + 1/2 - kx_d\}.$$

Here,  $R_{ij} = \sum_{t=1}^n \mathbb{1}\{X_{tj} \leq X_{ij}\}$  denotes the rank of  $X_{ij}$  among  $X_{1j}, \dots, X_{nj}$ . This estimator, the *empirical tail dependence function*, was introduced in slightly different form for  $d = 2$  in Huang (1992) and studied further in Drees and Huang (1998). Using

$n + 1/2$  rather than  $n$  allows for better finite-sample properties, that is, for a lower mean squared error (Einmahl et al., 2012).

Another nonparametric estimator is the *beta tail dependence function*, which was very recently proposed in Kiriliouk et al. (2018b). Contrary to the empirical tail dependence function, the beta tail dependence function is a smooth estimator which leads to some improvement in its finite-sample behavior. Its name stems from the fact that it is based on the empirical beta copula (Segers et al., 2017). We define

$$\tilde{\ell}_{n,k}^{\beta}(\mathbf{x}) = \frac{n}{k} \left\{ 1 - \mathbb{C}_n^{\beta} \left( 1 - \frac{k\mathbf{x}}{n} \right) \right\}, \quad \text{where} \quad \mathbb{C}_n^{\beta}(\mathbf{u}) = \frac{1}{n} \sum_{i=1}^n \prod_{j=1}^d F_{n,R_{ij}}(u_j),$$

and for  $r \in \{1, \dots, n\}$ ,  $F_{n,r}(u) = \sum_{s=1}^r \binom{n}{s} u^s (1-u)^{n-s}$  is the cumulative distribution function of a Beta( $r, n+1-r$ ) random variable.

A drawback of  $\tilde{\ell}_{n,k}$  or  $\tilde{\ell}_{n,k}^{\beta}$  might be their possibly growing bias as  $k$  increases: Fougères et al. (2015) show that, under suitable conditions, the estimator  $\tilde{\ell}_{n,k}$  satisfies the asymptotic expansion

$$\tilde{\ell}_{n,k}(\mathbf{x}) - \ell(\mathbf{x}) \approx k^{-1/2} Z_{\ell}(\mathbf{x}) + \alpha(n/k)M(\mathbf{x}),$$

where  $Z_{\ell}$  is a continuous centered Gaussian process,  $\alpha$  is a function such that  $\lim_{x \rightarrow \infty} \alpha(x) = 0$  and  $M$  is a continuous function. When  $\sqrt{k}\alpha(n/k)$  tends to a non-zero constant, an asymptotic bias appears. In Fougères et al. (2015) and Beirlant et al. (2016), this bias is estimated and subtracted from the estimator  $\tilde{\ell}_{n,k}(\mathbf{x})$  in order to propose new bias-corrected estimators.

Finally, many other nonparametric estimators exist of the Pickands dependence function, which is the restriction of  $\ell$  to the unit simplex. See, for instance, Capérea et al. (1997), Zhang et al. (2008), Gudendorf and Segers (2012), Berghaus et al. (2013), Vettori et al. (2016) and Marcon et al. (2017). While these can be transformed into estimators of the stable tail dependence function, they rely on stronger assumptions, i.e., they are based on data from an extreme-value distribution, and we will not consider them here.

### 2.2.2. Semi-parametric estimation of the stable tail dependence function

Nonparametric estimation forms a stepping stone for semi-parametric estimation (Einmahl et al., 2012; 2016; 2018). Here, we focus on the method proposed in Einmahl et al. (2018). Let  $\hat{\ell}_{n,k}$  denote any initial estimator of  $\ell$  based on  $\mathbf{X}_1, \dots, \mathbf{X}_n$ . Let  $\mathbf{c}_1, \dots, \mathbf{c}_q \in [0, \infty)^d$ , with  $\mathbf{c}_m = (c_{m1}, \dots, c_{md})$  for  $m = 1, \dots, q$ , be  $q$  points in which we will evaluate  $\ell$  and  $\hat{\ell}_{n,k}$ . Consider for  $\boldsymbol{\theta} \in \Theta$  the  $q \times 1$  column vectors

$$\hat{\mathbf{L}}_{n,k} := (\hat{\ell}_{n,k}(\mathbf{c}_m))_{m=1}^q, \quad L(\boldsymbol{\theta}) := (\ell(\mathbf{c}_m; \boldsymbol{\theta}))_{m=1}^q, \quad D_{n,k}(\boldsymbol{\theta}) := \hat{\mathbf{L}}_{n,k} - L(\boldsymbol{\theta}). \quad (4)$$

The points  $\mathbf{c}_1, \dots, \mathbf{c}_q$  need to be chosen in such a way that the map  $L: \Theta \rightarrow \mathbb{R}^q$  is one-to-one, i.e.,  $\boldsymbol{\theta}$  is identifiable from  $\ell(\mathbf{c}_1; \boldsymbol{\theta}), \dots, \ell(\mathbf{c}_q; \boldsymbol{\theta})$ . In particular, we will need to assume that  $q \geq p$ .

For  $\boldsymbol{\theta} \in \Theta$ , let  $\Omega(\boldsymbol{\theta})$  be a symmetric, positive definite  $q \times q$  matrix and define

$$f_{n,k}(\boldsymbol{\theta}) := D_{n,k}^T(\boldsymbol{\theta}) \Omega(\boldsymbol{\theta}) D_{n,k}(\boldsymbol{\theta}). \quad (5)$$

The continuous updating weighted least squares estimator for  $\boldsymbol{\theta}_0$  is defined as

$$\hat{\boldsymbol{\theta}}_{n,k} := \arg \min_{\boldsymbol{\theta} \in \Theta} f_{n,k}(\boldsymbol{\theta}) = \arg \min_{\boldsymbol{\theta} \in \Theta} \{ D_{n,k}(\boldsymbol{\theta})^T \Omega(\boldsymbol{\theta}) D_{n,k}(\boldsymbol{\theta}) \}. \quad (6)$$

In Section 4, we will study the performance of this estimator for  $\Omega(\boldsymbol{\theta}) = I_q$ , the  $q \times q$  identity matrix. Expression (6) then simplifies to

$$\hat{\boldsymbol{\theta}}_{n,k} = \arg \min_{\boldsymbol{\theta} \in \Theta} \sum_{m=1}^q (\hat{\ell}_{n,k}(\mathbf{c}_m) - \ell(\mathbf{c}_m; \boldsymbol{\theta}))^2.$$

Einmahl et al. (2018) show the consistency and asymptotic normality of  $\hat{\boldsymbol{\theta}}_{n,k}$  under the assumption that the true parameter vector  $\boldsymbol{\theta}_0$  is in the interior of  $\Theta$ . In the next section we give the asymptotic distribution of  $\hat{\boldsymbol{\theta}}_{n,k}$  without this restriction.

## 3. Inference on tail dependence parameters on the boundary of the parameter space

The asymptotic results presented in Section 3.1 build on Andrews (1999), who established the asymptotic distribution of a general extremum estimator when one or more parameters lie on the boundary of the parameter space. The methodology to obtain the asymptotic distribution of  $\sqrt{k}(\hat{\boldsymbol{\theta}}_{n,k} - \boldsymbol{\theta}_0)$  can be summarized as follows: first, one shows that as  $n \rightarrow \infty$ , the criterion function  $f_{n,k}(\boldsymbol{\theta})$  in (5) is equal to a quadratic function  $q_{n,k}(\sqrt{k}(\boldsymbol{\theta} - \boldsymbol{\theta}_0))$  plus a term that does not depend on  $\boldsymbol{\theta}$ . If  $\hat{\boldsymbol{\theta}}_{n,k}$  is consistent, then its asymptotic distribution is only affected by the part of  $\Theta$  close to  $\boldsymbol{\theta}_0$ ; equivalently, we are only concerned with the shifted parameter space  $\Theta - \boldsymbol{\theta}_0$  near the origin. If  $\Theta - \boldsymbol{\theta}_0$  can be approximated near the origin by a convex cone  $\Lambda$ , one can show that minimizing  $f_{n,k}(\boldsymbol{\theta})$  over  $\boldsymbol{\theta} \in \Theta$  is asymptotically equivalent to minimizing  $q_{n,k}(\boldsymbol{\lambda})$  over  $\boldsymbol{\lambda} \in \Lambda$ . Finally,  $\sqrt{k}(\hat{\boldsymbol{\theta}}_{n,k} - \boldsymbol{\theta}_0)$  converges in distribution to the argument minimizing the limit of  $q_{n,k}(\boldsymbol{\lambda})$  as  $n \rightarrow \infty$ .

In [Section 3.2](#), we show how a closed-form expression can be obtained for the asymptotic distribution of  $\sqrt{k}(\hat{\theta}_{n,k} - \theta_0)$ . For  $\beta \in \Theta$ , we show in [Section 3.3](#) how to test  $H_0: \beta = \beta_*$  against  $H_1: \beta \neq \beta_*$  when, under the null hypothesis,  $\beta_*$  is on the boundary of the alternative hypothesis.

We set up some notation first. Let  $B_\varepsilon(\theta)$  denote an open ball centered at  $\theta$  with radius  $\varepsilon$  and let  $C_\varepsilon(\theta)$  denote an open cube centered at  $\theta$  with sides of length  $2\varepsilon$ . Let  $\text{cl}(\Theta)$  denote the closure of  $\Theta$ . A set  $\Gamma \subset \mathbb{R}^p$  is said to be *locally equal* to a set  $\Lambda \subset \mathbb{R}^p$  if  $\Gamma \cap B_\varepsilon(\theta) = \Lambda \cap B_\varepsilon(\theta)$  for some  $\varepsilon > 0$ . Finally, a set  $\Lambda \subset \mathbb{R}^p$  is a *cone* if  $\lambda \in \Lambda$  implies  $a\lambda \in \Lambda$  for all  $a > 0$ .

### 3.1. Estimation, consistency and asymptotic normality

When  $\theta_0$  is on the boundary of  $\Theta$ , the map  $L$  in (4) is not defined and thus not differentiable on a neighbourhood of  $\theta_0$ . We will need the following assumption.

(A1)  $\Theta$  includes a set  $\Theta^+$  such that  $\Theta^+ - \theta_0$  equals the intersection of a union of orthants and an open cube  $C_\varepsilon(\theta_0)$  for some  $\varepsilon > 0$ . Moreover,  $\Theta \cap B_{\varepsilon_1}(\theta_0) \subset \Theta^+$  for some  $\varepsilon_1 > 0$ .

If  $\Theta - \theta_0$  happens to be locally equal to a union of orthants, we can simply set  $\Theta^+ = \Theta \cap C_\varepsilon(\theta_0)$ . This is the case for the models considered in [Section 4](#). We will assume existence of the so-called *left/right (l/r) partial derivatives* on  $\Theta^+$ ; a formal definition is given in the appendix. The shape of  $\Theta^+$  is such that these can always be defined.

Write  $\dot{L} := (\partial/\partial\theta)L(\theta) \in \mathbb{R}^{q \times p}$  for  $\theta \in \Theta^+$ , where  $\dot{L}$  denotes the matrix of l/r partial derivatives. Let  $\lambda_1(\theta) > 0$  denote the smallest eigenvalue of  $\Omega(\theta)$ .

**Theorem 3.1** (Existence, uniqueness and consistency). *Let  $\mathbf{c}_1, \dots, \mathbf{c}_q \in [0, \infty)^d$  be  $q \geq p$  points such that the map  $L: \theta \mapsto (\ell(\mathbf{c}_m; \theta))_{m=1}^q$  is a homeomorphism. Let  $\theta_0 \in \text{cl}(\Theta)$  and assume that (A1) holds, that each element of  $L(\theta)$  has continuous l/r partial derivatives of order two on  $\Theta^+$ , that  $\dot{L}(\theta_0)$  is of full rank, that  $\Omega: \Theta \rightarrow \mathbb{R}^{q \times q}$  has continuous l/r partial derivatives on  $\Theta^+$  and that  $\inf_{\theta \in \Theta} \lambda_1(\theta) > 0$ . Finally assume, for  $m = 1, \dots, q$ ,*

$$\hat{\ell}_{n,k}(\mathbf{c}_m) \xrightarrow{p} \ell(\mathbf{c}_m; \theta_0), \quad \text{as } n \rightarrow \infty. \quad (7)$$

Then with probability tending to one, the minimizer  $\hat{\theta}_{n,k}$  in (6) exists and is unique. Moreover,

$$\hat{\theta}_{n,k} \xrightarrow{p} \theta_0, \quad \text{as } n \rightarrow \infty.$$

We omit the proof of this theorem since it is directly obtained by replacing  $B_\varepsilon(\theta_0)$  by  $\Theta^+$  in the proof of [Einmahl et al. \(2018, Theorem 1\)](#).

When  $f_{n,k}$  is not defined on a neighbourhood of  $\theta_0$ , but there exists a set  $\Theta^+$  which satisfies (A1), a Taylor expansion of  $f_{n,k}(\theta)$  around  $f_{n,k}(\theta_0)$  holds ([Andrews, 1999, Theorem 6](#)). For each  $\theta \in \Theta^+$ , we have

$$f_{n,k}(\theta) = f_{n,k}(\theta_0) + Df_{n,k}(\theta_0)^T(\theta - \theta_0) + (\theta - \theta_0)^T D^2 f_{n,k}(\theta_0)(\theta - \theta_0)/2 + R_{n,k}(\theta),$$

where  $Df_{n,k}$  and  $D^2 f_{n,k}$  are based on l/r partial derivatives and  $R_{n,k}(\theta)$  is the remainder term. We suppress dependence of  $\Omega$ ,  $\dot{L}$  and  $D_{n,k}$  on  $\theta_0$  for ease of notation. From [Einmahl et al. \(2018, Proof of Theorems 3.2.1 and 3.2.2\)](#) we know that

$$Df_{n,k}(\theta_0) = -2D_{n,k}^T \Omega \dot{L} + o_p(1), \quad D^2 f_{n,k}(\theta_0) = 2\dot{L}^T \Omega \dot{L} + o_p(1),$$

so the quadratic expansion above is equal to

$$f_{n,k}(\theta) = f_{n,k}(\theta_0) - 2D_{n,k}^T \Omega \dot{L}(\theta - \theta_0) + (\theta - \theta_0)^T \dot{L}^T \Omega \dot{L}(\theta - \theta_0) + R_{n,k}(\theta).$$

Define

$$J := \dot{L}^T \Omega \dot{L} \in \mathbb{R}^{p \times p}, \quad \mathbf{Y}_{n,k} := \sqrt{k} J^{-1} \dot{L}^T \Omega D_{n,k} \in \mathbb{R}^p,$$

and

$$q_{n,k}(\lambda) = (\lambda - \mathbf{Y}_{n,k})^T J (\lambda - \mathbf{Y}_{n,k}).$$

Then  $f_{n,k}(\theta)$  can be written as

$$f_{n,k}(\theta) = f_{n,k}(\theta_0) - k^{-1} \mathbf{Y}_{n,k}^T J \mathbf{Y}_{n,k} + k^{-1} q_{n,k}(\sqrt{k}(\theta - \theta_0)) + R_{n,k}(\theta).$$

We see that  $f_{n,k}(\theta)$  is equal to a quadratic function plus a term that does not depend on  $\theta$  plus  $R_{n,k}(\theta)$ . In [Andrews \(2002, Lemma 3\)](#), it is shown that under the assumptions of [Theorem 3.1](#) and (A3) below, the remainder term  $R_{n,k}(\theta)$  is sufficiently small so that minimizing the quadratic approximation to  $f_{n,k}(\theta)$  is equivalent to minimizing  $q_{n,k}(\sqrt{k}(\theta - \theta_0))$ . Now, assume that

(A2)  $\Theta - \theta_0$  is locally equal to a convex cone  $\Lambda \subset \mathbb{R}^p$ .

Then

$$\inf_{\theta \in \Theta} q_{n,k}(\sqrt{k}(\theta - \theta_0)) = \inf_{\lambda \in \Lambda} q_{n,k}(\lambda) + o_p(1),$$

for some cone  $\Lambda$ . Finally, assume that

$$(A3) \quad \sqrt{k}D_{n,k}(\theta_0) \xrightarrow{d} \mathbf{D} \sim \mathcal{N}_q(0, \Sigma(\theta_0)) \text{ as } n \rightarrow \infty, \text{ for some covariance matrix } \Sigma(\theta_0).$$

Assumptions (A3) and (7) hold for the nonparametric estimators presented in Section 2.2.1 under some general regularity conditions. The matrix  $\Sigma$  can be expressed in terms of  $\ell$  and its first-order partial derivatives. We have

$$\mathbf{Y}_{n,k} \xrightarrow{d} \mathbf{Y} := J^{-1} \dot{L}^T \Omega \mathbf{D}, \quad q_{n,k}(\lambda) \xrightarrow{d} q(\lambda) := (\lambda - \mathbf{Y})^T J (\lambda - \mathbf{Y}),$$

for all  $\lambda \in \mathbb{R}^p$ .

**Theorem 3.2** (Asymptotic normality). *If all of the above assumptions hold,*

$$\sqrt{k}(\hat{\theta}_{n,k} - \theta_0) \xrightarrow{d} \hat{\lambda} = \arg \min_{\lambda \in \Lambda} q(\lambda), \quad \text{as } n \rightarrow \infty.$$

Here  $\hat{\lambda}$  can be interpreted as the projection of  $\mathbf{Y}$  on  $\Lambda$  with respect to the norm  $\|y\|_J := y^T J y$ . This theorem is a special case of Andrews (1999, Theorem 3). In the appendix, we verify that our assumptions are sufficient. Note that if  $\Theta$  includes a neighbourhood of  $\theta_0$ , then  $\Lambda = \mathbb{R}^p$  and we find the same distribution as in Einmahl et al. (2018, Theorem 2) since  $\hat{\lambda} = \mathbf{Y}$ .

### 3.2. Simplifying the asymptotic distribution

The goal of this section is to simplify the asymptotic distribution of  $\sqrt{k}(\hat{\theta}_{n,k} - \theta_0)$  and to give conditions under which (part of)  $\hat{\lambda}$  has a closed-form expression. Let  $\mathbf{G} := \dot{L}^T \Omega \mathbf{D}$ , i.e.,  $\mathbf{Y} = J^{-1} \mathbf{G}$ . We start by partitioning the vector  $\theta_0 \in \mathbb{R}^p$  into two subvectors,  $\beta_0 \in \mathbb{R}^c$  and  $\delta_0 \in \mathbb{R}^{p-c}$  for  $c \in \{1, \dots, p\}$ , where  $\delta_0$  consists of all parameters that are in the interior of the parameter space. We partition  $\theta_{n,k}$ ,  $\mathbf{Y}$ ,  $J$ ,  $\mathbf{G}$  and  $\lambda$  accordingly:

$$\theta_{n,k} = \begin{pmatrix} \beta_{n,k} \\ \delta_{n,k} \end{pmatrix}, \quad \mathbf{Y} = \begin{pmatrix} \mathbf{Y}_\beta \\ \mathbf{Y}_\delta \end{pmatrix}, \quad J = \begin{pmatrix} J_\beta & J_{\beta\delta} \\ J_{\delta\beta} & J_\delta \end{pmatrix}, \quad \mathbf{G} = \begin{pmatrix} \mathbf{G}_\beta \\ \mathbf{G}_\delta \end{pmatrix}, \quad \lambda = \begin{pmatrix} \lambda_\beta \\ \lambda_\delta \end{pmatrix}.$$

Let  $I_c$  denote the  $c \times c$  identity matrix. For  $H := (I_c : \mathbf{0}) \in \mathbb{R}^{c \times p}$ , define

$$q_\beta(\lambda_\beta) := (\lambda_\beta - \mathbf{Y}_\beta)^T (HJ^{-1}H^T)^{-1} (\lambda_\beta - \mathbf{Y}_\beta).$$

Suppose that

$$(A4) \quad \text{The cone } \Lambda \text{ of assumption (A2) is equal to the product set } \Lambda_\beta \times \mathbb{R}^{p-c}, \text{ where } \Lambda_\beta \subset \mathbb{R}^c \text{ is a cone.}$$

**Corollary 3.3.** *If all of the above assumptions hold, then*

$$\begin{aligned} \sqrt{k}(\hat{\beta}_{n,k} - \beta_0) &\xrightarrow{d} \hat{\lambda}_\beta := \arg \min_{\lambda_\beta \in \Lambda_\beta} q_\beta(\lambda_\beta), \\ \sqrt{k}(\hat{\delta}_{n,k} - \delta_0) &\xrightarrow{d} J_\delta^{-1} \mathbf{G}_\delta - J_\delta^{-1} J_{\delta\beta} \hat{\lambda}_\beta. \end{aligned}$$

When  $\Lambda_\beta$  is defined by equality and/or inequality constraints, a closed-form expression for  $\hat{\lambda}_\beta$  can be computed; we give some examples that are relevant for Section 4. For a more formal solution, see Andrews (1999, Theorem 5).

**Example 3.1** ( $c = 1$ ). Suppose that  $\Theta = [0, 1]^p$ . If  $\beta_0 = 0$ , then  $\Lambda_\beta = [0, \infty)$  and  $\hat{\lambda}_\beta = \max(Y_\beta, 0)$ . If  $\beta_0 = 1$ , then  $\Lambda_\beta = (-\infty, 0]$  and  $\hat{\lambda}_\beta = \min(Y_\beta, 0)$ .

**Example 3.2** ( $c = 2$ ). Suppose that  $\Theta = [0, 1]^p$ . If  $\beta_0 = (1, 1)$ , then  $\Lambda_\beta = (-\infty, 0]^2$ . Let  $\rho := HJ^{-1}H^T$ . Then

$$\begin{aligned} \hat{\lambda}_\beta &:= \mathbb{1}\{Y_{\beta 1} < 0, Y_{\beta 2} < 0\} \mathbf{Y}_\beta + \mathbb{1}\{Y_{\beta 1} - \rho_{21}Y_{\beta 2} < 0, Y_{\beta 2} \geq 0\} (Y_{\beta 1} - \rho_{21}Y_{\beta 2}, 0)^T \\ &\quad + \mathbb{1}\{Y_{\beta 1} \geq 0, Y_{\beta 2} - \rho_{12}Y_{\beta 1} < 0\} (0, Y_{\beta 2} - \rho_{12}Y_{\beta 1})^T. \end{aligned}$$

If  $\beta_0 = (0, 0)$ ,  $\beta_0 = (0, 1)$  or  $\beta_0 = (1, 0)$ , similar expressions can be obtained by reversing the first and/or the second inequalities in each of the indicator functions above.



### 3.3. Hypothesis testing for tail dependence parameters on the boundary of the parameter space

We are interested in constructing hypothesis tests for parameter values that, under the null hypothesis, are on the boundary of the alternative hypothesis. We propose two test statistics, whose asymptotic distribution follows from the results in Andrews (2001).

Assume that there are no nuisance parameters on the boundary, i.e., all components of  $\theta_0$  that lie on the boundary are part of the null hypothesis. We are interested in testing

$$H_0 : \beta = \beta_* \quad \text{vs} \quad H_1 : \beta \neq \beta_*, \quad \beta_* \in \mathbb{R}^c.$$

Let  $\Theta_0 := \{\theta \in \Theta : \theta = (\beta_*, \delta) \text{ for some } \delta \in \mathbb{R}^{p-c}\}$  denote the restricted parameter space. Assume that

(A5) For all  $\theta \in \Theta_0$ ,  $\Theta$  is a product set with respect to  $(\beta, \delta)$  local to  $\theta$ . That is, for all  $\theta \in \Theta_0$ ,

$$\Theta \cap B_\varepsilon(\theta) = (B \times \Delta) \cap B_\varepsilon(\theta) \quad \text{for some } B \subset \mathbb{R}^c, \Delta \subset \mathbb{R}^{p-c} \text{ and } \varepsilon > 0.$$

Define  $\hat{\theta}_{n,k}^{(0)} := \arg \min_{\theta \in \Theta_0} f_{n,k}(\theta)$ . A deviance test statistic can be defined as

$$T_{n,k}^{(1)} := k \left( f_{n,k}(\hat{\theta}_{n,k}^{(0)}) - f_{n,k}(\hat{\theta}_{n,k}) \right).$$

**Corollary 3.4.** Suppose  $\theta_0 \in \Theta_0$  and all previous assumptions hold. Then

$$T_{n,k}^{(1)} \xrightarrow{d} \hat{\lambda}_\beta^T (HJ^{-1}H^T)^{-1} \hat{\lambda}_\beta.$$

Recall that  $G = \dot{L}^T \Omega \dot{D}$  and let  $\mathcal{J}$  denote its covariance matrix, i.e.,  $G \sim \mathcal{N}_p(\mathbf{0}, \mathcal{J})$  with  $\mathcal{J} = \dot{L}^T \Omega \Sigma \Omega \dot{L} \in \mathbb{R}^{p \times p}$ . Note that  $\mathcal{J} = J$  if and only if  $\Omega = \Sigma^{-1}$ . If  $\Lambda_\beta = \mathbb{R}^c$  (no parameters on the boundary) and  $\mathcal{J} = J$ , then

$$T_{n,k}^{(1)} \xrightarrow{d} Y_\beta^T (HJ^{-1}H^T)^{-1} Y_\beta \sim \chi_c^2,$$

where  $\chi_c^2$  denotes a chi-squared random variable with  $c$  degrees of freedom.

A Wald-type test statistic can be based on the quadratic form in  $\hat{\beta}_{n,k} - \beta_*$ . Let  $\hat{V}_n^{-1} := (HJ_n^{-1}H^T)^{-1} \in \mathbb{R}^{c \times c}$  denote a weight matrix where  $J_n := J(\hat{\theta}_{n,k})$ . Define

$$T_{n,k}^{(2)} := k \left( \hat{\beta}_{n,k} - \beta_* \right)^T \hat{V}_n^{-1} \left( \hat{\beta}_{n,k} - \beta_* \right).$$

**Corollary 3.5.** Suppose  $\theta_0 \in \Theta_0$  and all of the above assumptions hold. Then

$$T_{n,k}^{(2)} \xrightarrow{d} \hat{\lambda}_\beta^T (HJ^{-1}H^T)^{-1} \hat{\lambda}_\beta.$$

Corollary 3.4 is a special case of Andrews (2001, Theorem 4c) and Corollary 3.5 is a special case of Andrews (2001, Theorem 6d); their proofs are direct and thus omitted.

## 4. Simulation studies

We conduct simulation experiments where we simulate 1000 samples of size  $n = 5000$  from a parametric model and we assess the quality of the weighted least squares estimator in terms of its root mean squared error (RMSE) for settings where one or more of the parameters are on the boundary of the parameter space. We take  $k \in \{25, 50, \dots, 300\}$  and we use the empirical and the beta tail dependence function as initial estimators. Next, we study the performance of the two test statistics introduced in Section 3.3 in terms of empirical level and power for  $k \in \{25, 50, 75, 100\}$ , as we observe that higher  $k$  leads to a steadily growing bias that quickly deteriorates the performance of the hypothesis test. In all experiments, we take  $\Omega = I_q$ , the  $q \times q$  identity matrix, because using an optimal weight matrix has a very minor effect on the quality of estimation while severely slowing down the estimation procedure. Moreover, inverting  $\Sigma$  may be hindered by numerical problems in the case of a max-linear model (Einmahl et al., 2018).

### 4.1. Brown–Resnick model

We simulate data from a Brown–Resnick model with  $\alpha = 2$  and  $\rho = 1$  on a regular  $3 \times 2$  grid ( $d = 6$ ) and on a regular  $4 \times 4$  grid ( $d = 16$ ). As in Einmahl et al. (2018), we let  $\mathbf{c}_m \in \{0, 1\}^d$  such that exactly two components of  $\mathbf{c}_m$  are equal to one (meaning that we consider a pairwise estimator) and we focus on pairs of neighbouring locations only, i.e., locations that are at most a distance of  $\sqrt{2}$  apart. This leads to  $q = 11$  and  $q = 42$  respectively.


Fig. 1 shows the RMSE of the parameter estimates based on the empirical tail dependence function (solid lines) and the beta tail dependence function (dashed lines). We see that the empirical tail dependence function outperforms the beta tail dependence function for  $\alpha$ , while the opposite is true for  $\rho$ , although differences are very minor. For small values of  $k$ , the RMSE is lower for  $d = 16$  than for  $d = 6$ .




**Table 1**

Empirical level of  $T_{n,k}^{(2)}$  based on the empirical and the beta tail dependence function for a significance level of 0.05.

	$k = 25$		$k = 50$		$k = 75$		$k = 100$	
	emp	beta	emp	beta	emp	beta	emp	beta
$d = 6$	6.0	4.3	5.1	5.0	7.0	5.5	8.8	8.0
$d = 16$	6.7	10.3	8.3	10.2	9.8	12.4	13.2	16.0



**Fig. 1.** RMSE for parameter estimators of the Brown-Resnick model based on the empirical tail dependence function (solid lines) and the beta tail dependence function (dashed lines) for  $(\alpha, \rho) = (2, 1)$ ;  $d = 6$  (left) and  $d = 16$  (right).



**Fig. 2.** Empirical power of  $T_{n,k}^{(2)}$  based on the empirical tail dependence function (solid lines) and the beta tail dependence function (dashed lines) for  $k \in \{25, 50, 75, 100\}$ ;  $d = 6$  (top) and  $d = 16$  (bottom).

Next, we study the empirical level and power of the test  $H_0: \alpha = 2$ . Assumption (A4) holds with  $\Lambda = (-\infty, 0] \times \mathbb{R}$ . Table 1 shows the empirical level of the test statistic  $T_{n,k}^{(2)}$  based on a significance level of  $\alpha = 0.05$ . We do not display  $T_{n,k}^{(1)}$  as its behaviour is identical to that of  $T_{n,k}^{(2)}$ . Surprisingly, the empirical tail dependence function is preferred for  $d = 16$  while the beta tail dependence function performs best for  $d = 6$ . In general, we can conclude that low values of  $k$  are to be preferred and that the test performs better for lower dimension. Fig. 2 shows the empirical power of  $T_{n,k}^{(2)}$  as a function of  $\alpha \in [1.5, 2)$ . The power increases both with the dimension and with  $k$ .

**Table 2**

Empirical level of  $T_{n,k}^{(1)}$  and  $T_{n,k}^{(2)}$  for models 1–3 (M1–M3) based on the empirical and the beta tail dependence function for a significance level of 0.05.

$T_{n,k}^{(1)}$	$k = 25$		$k = 50$		$k = 75$		$k = 100$	
	emp	beta	emp	beta	emp	beta	emp	beta
M1	2.9	2.1	6.1	4.2	5.9	5.3	8.0	6.4
M2	0.3	2.3	8.7	5.0	12.4	12.6	40.3	27.7
M3	3.3	7.9	7.9	9.7	6.5	9.8	9.8	10.0
$T_{n,k}^{(2)}$	$k = 25$		$k = 50$		$k = 75$		$k = 100$	
	emp	beta	emp	beta	emp	beta	emp	beta
M1	3.0	2.9	6.2	5.0	6.1	6.1	8.5	7.1
M2	0.3	2.4	9.2	5.3	12.5	13.0	41.3	28.3
M3	5.4	9.9	10.3	10.2	8.4	10.7	10.7	11.1

#### 4.2. Max-linear model

The constants  $\mathbf{c}_1, \dots, \mathbf{c}_q$  need to be chosen such that the max-linear model is identifiable. In Einmahl et al. (2018) it was observed that taking extremal coefficients for  $\mathbf{c}_m$ , i.e.,  $\mathbf{c}_m \in \{0, 1\}^d$  which at least two non-zero elements, is not enough for identifiability. Instead, one needs to choose  $\mathbf{c}_m$  such that its non-zero elements are unequal. For some theoretical considerations, see also Einmahl et al. (2018, Appendix B). Simulation experiments showed that in practice, taking a large grid of values (meaning that  $q \gg p$ ) on  $[0, 1]^d$  will lead to good estimators in terms of RMSE. Hence, in all following simulation experiments, we choose  $\mathbf{c}_1, \dots, \mathbf{c}_q$  such that

$$\mathbf{c}_m \in \{0, 0.01, 0.1, 0.2, \dots, 0.8, 0.9, 0.99, 1\}^2, \quad m \in \{1, \dots, q\}.$$

##### 4.2.1. Testing the structure of a max-linear model

We consider three different scenarios:

**Model 1** Let  $r = 2$  and  $d = 3$ , so that  $\Theta = [0, 1]^3$ . Let  $\theta_0 = (b_{11}, b_{21}, b_{31}) = (1, 0.7, 0.2)$ . We test  $H_0 : b_{11} = 1$ . Assumption (A4) holds with  $\Lambda = (-\infty, 0] \times \mathbb{R}^2$ .

**Model 2** Let  $r = 2$  and  $d = 3$ , so that  $\Theta = [0, 1]^3$ . Let  $\theta_0 = (b_{11}, b_{21}, b_{31}) = (1, 0.7, 0)$ . We test  $H_0 : (b_{11}, b_{31}) = (1, 0)$ . Assumption (A4) holds with  $\Lambda = (-\infty, 0] \times [0, \infty) \times \mathbb{R}$ .

**Model 3** Let  $r = 3$  and  $d = 2$ , so that  $\Theta = [0, 1]^4$ . Let  $\theta_0 = (b_{11}, b_{21}, b_{12}, b_{22}) = (0, 0.8, 0.6, 0)$ . We test  $H_0 : (b_{11}, b_{22}) = (0, 0)$ . Assumption (A4) holds with  $\Lambda = [0, \infty) \times \mathbb{R}^2 \times [0, \infty)$ . If the null hypothesis cannot be rejected, it means that a Marshall–Olkin model suffices.

Fig. 3 shows the RMSE of the parameter estimates of the three max-linear models explained above, based on the empirical tail dependence function (solid lines) and the beta tail dependence function (dashed lines). We see again that these two initial estimators lead to similar results. Parameters whose true values are on the boundary are better estimated for low  $k$ , while higher values of  $k$  are preferred for parameters whose true values are in the interior of the parameter space.

Table 2 shows the empirical level of the test statistics  $T_{n,k}^{(1)}$  and  $T_{n,k}^{(2)}$  for  $k \in \{25, 50, 75, 100\}$ , using a significance level of 0.05. The tests perform well for model 1 in general, while low values of  $k$  are necessary for models 2 and 3. The beta tail dependence function outperforms the empirical tail dependence function for model 2, while the opposite is the case for model 3. Overall, test statistic  $T_{n,k}^{(1)}$  performs somewhat better than  $T_{n,k}^{(2)}$ .

To assess the power of the two test statistics, we consider the models described above, but we let

$$\begin{aligned} b_{11} &\in \{0.8, 0.85, \dots, 1\} \quad \text{for model 1,} \\ b_{11} &\in \{0.8, 0.85, \dots, 1\}, \quad b_{31} \in \{0, 0.05, \dots, 0.2\} \quad \text{for model 2, and} \\ b_{11}, b_{22} &\in \{0, 0.05, \dots, 0.2\} \quad \text{for model 3.} \end{aligned}$$

Fig. 4 shows the empirical power of  $T_{n,k}^{(1)}$  for models 1–3 for the beta tail dependence function (results for the empirical tail dependence function are very similar). The most striking result of the lower-left panel is that the graph is not symmetric, i.e., the power is higher when  $b_{22}$  is near zero (and  $b_{11}$  is not) than when  $b_{11}$  is near zero (and  $b_{22}$  is not).

##### 4.2.2. Testing the number of factors in a max-linear model

We studied the performance of the test statistics in case we wanted to identify a specific submodel of the max-linear model. We would also like to investigate the capability of the test statistics to correctly retrieve the true number of factors, i.e., for  $s \in \{1, \dots, r\}$ , we wish to test the hypothesis  $H_0 : (b_{1s}, \dots, b_{ds}) = \mathbf{0}$ . However, as mentioned in Section 2.1, the max-linear model is only defined for parameter values with  $\sum_{j=1}^d b_{js} > 0$  for all  $s \in \{1, \dots, d\}$ ; when this condition is not met, there are zeroes on the diagonal of  $\Sigma(\theta_0)$ , making computation of  $\hat{\lambda}_\beta$  impossible. We solve this problem by computing




Fig. 3. RMSE for models 1–3 based on the empirical tail dependence function (solid lines), and the beta tail dependence function (dashed lines).

Table 3


Empirical level of  $T_{n,k}^{(2)}$  based on the empirical and the beta tail dependence function for a significance level of 0.05.

	$k = 25$		$k = 50$		$k = 75$		$k = 100$	
	emp	beta	emp	beta	emp	beta	emp	beta
$T_{n,k}^{(2)}$	4.6	23.9	2.7	24.9	2.4	23.8	2.5	20.0


the asymptotic distribution of the test statistics using  $\hat{\theta}_{n,k}$  rather than  $\theta_0$ . We consider a model with  $d = 2$  and  $r = 3$ , with parameter vector  $\theta_0 = (b_{11}, b_{21}, b_{12}, b_{22}) = (0.8, 0.6, 0.2, 0.4)$ , i.e., the third column of  $B$  contains only zeroes and the model has effectively two factors. We estimate a three-factor model and we test  $H_0 : (b_{13}, b_{23}) = (0, 0)$ .

Fig. 5 shows the RMSE of the parameter estimates based on the empirical tail dependence function (solid lines) and the beta tail dependence function (dashed lines). We see that, contrary to previous experiments, the beta tail dependence has much higher RMSE than the empirical tail dependence function: oversmoothing leads to a large bias for  $(b_{13}, b_{23})$  (and hence for the other parameters as well).

Table 3 shows the empirical level of the test statistic  $T_{n,k}^{(2)}$  using a significance level of 0.05. We included the results based on the beta tail dependence function for completeness: because it tends to overestimate  $(b_{13}, b_{23})$ , it rejects the null hypothesis far too often. Test statistic  $T_{n,k}^{(1)}$  is not presented because for any  $k$  and any initial estimator, it has an empirical level of 0. Even when  $|\hat{\theta}_{n,k} - \hat{\theta}_{n,k}^{(0)}|$  is large,  $|f_{n,k}(\hat{\theta}_{n,k}) - f_{n,k}(\hat{\theta}_{n,k}^{(0)})|$  is small and hence the deviance-type test is not adapted to this type of model. Finally, Fig. 6 shows the empirical power of  $T_{n,k}^{(2)}$  for  $b_{11}, b_{22} \in \{0, 0.05, \dots, 0.2\}$  for two values of  $k$ .



**Fig. 4.** Top: empirical power in % for model 1 and  $k = 50$  (dashed line),  $k = 75$  (solid line), based on the empirical tail dependence function. Bottom: empirical power in % for model 2 (left) and model 3 (right) and  $k = 50$  based on the beta tail dependence function.



**Fig. 5.** RMSE based on the empirical tail dependence function (solid lines) and the beta tail dependence function (dashed lines).

We remark that the power is similar when one parameter is far from its boundary and when both parameters are far from their boundary.

## 5. Application to stock market indices

Consider two major European stock market indices, the German DAX and the French CAC40. We take the daily negative log-returns of the prices of these two indices from <https://finance.yahoo.com> for the period of January 1st, 1997 to December 31st, 2017. We remove all dates for which at least one of the two series has missing values, ending up with a sample of size  $n = 5313$ . Fig. 7 shows the time series plots of log-returns and the dependence structure for the returns standardized to unit Pareto margins, plotted on the exponential scale. The time series plots look stationary: we perform a Augmented



Power of  $T_{n,k}^{(2)}$  based on empirical stdf,  $k = 50$ Power of  $T_{n,k}^{(2)}$  based on empirical stdf,  $k = 75$ 

Fig. 6. Empirical power in % of  $T_{n,k}^{(2)}$ , based on the empirical tail dependence function for  $k = 50$  (left) and  $k = 75$  (right).




Fig. 7. Left and middle: time series plots of daily negative log-returns of the DAX and the CAC40; right: scatterplot of daily negative log-returns of the DAX versus the CAC40 on the unit Pareto scale, plotted on the exponential scale.

Table 4

Parameter matrix for a bivariate max-linear model with  $r = 4$  factors for  $k = 40$ ; standard errors are in parentheses.

DAX	0.41 (0.11)	0.14 (0.05)	0.43 (0.11)	0.03 (0.05)
CAC40	0.43 (0.12)	0.44 (0.10)	0.13 (0.05)	0.00 (0.01)


Dickey–Fuller test, which gives  $p$ -values below 0.01 for both series, and a KPSS test, giving  $p$ -values above 0.1 for both series. Hence, the assumption of stationarity is reasonable. Let  $u_j$  denote the 0.95% quantile of  $X_{1j}, \dots, X_{nj}$  for  $j \in \{1, \dots, d\}$ . We fit a generalized Pareto distribution (GPD) to  $X_{ij} - u_j \mid X_{ij} > u_j$  and obtain the parameter estimates  $\hat{\sigma}_1 = 1.12$  (0.11),  $\hat{\gamma}_1 = 0.03$  (0.08),  $\hat{\sigma}_2 = 1.10$  (0.10) and  $\hat{\gamma}_2 = 0.02$  (0.07).

Table 4 shows the estimated parameter matrix  $B = (b_{jt})_{j,t}$  for  $j = 1, 2$  and  $t = 1, 2, 3, 4$  obtained by fitting a four-factor max-linear model to our data based on the empirical tail dependence function. Values of  $k \in \{25, 30, \dots, 70, 75\}$  were considered: we chose  $k = 40$  because parameter estimates are stable around this value. Standard errors were calculated using the asymptotic variance matrix of the estimator. The results in Table 4 suggest to test  $H_0 : (b_{14}, b_{24}) = (0, 0)$ ; the value of the test statistic  $T_{n,k}^{(2)}$  is 0.04. Comparing with a critical value of 17.4 based on a significance level of 0.05, we find that we cannot reject the null hypothesis. Critical values are calculated by simulation from the asymptotic distribution of the test statistics, which is given in Corollaries 3.4 and 3.5. We do not consider  $T_{n,k}^{(1)}$  or the beta tail dependence function because of their bad performance (see Section 4.2.2).

**Table 5**

Parameter matrix for a bivariate max-linear model with  $r = 3$  factors for  $k = 40$ ; standard errors are in parentheses.

DAX	0.41 (0.11)	0.14 (0.06)	0.46 (0.09)
CAC40	0.43 (0.12)	0.44 (0.10)	0.13 (0.05)



**Fig. 8.** Left: level sets  $\{(x_1, x_2) : \ell(x_1, x_2) = c\}$  for  $c \in \{0.2, 0.4, 0.8, 1\}$  based on the empirical tail dependence function (solid lines) and on the fitted max-linear model (dashed lines); middle: nonparametric (black dots) and model-based (red dots) estimates of the tail dependence coefficient  $\hat{\chi}_{n,k}$  for  $k \in \{200, 195, \dots, 30\}$ ; right: probability of a joint exceedance based on the fitted max-linear model (solid line) and on the empirical excesses (dashed line). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Table 5 shows the estimated parameter matrix  $B = (b_{jt})_{j,t}$  for  $j = 1, 2$  and  $t = 1, 2, 3$  for a three-factor model. We test if a Marshall–Olkin model suffices, i.e.,  $H_0 : (b_{12}, b_{23}) = (0, 0)$ . The value of the test statistic  $T_{n,k}^{(2)}$  is 75.9. Comparing to a critical value of 8.03, we reject the null hypothesis of a Marshall–Olkin model.

Fig. 8 shows two goodness-of-fit measures and the estimated probability of a joint exceedances of the two stocks. On the left, we plotted the level sets  $\{(x_1, x_2) : \ell(x_1, x_2) = c\}$  for  $c \in \{0.2, 0.4, 0.8, 1\}$  based on the empirical tail dependence function (solid lines) and on the fitted max-linear model (dashed lines).

A common summary measure of dependence is the tail dependence coefficient,

$$\chi(q) := \lim_{q \uparrow 1} \mathbb{P}[F_1(X_{11}) < q \mid F_2(X_{12}) < q] = 2 - \ell(1, 1),$$

see (2). Fig. 8 (middle) shows nonparametric estimates of the tail dependence coefficient (black dots)  $\tilde{\chi}_{n,k} = 2 - \tilde{\ell}_{n,k}(1, 1)$  and its model-based counterpart (red dots) for decreasing  $k$ . The horizontal grey line corresponds to the value  $k = 40$  that was used for the parameter estimators and the tests, corresponding to  $\hat{\chi}_{n,k} \approx 0.68$ , which is equal to the model-based  $\chi$ . The dotted lines correspond to 95% pointwise bootstrap confidence intervals.

The probability of a joint excess  $\bar{F}(x_1, x_2) = \mathbb{P}[X_{11} > x_1, X_{12} > x_2]$  can be approximated for large  $(x_1, x_2)$  by

$$\bar{F}(x_1, x_2) = x_1^* + x_2^* - 1 + \exp\{-\ell(x_1^*, x_2^*)\} \quad \text{for } x_j^* = 1 - F_j(x_j), \quad j = 1, 2.$$

Let  $\bar{H}(\cdot; \gamma_j, \sigma_j)$  denote the survival function of a GPD with shape  $\gamma_j$  and scale  $\sigma_j$ . For  $x_1 > u_1$  and  $x_2 > u_2$ , we can estimate  $\bar{F}(x_1, x_2)$  by

$$\hat{\bar{F}}(x_1, x_2) = \hat{x}_1^* + \hat{x}_2^* - 1 + \exp\left\{-\ell(\hat{x}_1^*, \hat{x}_2^*; \hat{\theta}_{n,k})\right\},$$

where  $\ell(\cdot; \hat{\theta}_{n,k})$  denotes the max-linear stable tail dependence function based on the fitted parameter estimates and  $\hat{x}_j^* = 0.05 \times \bar{H}(x_j - u_j; \hat{\gamma}_j, \hat{\sigma}_j)$ . The right-hand plot of Fig. 7 shows the probabilities  $\hat{\bar{F}}(x, x)$  on the log-scale for  $x \in \{2.5, 3, \dots, 12\}$  (solid line). These can be compared to the empirical excess probabilities  $n^{-1} \sum_{i=1}^n \mathbb{1}\{X_{1i} > x, X_{2i} > x\}$  (dashed line), which are zero for  $x > 8.5$ .

## 6. Discussion

We proposed two test statistics for parameters that, under the null hypothesis, are on the boundary of the alternative hypothesis. These test statistics are convenient because their asymptotic distribution has an explicit expression, and have been shown to perform well in simulation studies. In practice, their asymptotic distribution becomes cumbersome when testing for a higher dimensional ( $c > 4$ ) parameter vector. A possible solution might be to consider a multiple testing procedure with a Bonferroni-type of correction.

We applied the test statistics to the max-linear model and the Brown–Resnick model, but the results are generic and could be used on any multivariate extreme-value model where submodels appear at boundary values or where the number of “factors” is of interest. Examples include Fougères et al., (2009), where a mixture model is obtained based on a certain number of stable random variables, and Gissibl and Klüppelberg (2018), where a max-linear model is defined on a directed acyclic graph. For the latter, our methods could be useful in the procedure to reconstruct the graph structure of a dataset (Gissibl et al., 2018), since it is based on identifying pairwise tail dependence coefficients that are zero.

## Acknowledgements

The author would like to thank two reviewers and an associate editor for their careful reading of the paper and their constructive comments that greatly improved the generality of the paper. She would also like to thank Johan Segers for helpful comments on an earlier version of this paper.

## Appendix A. Proofs

**Definition** (Left/right partial derivatives). Let  $f$  be a function whose support includes  $\mathcal{X} \subset \mathbb{R}^p$  and let  $\mathbf{a} \in \mathcal{X}$ . Suppose that  $\mathcal{X} - \mathbf{a}$  equals the intersection of a union of orthants and an open cube  $C_\varepsilon(\mathbf{0})$  for some  $\varepsilon > 0$ , i.e.,  $\mathcal{X} - \mathbf{a}$  is locally equal to a union of orthants. The function  $f$  is said to have *left/right (l/r) partial derivatives of order 1 on  $\mathcal{X}$*  if

1. it has partial derivatives at each interior point of  $\mathcal{X}$ ;
2. it has partial derivatives at each boundary point of  $\mathcal{X}$  with respect to coordinates that can be perturbed to the left and right;
3. it has left (right) partial derivatives at each boundary point of  $\mathcal{X}$  with respect to coordinates that can be perturbed only to the left (right).

The shape of  $\mathcal{X}$  is such that for all  $\mathbf{x} \in \mathcal{X}$  and for all  $i \in \{1, \dots, p\}$ , it is possible to perturb  $x_i$  to the left, the right, or both and stay within  $\mathcal{X}$ . This means that it is always possible to define the left, right, or two-sided partial derivative of  $f$  with respect to  $x_i$ . We say that  $f$  has l/r partial derivatives of order  $k$  on  $\mathcal{X}$  for  $k \geq 2$  if  $f$  has l/r partial derivatives of order  $k - 1$  on  $\mathcal{X}$  and each of the latter has l/r partial derivatives on  $\mathcal{X}$ . When we say the  $f$  has continuous l/r partial derivatives, continuity is defined in terms of local perturbations within  $\mathcal{X}$  only.

**Proof of Theorem 3.2.** This theorem is a special case of Andrews (1999, Theorem 3b). A closely related work is Andrews (2002), where the focus is on generalized method of moment estimators. This setting is closer to ours but uses a convergence rate of  $\sqrt{n}$ , whereas (Andrews, 1999) allows for more generality. The quantities  $\ell_T$ ,  $B_T$  and  $R_T$  in Andrews (1999) correspond to  $-(k/2)f_{n,k}$ ,  $\sqrt{k}I_p$  and  $-(k/2)R_{n,k}$  respectively in this paper. Theorem 3b in Andrews (1999) holds under the assumptions 1, 2\*, 3\*, 5\*, and 6 of that paper. We show that these are implied by ours:

**Assumption 1** The assumptions made in Theorem 3.1 imply assumption 1.

**Assumption 2\*** Assumption GMM2 in Andrews (2002) implies Assumption 2\*; this is proven in Andrews (2002, Lemma 3). We show that our assumption imply GMM2: assumption GMM2(a) holds because  $D_{n,k}(\boldsymbol{\theta})$  converges in probability to  $L(\boldsymbol{\theta}_0) - L(\boldsymbol{\theta})$ . Assumption GMM2(b) holds if assumption GMM2\*(b) holds, which in turn is implied by the assumptions of Theorem 3.1 and (A1). Assumption GMM2(c) holds since  $\mathbf{D}(\boldsymbol{\theta}_0) = \mathbf{0}$ . Assumption GMM2(d) holds since  $D_{n,k}(\boldsymbol{\theta}) - \mathbf{D}(\boldsymbol{\theta}) - D_{n,k}(\boldsymbol{\theta}_0) = \mathbf{0}$ . Finally, Assumption GMM2(e) holds automatically since our weight matrix isn't random.

**Assumption 3\*** Assumption 3\* holds because of (A3) and because  $J$  is non-random, symmetric and non-singular.

**Assumption 5\*** Assumption 5\* holds because of (A2) and because  $B_T = \sqrt{k}I_p$  and  $\sqrt{k} \rightarrow \infty$  as  $n \rightarrow \infty$ .

**Assumption 6** Assumption (A2) implies assumption 6.  $\square$

**Proof of Corollary 3.3.** This corollary is a special case of Andrews (1999, Corollary 1b), where no parameter  $\psi$  appears. Assumptions 1, 2\*, 3\*, 5\* and 6–8 in that paper are needed: in the proof of Theorem 3.2, we've already shown that our assumptions imply 1, 2\*, 3\*, 5\* and 6; assumptions 7 and 8 hold by our assumption (A4).  $\square$

## References

- Andrews, D.W.K., 1999. Estimation when a parameter is on a boundary. *Econometrica* 67, 1341–1383.
- Andrews, D.W.K., 2001. Testing when a parameter is on the boundary of the maintained hypothesis. *Econometrica* 69, 683–734.
- Andrews, D.W.K., 2002. Generalized method of moments estimation when a parameter is on a boundary. *J. Bus. Econ. Stat.* 20, 530–544.
- Balkema, A.A., De Haan, L., 1974. Residual life time at great age. *Ann. Probab.* 2, 792–804.
- Beirlant, J., Escobar-Bach, M., Goegebeur, Y., Guillou, A., 2016. Bias-corrected estimation of stable tail dependence function. *J. Multivar. Anal.* 143, 453–466.
- Berghaus, B., Bücher, A., Dette, H., 2013. Minimum distance estimators of the Pickands dependence function and related tests of multivariate extreme-value dependence. *Journal de la Société Française de Statistique* 154, 116–137.
- Brigo, D., Mai, Jan, Scherer, M., Sloom, H., 2018. Consistent iterated simulation of multivariate defaults: Markov indicators, lack of memory, extreme-value copulas, and the Marshall–Olkin distribution. *Innovations in Insurance, Risk- and Asset Management*. World Scientific Publishing Co.
- Burtschell, X., Gregory, J., Laurent, J.P., 2009. A comparative analysis of CDO pricing models. *J. Deriv.* 16, 9–37.
- Capéraà, P., Fougères, A.L., Genest, C., 1997. A nonparametric estimation procedure for bivariate extreme value copulas. *Biometrika* 84, 567–577.



- Castruccio, S., Huser, R., Genton, M.G., 2016. High-order composite likelihood inference for max-stable distributions and processes. *J. Comput. Graph. Stat.* 25, 1212–1229.
- Chavez-Demoulin, V., Embrechts, P., Hofert, M., 2016. An extreme value approach for modeling operational risk losses depending on covariates. *J. Risk Insur.* 83, 735–776.
- Cui, Q., Zhang, Z., 2018. Max-linear competing factor models. *J. Bus. Econ. Stat.* 36, 62–74.
- Davison, A.C., Padoan, S.A., Ribatet, M., 2012. Statistical modeling of spatial extremes. *Stat. Sci.* 27, 161–186.
- Dombry, C., Engelke, S., Oesting, M., 2017. Bayesian inference for multivariate extreme value distributions. *Electron. J. Stat.* 11, 4813–4844.
- Draisma, G., Drees, H., Ferreira, A., De Haan, L., 2004. Bivariate tail estimation: dependence in asymptotic independence. *Bernoulli* 10, 251–280.
- Drees, H., Huang, X., 1998. Best attainable rates of convergence for estimators of the stable tail dependence function. *J. Multivar. Anal.* 64, 25–47.
- Einmahl, J.H., Kiriliouk, A., Krajina, A., Segers, J., 2016. An M-estimator of spatial tail dependence. *J. Royal Stat. Soc.: Ser. B (Stat. Methodol.)* 78, 275–298.
- Einmahl, J.H., Kiriliouk, A., Segers, J., 2018. A continuous updating weighted least squares estimator of tail dependence in high dimensions. *Extremes* 21, 205–233.
- Einmahl, J.H., Magnus, J.R., 2008. Records in athletics through extreme-value theory. *J. Am. Stat. Assoc.* 103, 1382–1391.
- Einmahl, J.H., Krajina, A., Segers, J., 2012. An M-estimator for tail dependence in arbitrary dimensions. *Ann. Stat.* 40, 1764–1793.
- Embrechts, P., Lindskog, F., McNeil, A., 2003. Modelling dependence with copulas and applications to risk management. In: Rachev, S. (Ed.), *Handbook of Heavy Tailed Distributions in Finance* chapter 8. Elsevier, pp. 329–384.
- Falk, M., Hüslér, J., Reiss, R.D., 2010. *Laws of Small Numbers: Extremes and Rare Events*. Springer Science & Business Media.
- Fisher, R.A., Tippett, L.H.C., 1928. Limiting forms of the frequency distribution of the largest or smallest member of a sample. In: *Mathematical Proceedings of the Cambridge Philosophical Society*, (pp. 180–190) volume 24. Cambridge University Press.
- de Fondeville, R., Davison, A.C., 2018. High-dimensional peaks-over-threshold inference. *Biometrika* 105, 575–592.
- Fougères, A.L., de Haan, L., Mercadier, C., 2015. Bias correction in multivariate extremes. *Ann. Stat.* 43, 903–934.
- Fougères, A.L., Nolan, J.P., Rootzén, H., 2009. Models for dependent extremes using stable mixtures. *Scand. J. Stat.* 36, 42–59.
- Gissibl, N., Klüppelberg, C., Otto, M., 2018. Tail dependence of recursive max-linear models with regularly varying noise variables. *Econom. Stat.* 6, 149–167.
- Gissibl, N., Klüppelberg, C., 2018. Max-linear models on directed acyclic graphs. *Bernoulli* 24, 2693–2720.
- Gnedenko, B., 1943. Sur la distribution limite du terme maximum d'une serie aleatoire. *Ann. Math.* 22, 423–453.
- Gudendorf, G., Segers, J., 2012. Nonparametric estimation of multivariate extreme-value copulas. *J. Stat. Plan. Inference* 142, 3073–3085.
- Guillou, A., Padoan, S.A., Rizzelli, S., 2018. Inference for asymptotically independent samples of extremes. *J. Multivar. Anal.* 167, 114–135.
- Gumbel, E.J., 1960. Bivariate exponential distributions. *J. Am. Stat. Assoc.* 55, 698–707.
- de Haan, L., Ferreira, A., 2006. *Extreme Value Theory: an Introduction*. Springer-Verlag Inc.
- Huang, X., 1992. *Statistics of Bivariate Extreme Values*. Tinbergen Institute Research Series (Ph.D. thesis).
- Huser, R., Davison, A., 2013. Composite likelihood estimation for the Brown–Resnick process. *Biometrika* 100, 511–518.
- Hüslér, J., Li, D., 2009. Testing asymptotic independence in bivariate extremes. *J. Stat. Plan. Inference* 139, 990–998.
- Kabluchko, Z., Schlather, M., de Haan, L., 2009. Stationary max-stable fields associated to negative definite functions. *Ann. Probab.* 37, 2042–2065.
- Kiriliouk, A., Rootzén, H., Segers, J., Wadsworth, J.L., 2018a. Peaks over thresholds modelling with multivariate generalized Pareto distributions. *Technometrics* 61, 123–135.
- Kiriliouk, A., Segers, J., Tafakori, L., 2018b. An estimator of the stable tail dependence function based on the empirical beta copula. *Extremes* 21, 581–600.
- Ledford, A.W., Tawn, J.A., 1996. Statistics for near independence in multivariate extreme values. *Biometrika* 83, 169–187.
- Marcon, G., Padoan, S., Naveau, P., Muliere, P., Segers, J., 2017. Multivariate nonparametric estimation of the Pickands dependence function using Bernstein polynomials. *J. Stat. Plan. Inference* 183, 1–17.
- Padoan, S., Ribatet, M., Sisson, S., 2010. Likelihood-based inference for max-stable processes. *J. Am. Stat. Assoc. (Theory Methods)* 105, 263–277.
- Pickands III, J., 1975. Statistical inference using extreme order statistics. *Ann. Stat.* 3, 119–131.
- Rootzén, H., Tajvidi, N., 2006. Multivariate generalized Pareto distributions. *Bernoulli* 12, 917–930.
- Rootzén, H., Zholud, D., 2017. Human life is unlimited—but short. *Extremes* 20, 713–728.
- Segers, J., 2012. Max-stable models for multivariate extremes. *REVSTAT — Stat. J.* 10, 61–92.
- Segers, J., Sibuya, M., Tsukahara, H., 2017. The empirical beta copula. *J. Multivar. Anal.* 155, 35–51.
- Smith, R.L., 1989. Extreme value analysis of environmental time series: an application to trend detection in ground-level ozone. *Stat. Sci.* 4, 367–377.
- Smith, R.L., 1990. Max-stable processes and spatial extremes. Unpublished manuscript.
- Su, J., Furman, E., 2017. Multiple risk factor dependence structures: copulas and related properties. *Insur.: Math. Econ.* 74, 109–121.
- Tawn, J.A., 1990. Modelling multivariate extreme value distributions. *Biometrika* 77, 245–253.
- Thomas, M., Lemaitre, M., Wilson, M.L., Viboud, C., Yordanov, Y., Wackernagel, H., Carrat, F., 2016. Applications of extreme value theory in public health. *PLoS One* 11, e0159312.
- Towler, E., Rajagopalan, B., Gilleland, E., Summers, R.S., Yates, D., Katz, R.W., 2010. Modeling hydrologic and water quality extremes in a changing climate: a statistical approach based on extreme value theory. *Water Resour. Res.* 46, 1–11.
- Vettori, S., Hüser, R., Genton, M.G., 2016. A comparison of dependence function estimators in multivariate extremes. *Statistics and Computing* 28, 525–538.
- Wadsworth, J., Tawn, J., Davison, A., Elton, D., 2016. Modelling across extremal dependence classes. *J. Royal Stat. Soc.: Ser. B (Stat. Methodol.)* 79, 149–175.
- Wadsworth, J.L., Tawn, J.A., 2014. Efficient inference for spatial extreme-value processes associated to log-Gaussian random functions. *Biometrika* 101, 1–15.
- Zhang, D., Wells, M.T., Peng, L., 2008. Nonparametric estimation of the dependence function for a multivariate extreme value distribution. *J. Multivar. Anal.* 99, 577–588.
- Zhao, Z., Zhang, Z., 2018. Semiparametric dynamic max-copula model for multivariate time series. *J. Royal Stat. Soc.: Ser. B (Stat. Methodol.)* 80, 409–432.