



## THESIS / THÈSE

### MASTER IN BIOLOGY OF ORGANISMS AND ECOLOGY

#### Impact of low temperature during embryonic development on the sex determination and de novo DNA methyltransferases in the androdiecious fish *Kryptolebias marmoratus*

Renard, Tiana

*Award date:*  
2022

*Awarding institution:*  
University of Namur  
Université Catholique de Louvain

[Link to publication](#)

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Université catholique de Louvain  
École de biologie

Université de Namur  
Département de biologie



---

Impact of low temperature during embryonic development on the sex determination and *de novo* DNA methyltransferases in the androdieicious fish *Kryptolebias marmoratus*

Renard Tiana

Mémoire présenté en vue de l'obtention du diplôme de  
Master en Biologie des Organismes et Ecologie

Promoteur : Frédéric Silvestre (UNamur)  
Encadrant : Yves Blanco-Alvarez (UNamur)

Année académique  
2021-2022

# **Impact of low temperature during embryonic development on the sex determination and *de novo* DNA methyltransferases in the androdiecious fish *Kryptolebias marmoratus***

## **ABSTRACT**

The teleost fish *Kryptolebias marmoratus* (mangrove rivulus) presents an unusual reproductive system as it is one of only two vertebrates in which hermaphroditic individuals can reproduce by self-fertilization but also outcross with males. Between different populations of rivulus, sex ratios are variable as sex determination processes are based on genetic and environmental factors. In laboratory, isogenic lineages that avoid genetic diversity can be produced, allowing to study the effect of environment only. Even though low temperature is known to impact the sex ratio by inducing male development, and epigenetic mechanisms are suspected to make the connection between environment and sexual gene expression, most of the mechanisms behind sex determination of *K. marmoratus* by temperature are still unknown. To better understand these processes, we compared the expression level of the *de novo* DNA methyltransferases dnmt3aa and dnmt3ab as well as those of the sexual genes sox9a and cyp19b in rivulus embryos incubated at 18°C and 25°C. Even though we could not show a correlation between dnmt3a and sexual genes expression, we found that, while dnmt3aa and dnmt3ab are similarly expressed for a same temperature, the level expression of both dnmt3a variants seems to increase at a faster rate at 18°C compared to 25°C, at least during the early stages of embryonic development. In parallel, we observed that in the last stages of embryonic development, sox9a was overexpressed at 18°C while cyp19b was overexpressed as 25°C. This study may be a starting point for further research to prove a correlation, or even a causality, between *de novo* methyltransferases and sexual genes as a pathway to explain temperature-dependent sex determination in mangrove rivulus.

**Key words:** androdioecy; DNA methylation; *Kryptolebias marmoratus*; masculinization; sex determination

## 1. Introduction

Sexual reproduction is an ancient and intrinsic feature of eukaryotic life (Speijer et al., 2015). Yet, sexual systems, i.e. the pattern of distribution of the male and female function among the individuals of a given species, are diverse and complex (Bachtro et al., 2014; Pla et al., 2022). While common systems, like gonochorism or hermaphroditism, are well known, androdioecy, i.e., the presence of male and hermaphroditic individuals, is rare and still poorly understood (Weeks et al., 2006; Moreira-Pinto & Pires-daSilva, 2018; Pla et al., 2022). The first cases of documented androdiecious species only appeared in the '60s – '70s, mainly in plants (Charlesworth, 1984). Since then, other examples of androdiecious species have been found in plants and animals, especially in nematodes and crustaceans (Pannell, 2002; Weeks, 2012; Subranomian, 2017; Moreira-Pinto & Pires-daSilva, 2018). Even though this sexual pattern appears to be restricted taxonomically, extended research on androdiecious metazoans provided evidence that it has evolved many.(Weeks 2012; Leonard, 2018). The most common form of androdioecy in animals, which seems to have evolved from dioecy, involves hermaphrodites that either self-fertilize or outcross with males, like the nematode *Caenorhabditie elegans* and the fish *Kryptolebias marmoratus* (Harrington, 1967; Weeks 2012; Leonard, 2018; Moreira-Pinto & Pires-daSilva, 2018).




Figure 1 - Representatives of the three sexual phenotypes encountered in *K. marmoratus*. full bodies, caudal fin, and ocellus beneath. Bottom panels show a ventral view of a dissected animal. The ocellus is shown at the red arrow for the hermaphrodite (Marson *et al.*, 2019)

*Kryptolebias marmoratus* (Poey, 1880), also called mangrove rivulus or mangrove killifish, is a teleost fish mostly known for being one of the only androdieocious vertebrates (Harrington, 1960). This species is characterized by self-fertilizing hermaphrodites, primary males, which directly develop testicular tissues, and secondary males, which correspond to hermaphroditic adults characterized by a resorption of the ovarian tissue and a colour change (Figure 1) ((Weeks et al., 2006; Kelley et al., 2016; Marson *et al.*, 2019). Present in the tropical and subtropical West Atlantic, the rivulus distribution range extent from Florida to the north-eastern coast of Brazil, through the Yucatan Peninsula and the Bay of Campeche in Mexico (Taylor, 2012; Costa, 2016). This species, associated to mangroves, can colonize a large variety of micro-habitat, such as shallow water, crab burrows or mangrove litter (Tayler, 2000; Taylor *et al.*, 2008). The great phenotypic plasticity of the rivulus allows it to thrive in the large range of physico-chemical parameters like mangroves that are a highly variable habitat in time and space, (Taylor, 2012). The hard living conditions might have led to low density populations during *K. marmoratus* life history, and thus mating difficulties. Androdioecy could have appeared as a potential strategy of reproductive assurance, as self-fertilizing individuals do not have to find a mate anymore and can produce almost homozygotic populations, which can be highly adapted to their environment, as multiple generations of selfing, starting from a heterozygote individual, can lead to the fixation of sets of alleles that are particularly advantageous in a specific biotope. (Taylor et al., 2001; Weeks; 2012; Ellison et al., 2013). On the other hand, rivulus populations that have low genetic diversity could be more sensitive to environmental variation as less polymorphism might prevent the emergence of new phenotypes in case of changing environments (Tatarenkov et al., 2012). In the wild, not all population of rivulus are isogenic. Sex ratio, which is the result of complex interactions between genetic and environmental factors, finally resulting in the expression of the hermaphroditic or male phenotype at the individual level, is dependent on the population localization and varies from 1-2% to 45% (Turner et al., 1990; Taylor et al., 2001; Pers. com. Chapelle & Silvestre). The sexual phenotype is influenced by the developmental plasticity of the rivulus, as primary males are induced by incubation temperatures of around 18-20°C and hermaphrodites by temperatures of around 25-29°C, but also by the phenotypic flexibility in adults, as hermaphrodites can turn into secondary males due to environmental factors (Harrington, 1967; Harrington & Kallman, 1968; Taylor, 2012; Ellison *et al.*, 2015; Kelley *et al.*, 2016). As more heterozygous individuals lead to a lower chance of inbreeding depression, sex ratio plays an important role in population fitness, and thus, species evolution (Taylor et al., 2001; Piferrer *et*

*al.*, 2012; Ellison *et al.*, 2013). Unravelling the mechanisms behind sex determination for *K. marmoratus* is thus essential to better understand its evolution.

Sex determination (SD) can be defined as the physiological processes that lead to the differentiation of the reproductive system under the influence of genetic or environmental factors (Hayes, 1998; Sandra & Norma, 2010; Santi *et al.*, 2016). SD is fundamental for sexual reproduction, species survival and evolution. As one of the most plastic evolutionary processes, many different mechanisms exist (Haag & Doty, 2005; Capel, 2017; Bertho *et al.*, 2018). In vertebrates alone, SD mechanisms are highly variable, ranging from strict genotypic sex determination (GSD) to strict environmental sex determination (ESD), including combinations of genetic and environmental determination (GSD+ESD) (Penman & Piferrer, 2008; Barchtog *et al.*, 2014; Capel, 2017). At first, sex determination was thought to be strictly control by genetic (Barchtog *et al.*, 2014). Most vertebrates have either a XX/XY-system in which the gene SRY, located on Y-male chromosome, act as a trigger for testis formation (mammals), or a ZZ/ZW-system in which the overexpression of the Z-linked gene *dmrt1* leads to male sex determination (birds) (Matsuda *et al.*, 2002; Smith *et al.*, 2009; Yoshimoto & Ito, 2011; Barchtog *et al.*, 2014; Li & Gui, 2018). In fish and other non-mammal vertebrates, *dmrt1* is involved in testis formation (Nagahama, 2005; Barchtog *et al.*, 2014; Capel, 2017). In certain species of fish, reptiles, and amphibians, rather than GSD, the incubation temperature of the eggs induced the sex of the progeny. For these species, sex is strictly determined by the environment, and in most cases by the temperature (Korpelainen, 1990; Ospina-Alvarez & Piferrer, 2008). Other species present a gradient between GSD and ESD. Environmental parameters, especially temperature, can override genotypic control (Barchtog *et al.*, 2014; Li & Gui, 2018). For example, it has been shown that, in the Nile Tilapia *Oreochromis niloticus*, high temperatures lead to the masculinization of the embryos, resulting in XX males (Baroiller, 1995; Baroiller *et al.*, 2009a). Sex determination is mainly studied in gonochoric species, but combination between GSD and ESD also exists in other sexual systems. In our model, *K. marmoratus*, masculinization is induced by low temperature and a modification of the level of expression of the sexual genes *cyp19a*, associated with the ovarian development, and *sox9a*, associated with the development of testicular tissues (Ellison *et al.*, 2015). This is also the case for the Nile tilapia in which temperature regulates the expression level of genes included in sex determination pathways, and induces male phenotype through DNA methylation (Baroiller *et al.*, 2009a; Baroiller *et al.*, 2009b)

DNA methylation is an epigenetic mechanism that seems to bridge the gap between GSD and ESD (Ellison *et al.*, 2015; Fellous *et al.*, 2018). Epigenetics examines heritable

changes in gene expression that occur without modification of the DNA sequence (Berger *et al.*, 2009). Among epigenetic mechanisms, histone modifications and DNA methylation seem to be involved in the regulation of sex-related gene expression by the environment. Histone modifications regulate gene transcription by altering the structure of the chromatin (Bollati & Baccarelli, 2010; Bannister & Kouzarides 2011). For example, histone demethylase *kdm6B* regulates temperature-dependent sex determination in the red-eared slider turtle *Trachemys scripta elegans*. In this species, incubation temperature determines the turtle's sex. When *kdm6b* is knocked down and the embryos incubated at 26°C, females are sired instead of the expected males (Ge *et al.*, 2018). In the same species, DNA methylation of *dmrt1* is also involved in the sex determination by temperature during embryo development (Ge *et al.*, 2017). As we saw before, DNA methylation is also involved in sex determination of fish. More precisely, masculinization by high temperature in the Nile tilapia is correlated with an increase of the DNA methylation level of the sexual gene *cyp19a1a* and a reduction of its expression level (Wang *et al.*, 2017). DNA methyltransferases (dnmts) catalyze the transfer of a methyl-group onto a cytosine, which usually prevents transcription and thus reduces or even suppresses gene expression (Moore & Fan, 2013). DNA methylation is also thought to be involved in sex determination in *K. marmoratus*. Ellison *et al.* (2015) have shown that DNA methylation could be the intermediary through which temperature modulates sexual identity. In the rivulus, *dnmt3a* variants, responsible for *de novo* methylation, were shown to be differentially expressed between in male and hermaphrodite adult tissues, with *dnmt3ab* overexpressed in males and *dnmt3aa* overexpressed in hermaphrodites. The *dnmt3a* gene family plays a role in the establishment of DNA methylation patterns that occurs during early development, mainly during the reprogramming (Okano *et al.*, 1999; Santos & Dean 2004). In the rivulus, the two *dnmt3a* variants are differentially expressed during the early development as *dnmt3ab* is overexpressed during the last stages of development, i.e., after the DNA methylation reprogramming event, compared to *dnmt3aa* (Fellous *et al.*, 2018). As it can be seen as an erasure of epigenetic marks required for correct development of the embryo and establishment of DNA methylation patterns in the new individual, the reprogramming represents a window in which the genome is particularly sensitive to the environment (Santos & Dean, 2004). In *K. marmoratus*, this window is later and longer than in other vertebrates, especially compared to the zebrafish *Danio rerio* (Fellous *et al.*, 2018). This characteristic could be one of the reasons that allow the important phenotypic plasticity in the rivulus in a highly variable habitat as a wider window could provide sufficient time to environmental cues to be assimilated in the epigenome during embryonic development, and thus to impact genetic level expression (Fellous

*et al.*, 2018). In our model species, DNA methylation as a potential explanation for the influence of the environment could explain maintenance of variable sexing rates by modulating mate availability (Ellison *et al.*, 2015; Fellous *et al.*, 2018). To better understand the impact of environment on sex ratio in *K. marmoratus*, interactions between GSD, ESD and epigenetic mechanisms require further study.

As an emergent model for research, the origin, the evolution, and the determination of *K. marmoratus* reproduction mode are still unknown. The main objective of this study is to investigate the impact of low temperature on sex determination during *Kryptolebias marmoratus*' embryonic development. To achieve this, we compared the expression of the *de novo* DNA methyltransferases *dnmt3aa* and *dnmt3ab* at 18°C and 25°C, as well as of the sexual genes *sox9a* and *cyp19b*. The relative expression of *sox9a* and *cyp19b* were analyzed during the last stages of development, i.e., after the reprogramming period (stage 28 to 33). Comparing the relative expression of *dnmt3a* variants and sex-related genes during the embryonic development will allow to assess whether *dnmt3a* could be involved in the regulation of sexual genes by temperature, which is in turn an important mechanism allowing the existence of androdioecy in the rivulus. In a further extent, this can give an insight into the significance of epigenetic mechanisms as driver of evolution in *K. marmoratus*, as environment is known to have an impact on sex determination, and thus on sex ratio, species fitness and species evolution.

## **2. Materials and methods**

### *2.1 Ethic statement*

All rivulus husbandry were performed in accordance with the Belgian animal protection standards and were approved by the University of Namur Local Research Ethics Committee (UN 21 367 KE). The agreement number of the laboratory for fish experiments is LA1900048.

### *2.2 Experimental animals and housing*

The individuals used in this study were offsprings from adult hermaphrodites of the EPP population of *Kryptolebias marmoratus* located at the University of Namur (UNamur). The EPP stock population was initially obtained from fish sampled by Kristy Marson, Frédéric Silvestre, and Valentine Chapelle in 2019 in the Florida Keys (Emerson Point Preserve; N27°53'29.80", W82°62'55.01"). They are currently housed at 26 ± 1 C° in 12 ± 1 parts per



thousand (ppt) saltwater (Instant Ocean™ sea salt), 12:12 light:dark cycle and fed every day *ad libitum* with live *Artemia salina* nauplii. Spawning cottons were added to the tanks to provide a substrate for oviposition. These housing conditions are kept for the rest of the experiment.

### 2.3 Egg collection and temperature treatment

From April 2022 to November 2022, eggs were manually collected from adult hermaphrodite individuals. The developmental stage before incubation was determined according to the criteria described in Mourabit *et al.* (2011) (Table 1). Once identified, the eggs were individually incubated at 18°C or 25°C for 3 days. The stage of development after incubation was determined again and embryos that reached a stage between 16 and 33 were selected. Eggs were pooled by 2 or 3 depending on the stage to ensure a sufficient amount of RNA for the extraction, frozen in liquid nitrogen and conserved at -80°C.

Table 1 – Number, corresponding name and hours post-fertilization (hpf) at 25°C for stage 16 to stage 33 of *K. marmoratus* embryotic development (Mourabit *et al.*, 2011)

Stage Number	Stage Name	HPF (25°C)
16	Recognition of head and tail regions	34,5
17	Optic vesicle and somite formation	36
18	Otic vesicle formation	43,5
19	Lens formation	53
20	Heart beats	55,5
21	Body movements	58
22	Circulation	62
23	Increased vitelline circulation	71
24	Otolith formation	73
25	Pectoral fin development and erythrocyte formation	77
26	Liver formation	90
27	Increased pigmentation and body movement	105
28	Caudal fin formation	140
29	Air bladder and anal fin formation	180
30	Jaw formation	211
31	Pectoral fin movement	240
32	Hatching	310
33	Diapause	> 310

## *2.4 Sample preparation, RNA extraction and RNA quality assessment*

RNA was extracted using a TRIzol reagent. Briefly, pooled eggs were crushed using a mortar and homogenized with 150  $\mu$ l of TRI Reagent. Another 150  $\mu$ l of TRI Reagent was then added to reach a total volume of 300  $\mu$ l. After a 5 minute incubation at room temperature, 35  $\mu$ l of 1-bromo-3-chloropropane were added to separate the samples in 3 phases (aqueous phase containing the RNA, interphase consisting of DNA and organic phase containing the proteins) after incubation and centrifugation. The clear supernatant, containing the RNA, was extracted, and transferred into a new Eppendorf. 175  $\mu$ l of room temperature isopropanol were added to the supernatant to precipitate the RNA. After an incubation at room temperature and a centrifugation, the isopropanol was removed, and samples were rinsed with 350  $\mu$ l of ice-cold ethanol (75%). Samples were centrifuged, the ethanol removed, and the remaining RNA pellet dried out. After evaporation of the ethanol, pellets were resuspended in 30  $\mu$ l of nuclease-free water and frozen at  $-80^{\circ}\text{C}$ . ARN concentration and sample contamination were checked using a Nanodrop spectrophotometer.

## *2.5 Quantitative reverse transcription PCR (RT-qPCR)*

### *2.5.1. DNase and Reverse Transcription*

First, samples were treated with DNase to remove residual genomic DNA that could interfere with the technique. The RQ1 RNase-Free Dnase kit  $\text{\textcircled{R}}$  (PROMEGA) was used. Briefly, 2.5  $\mu$ l of 10x Reaction Buffer, 0.5  $\mu$ l of RNase-free DNase and nuclease-free water were added to a maximum of 5  $\mu$ g of RNA to obtain a final volume of 25  $\mu$ l. After an incubation at  $37^{\circ}\text{C}$  for 25 min, 2.5  $\mu$ l of DNase Inactivation Reagent was added to stop the activity of the DNase and thus, the degradation of residual genomic DNA. After a 2 min incubation at room temperature and a centrifugation, 11  $\mu$ l of RNA was transferred to a fresh tube to perform the reverse transcription. 11  $\mu$ l were transferred to a second tube to subsequently serve as negative control for the qPCR. The DNA-free RNA from the first tube was reverse-transcribed using the RevertAid RT Reverse Transcription kit  $\text{\textcircled{R}}$  (Thermo Fisher Scientific). Briefly, 1  $\mu$ l of Random Hexamer primer was added to each sample and incubated at  $65^{\circ}\text{C}$  for 5 min. 8  $\mu$ l of a MasterMix composed of 4  $\mu$ l of 5x Reaction Buffer, 2  $\mu$ l of 10 mM dNTP Mix, 1  $\mu$ l of RiboLock Rnase Inhibitor and 1  $\mu$ l of RevertAid RT was added to each sample to reach a final volume of 20  $\mu$ l. The samples were incubated at  $25^{\circ}\text{C}$  for 5 min,  $42^{\circ}\text{C}$  for 60 min and then,  $70^{\circ}\text{C}$  for 5 min to

catalyze the cDNA synthesis. After cooling down to 4°C, cDNA samples were conserved at -20°C.

### 2.5.2 Primer design

Specific primers have been designed to amplify the cDNA of genes of interest. The ARNm sequence of each gene was downloaded from <https://www.ncbi.nlm.nih.gov> in FASTA format (exons). Pairs of primers were designed using AmplifX®, the specificity was tested via Primer-Blast (Ye *et al.*, 2012) and Multiple Primer Analyzer (Breslauer, 1986) was used to detect potential undesirable primer-dimers. The primer efficiency was tested via a standard qPCR. To do so, cDNA from each sample is pooled together and serially diluted (5x, 25x, 125x, 625x, 3125x). Triplicates were done for each sample. Two negative controls (nuclease-free water and DNA-free RNA) were also carried out in triplicate. 2.5 µl of sample (cDNA, nuclease-free water or DNA-free RNA), 2.5 µl of primer (2.5 mM) and 5 µl of SYBER green (Bio Rad®) are added in each well. The qPCRs were performed with a QuantStudio 5 Real-Time PCR System® (Thermo Fisher Scientific), and the results were retrieved and analyzed via QuantStudio Design and Analysis® (Thermo Fisher Scientific). Only primers with an efficacy between 95% and 105% and a  $R^2 > 0,98$  were selected (see appendix 1).

### 2.5.3 qPCR amplification

After the primer assessment, the RT-qPCR was carried out. The cDNA samples were diluted 100x and each sample was analyzed in triplicate. 2.5 µl of sample (cDNA, nuclease-free water or DNA-free RNA), 2.5 µl of primer (2,5 mM) and 5 µl of SYBER green (Bio Rad®) were added in a 384-well plate, centrifuged and placed in the Real-Time qPCR system.

### 2.6 Data analysis

To calculate the relative gene expression, we used the  $\Delta\Delta C_t$  method with 18S as housekeeping gene. The best housekeeping gene was chosen by using RefFinder (Xie *et al.*, 2012) (See appendix 1). To do so, we calculated the  $\Delta C_t$  by subtracting the  $C_t$  of the housekeeping gene from the  $C_t$  of the gene of interest for the same condition. We calculated the  $\Delta\Delta C_t$  by subtracting the  $\Delta C_t$  of the reference group, corresponding to the treatment 25°C for the earliest stage

analyzed (Stage 16 or Stage 28). We calculated the fold change using  $2^{-\Delta\Delta C_t}$ . Samples with a Ct standard deviation between replicates superior to 0,5 were removed. A linear model was done on fold change values to eliminate any outliers. For each gene and temperature, the function outlierTest() (package *car*; R version 4.0.3) was used on the model  $lm(FC \sim Stage)$ . One dnmt3aa sample (p-value < 0,05) was removed (See appendix 2).

## 2.7 Statistic analysis

Normality and homogeneity hypotheses were tested with Shapiro-Wilk and Levene's test respectively (See appendix 3). Data for dnmt3aa and dnmt3ab were log2 transformed to achieve normality. A two-way ANOVA was used to test the interaction between temperatures and stages. A post-hoc Tukey's test was used to assess differences across temperature and stages. For all statistical analyses, a p-value  $p < 0,05$  was considered significant. All analyses were performed on R (version 4.0.3).

## 3. Results

### 3.1 *de novo* DNA methylation gene expression

#### 3.1.1 *dnmt3aa*

The two-way ANOVA showed that development stage and temperature had a significant influence on the gene expression of dnmt3aa. The interaction between the developmental stage and the temperature was also significant (Table 2). At 18°C, dnmt3aa expression increased from stage 16 (FC =  $0.90 \pm 0.29$ ) to reach a first peak at stage 21 (FC =  $4.97 \pm 1.77$ ). Even though the maximum expression level is obtained during stage 32 (FC =  $10.15 \pm 2.45$ ), it is not significantly different from stage 21. The expression of dnmt3aa stayed constant from stage 21 to stage 33. At 25°C, dnmt3aa expression increased from stage 16 (reference group; FC =  $1.04 \pm 0.38$ ) to reach a maximum during stage 30 (FC =  $4.36 \pm 1.45$ ), without reaching a peak at stage 21. In the first stages of the development, the increase of dnmt3aa expression was slower at 25°C compared to 18°C. Regarding the impact of temperature on gene expression, dnmt3aa was overexpressed at 18°C compared to 25°C at stage 21 only (Figure 2; Figure 3; Table 1).

### A. 18°C





Figure 2 – Fold change of dnmt3aa (solid line) and dnmt3ab (dash line) during embryo development at (A) 18°C and (B) 25°C.

Table 2 – Results of the ANOVA II and significant results of post hoc Tukey’s tests for the gene dnmt3aa (p-value < 0.05). For the Tukey’s test, the group name corresponds to stage – temperature.

ANOVA II			
	F	p-value	
Stage	8.884	4.36e-13	***
Temperature	6.954	9.80e-03	**
Stage : Temperature	2.771	9.23e-04	***

Tukey's test		
Group 1	Group 2	p-value (Tukey's correction)
16-18	21-18	3.97e-05
16-18	22-18	2.13e-03
16-18	23-18	1.83e-02
16-18	24-18	1.51e-02
16-18	25-18	1.43e-02
16-18	29-18	1.52e-02
16-18	30-18	4.44e-02
16-18	31-18	7.09e-04
16-18	32-18	7.70e-10
16-18	33-18	1.07e-03
16-25	25-25	3.65e-02
16-25	26-25	1.81e-02
16-25	27-25	3.51e-02
16-25	29-25	1.55e-02
16-25	30-25	1.60e-03
16-25	31-25	4.28e-03
16-25	32-25	2.90e-03
17-18	21-18	1.07e-02
17-18	32-18	9.31e-08
18-18	21-18	3.48e-02
18-18	32-18	4.87e-07
19-18	32-18	1.21e-02
20-18	32-18	3.07e-05
21-18	21-25	9.71e-04
21-25	30-25	1.86e-02
21-25	31-25	4.28e-02
21-25	32-25	3.09e-02
23-18	32-18	9.74e-03
25-18	32-18	1.29e-02
26-18	32-18	2.95e-04
28-18	32-18	3.29e-05
29-18	32-18	1.20e-02
30-18	32-18	3.31e-03




Figure 3 - Fold change of *dnmt3aa* during the embryo development of *K. marmoratus* at (A) 18°C and (B) 25°C. A two-way ANOVA showed that the interaction between the developmental stage and the temperature is significant ( $p$ -value =  $9.23e-04$ ). 18S was used housekeeping gene. The red box represents the stage for which temperature as a significant impact on gene expression. Significant differences between stages and temperature were checked using post hoc Tukey's test.

### 3.1.2 *dnmt3ab*

The development stage and the temperature had a significant influence on the gene expression of *dnmt3ab*, but no interaction was found between stages and the temperatures (Table 2). At 18°C, the gene expression rises from stage 16 (FC =  $1.12 \pm 0.32$ ) to hit the first peak at stage 21 (FC =  $4.43 \pm 2.2$ ). The maximum expression level is obtained at stage 32 (FC =  $4.6 \pm 1.74$ ) but the difference with stage 21 was not significant. The expression level of *dnmt3ab* stayed constant from stage 21 to stage 33. At 25°C, stage 32 (FC =  $3.5 \pm 2.12$ ) is the only one considered as significantly different from the reference group (T = 25°C; Stage = 16). As no

peak is reached at stage 21, the increase of dnmt3ab expression was slower at 25°C compared to 18°C during the first stages of development. Regarding the impact of temperature on gene expression, dnmt3ab was overexpressed at 18°C compared to 25°C at stage 21 only (Figure 4; Table 3).

Table 3 – Results of the ANOVA II and significant results of post hoc Tukey’s tests for the gene dnmt3ab (p-value < 0.05). For the Tukey’s test, the group name corresponds to stage – temperature.

ANOVA II		
	F	p-value
Stage	4.792	2.48e-07 ***
Temperature	18.715	3.60e-05 ***
Stage : Temperature	1.249	2.42e-01

Tukey's test		
Group 1	Group 2	p-value (Tukey's correction)
16-18	21-18	2.85e-04
16-18	31-18	1.30e-02
16-18	32-18	6.80e-05
16-18	33-18	3.20e-02
16-25	32-25	9.60e-03
21-18	21-25	1.08e-02






Figure 4 - Fold change of *dnmt3ab* during the embryo development of *K. marmoratus* at (A) 18°C and (B) 25°C. The interaction between the developmental stage and the temperature is not significant ( $p$ -value =  $2.42e-01$ ). 18S was used as housekeeping gene. The red box represents the stage for which temperature as a significant impact on gene expression. Significant differences between stages and temperature were checked using post hoc Tukey's test.

### 3.2 Sex-related gene expression

Regarding the sex-related genes, the stage and/or the temperature had a significant impact on gene expression (Table 4; Table 5). At 18°C, *sox9a* expression decreased from stage 28 (FC =  $2.75 \pm 0.5$ ) to reach a minimum during diapause (FC =  $0.62 \pm 0.21$ ). At 25°C, no significant differences could be highlighted. Regarding the impact of temperature on the gene expression, *sox9a* is overexpressed at 18°C during stage 28 only (Figure 5A; Table 4). For *cyp19b* expression, there were no significant differences between stages at 18°C. At 25°C, *cyp19b* expression increased from stage 28 (FC =  $1.03 \pm 0.03$ ) to reach a maximum during diapause

(FC = 2.89 ± 0.08). No significant differences in *cyp19b* expression could be highlighted between temperatures (Figure 5B; Table 5).

Table 4 – Results of the ANOVA II and significant results of post hoc Tukey’s tests for the gene *sox9a* (p-value < 0.05). For the Tukey’s test, the group name corresponds to stage – temperature.

ANOVA II		
	F	p-value
Stage	11.634	5.62e-07 ***
Temperature	0.012	9.13e-01
Stage : Temperature	7.561	4.47e-05 ***

Tukey's test		
Group 1	Group 2	p-value (Tukey's correction)
28-18	28-25	4.73e-04
28-18	32-18	4.83e-06
28-18	33-18	1.65e-06
29-18	32-18	2.18e-04
29-18	33-18	7.09e-05
30-18	32-18	2.97e-02
30-18	33-18	1.13e-02
31-18	32-18	1.21e-02
31-18	33-18	4.68e-03

Table 5 – Results of the ANOVA II and significant results of post hoc Tukey’s tests for the gene *cyp19b* (p-value < 0.05). For the Tukey’s test, the group name corresponds to stage – temperature.

ANOVA II		
	F	p-value
Stage	6.073	2.34e-04 ***
Temperature	0.517	4,76e-01
Stage : Temperature	3.128	1,68e-02 *

Tukey's test		
Group 1	Group 2	p-value (Tukey's correction)
28-25	32-25	1.76e-02
28-25	33-25	1.29e-03
29-25	33-25	8.71e-03
30-25	33-25	1.46e-02




Figure 5 – Fold change of (A) *sox9a* and (B) *cyp19b* during the last stages of *K. marmoratus* embryo development at 18°C (blue) and 25°C (red). A two-way ANOVA showed that the interaction between the developmental stage and the temperature is significant ( $p\text{-value}_{\text{sox9a}} = 4.47\text{e-}05$ ;  $p\text{-value}_{\text{cyp19b}} = 1.68\text{e-}02$ ). 18S was used as housekeeping gene. Significant differences between stages and temperature were checked using post hoc Tukey's test.

#### 4. Discussion

Previous studies on *K. marmoratus* showed that different populations have different sex ratio (Turner et al., 1990; Taylor et al., 2001) depending on the population's genetics and the environment (Harrington, 1967; Harrington & Kallman, 1968; Taylor, 2012; Ellison *et al.*, 2015; Kelley *et al.*, 2016). Harrington (1967; 1968) discovered that temperature plays an important role in the sex determination of the rivulus and situated the thermolabile period between the stage 31 and 32. When incubated at 18°-20°, i.e., 5 to 7 degrees below their optimum of 25°C, *K. marmoratus* embryos have tendency to differentiate into males. Ellison *et al.* (2015) also showed that masculinization is induced by low temperature, as well as a modification of the level of expression of the sexual genes *cyp19a*, associated with the ovarian development, and *sox9a*, associated with the development of testicular tissues, and that DNA methylation could be the intermediary through which temperature modulates sexual identity. DNA methylation is catalyzed by the activity of enzymes from the dnmts family, through which methyl-groups are added on DNA strands, and reduces or even suppresses gene expression by

blocking the binding of transcription factors (Moore & Fan, 2013). Fellous et al.. (2018) studied the expression level of different dnmts during the embryonic development of hermaphroditic embryos, and between male and hermaphroditic adult tissues. For embryos incubated at 25°C, they found that dnmt3ab was overexpressed during the last stages of embryonic development compared to previous stages and to dnmt3aa. According to them, that could indicate a potential role of rivulus dnmt3ab in oogenesis and in late DNA remethylation during embryonic development. They also found that dnmt3ab was overexpressed in male tissues and dnmt3aa was overexpressed in hermaphroditic tissues. In addition, Ellison *et al.*. (2015) observed that a significant interaction between sex, temperature and methylation patterns. They were also able to identify several genes differentially methylated in males and hermaphrodites, namely *sox9a* and *cyp19a*. By highlighting differences in dnmt3a expression between males and hermaphrodites (Fellous *et al.*, 2018) and gene candidates for the temperature-mediated sex regulation (Ellison *et al.*, 2015), these studies incited us to look at the impact of temperature on sex determination during the embryonic development of *K. marmoratus* to try to identify a possible correlation between the expression of sexual genes and of the enzymes directly responsible for *de novo* methylation. In the current study, we therefore investigated the effect of a low temperature (18°C) on embryonic SD by looking at the expression of *de novo* DNA methylation genes (dnmt3aa and dnmt3ab) and sex-related genes (*sox9a* and *cyp19b*). Based on the work of Harrington (1967; 1968), we incubated eggs at 18°C and 25°C.

First, we investigate the expression pattern of dnmt3aa and dnmt3ab in embryos incubated at 18°C and 25°C. In most of teleost fish, two dnmt3a paralogs exist, dnmt3aa and dnmt3ab, both responsible for *de novo* DNA methylation (Campos *et al.*, 2012; Liu *et al.*, 2020). In this study, we found that the two variants were similarly expressed during embryo development of *K. marmoratus* when incubated at the same temperature. This result differs from what Fellous *et al.*. (2018) found for *K. marmoratus* embryos incubated at 25°C, which is a differential expression between dnmt3aa and dnmt3ab during the last stages of embryonic development. This differential expression could indicate a potential role of rivulus dnmt3ab in oogenesis and in late DNA remethylation during embryonic development. It is important to note that the fish population that we use (EPP) differs from the one studied by Fellous et al.. (DC4). As sex determination of *K. marmoratus* depends on environmental but also on genetic factors (Ellison et al., 2015), further studies would be needed to assess the part of genetic and/or environmental factors that influence the difference in the expression level of dnmt3aa in EPP and DC4 population. The similitude of expression levels of the two variants does not correspond to the differences found by Fellous et al.. (2018) in adults tissues. This may be due to the fact

that Fellous et al.. (2018) did the analyses on secondary males while we looked at potential primary males. Sex determination mechanisms and sex differentiation mechanisms might differ in *K. marmoratus*. Even though, understanding these differences needs further investigation, it has already been shown that steroid hormones are key regulators for sex differentiation (Godwin, 2010; Garcia *et al.*, 2016). Regarding the expression of dnmt3a variants according to the development temperature, we found that the expression level of both variants increased with the development stage, but the augmentation seems to start earlier at 18°C (male-producing temperature) compared to 25°C (hermaphrodite-producing temperature), with a maximal expression of dnmt3a at stage 21 in potential future primary males. The early expression of the dnmt3s in male embryos might increase the methylation and thus downregulate genes responsible for ovarian development. This corresponds to what was found in the zebrafish where the inhibition of dnmts by exposition to 5-aza-2'-deoxycytidine during embryonic development induces the feminization of the fish (Ribas *et al.*, 2017). In the Nile tilapia, dnmts might also have a potential role in the epigenetic control of gonadal development (Li *et al.*, 2014). If present in both testis and ovary, dnmt3aa and dnmt3ab will a higher level of expression in testis than in ovary. The difference of expression between gonads can be compared with the dnmt3 profile expression of *K. marmoratus* where dnmt3a expression increases earlier in potential males compared to hermaphrodites. In the Nile tilapia, the use of a dnmt inhibitor reduced dnmt3aa and dnmt3ab expression level leading to a significant augmentation in expression of cyp19a1a and dmrt1, key genes of gonadal development for females and males respectively. If embryos incubated at 18°C give a majority of males, DNA methyltransferase, and particularly dnmt3a, might have a potential role in gonadal development of *K. marmoratus*.

Then we looked at the level expression of sexual genes that might play a role in sex determination, namely sox9a and cyp19b. Sox9a is part of the SRY-related high mobility group box (SOX) genes family which corresponds to transcription factors that play crucial roles during reproduction and development (Hu *et al.*, 2021). While sox9 has been demonstrated to play a key role in sex determination and differentiation in vertebrates, it is not yet proven in teleost fish and its function is still debated (Chiang *et al.*, 2001; Nakamura *et al.*, 2012). In our study, embryos incubated at 18°C underwent a diminution of sox9a expression level during the last stages of development as well as an overexpression of sox9a at stage 28 compared to embryos incubated at 25°C, i.e. sox9a is overexpressed in males compared to hermaphrodites. In hermaphrodites, a constant expression level of sox9a was kept during the last stages of their development. Harrington (1968) showed that rivulus thermolabile stage was the second part of stage 31. Here, we found that a sexual gene involved in testis development is overexpressed

prior to stage 31. As for *dnmt3a*, the fish population analyzed here is different from the ones used by Harrington. The genetic differences between the two population might translate into an earlier thermolabile period in EPP population. Ellison et al.. (2015) found that *sox9a* was hypermethylated in males incubated at low temperature, which lead to a reduction of *sox9a* expression level in males compared to hermaphrodites, as opposed to what we observed in our analyses. Several scenarios could explain this. Again, the genetic differences of the EPP population compared to the ones used by Ellison *et al.*, leads to differential expression of *sox9a*. If embryos incubated at 18°C do not yield 100% of males, the expression level observed might be superior to the expected one. Also, we see that, even though it is not significant compared to 25°C, expression level of *sox9a* drop significantly from stage 28 to stage 33 at 18°C. It could also be interesting to look at the second paralogs of *sox9*, *sox9b*. In the Nile tilapia, the two paralogs do not have the same role in sex determination. While *sox9a* may play major role in the regulation of ovarian differentiation, *sox9b* seems to be involved in the regulation of testicular differentiation (Wei et al., 2016).

The *cyp19* gene encodes cytochrome P450 aromatase, which catalyzes the synthesis of estrogens (Chiang *et al.*, 2001). Two variants have been highlighted in fish, *cyp19a* which is expressed in the ovary, and *cyp19b* which is expressed in the brain (Chiang *et al.*, 2001). As the regulation of this gene dictates the ratio of androgens to estrogens, an appropriate expression of this enzyme is critical for reproduction and sex differentiation for most vertebrates (Trant *et al.*, 2001). Here, we found that expression level of *cyp19b* increases during the last stages of development for embryos incubated at 25°C, i.e. the hermaphrodites. In embryos incubated at male-producing temperature, expression level of *cyp19b* stays constant. Ellison *et al.* (2015) showed that *cyp19a* could be involved in modulating sex ratio of the rivulus in response to temperature during the embryonic development as it is downregulated in the brain of fish incubated at 25°C. Here, we observed an augmentation of *cyp19b* expression at hermaphrodite producing temperature, which could indicate that this gene is indeed involved in the pathway leading to the differentiation of ovarian tissue in the rivulus, as it is in other species such as the zebrafish and the sea bass *Dicentrarchus labrax* (Trant *et al.*, 2001; Blázquez & Piferrer, 2005), but opposite to what Ellison et al.. (2015) suggest.. *Cyp19b* is however not always involved in sex determination or differentiation in all species, as for example for the Atlantic halibut *Hippoglossus hippoglossus*, where no dimorphic expression has been observed (van Nes *et al.*, 2005). In the Nile tilapia, another form of *cyp19b*, named tilapia *cyp19b* (*tCyp19b*) was shown to be exclusively expressed in female brain (Chang *et al.*, 2005). Further study still has to be done to determine the implication of *cyp19b* in sex determination by temperature.

## **Conclusion**

Through this study, we manage to highlight the difference in level expression of *dnmt3a* between embryos incubated at 18°C and at 25°C. We also showed that *sox9a* is more expressed in male embryos. No significant differences could be observed for *cyp19b*. Regarding our main objective, no significant differences could be identified in *dnmt3aa* and *dnmt3ab* expression level, neither for embryos incubated at 18°C nor for those incubated at 25°C, during the last stages of embryonic development. No correlation could be found between *dnmt3aa/ab* and sexual genes *sox9a/cyp19b* in our study. In further study, it will be interesting to look at other sexual genes that might be involved in the sex determination by temperature like *dmrt1*, *amh*, *foxl2* or *fig-alpha*, and to further investigate the methylation of those genes.

## **Acknowledgments**

I would like to thank my supervisor Yves Blanco-Alvarez for all the advice, support and help during my thesis. I also want to thank Dr. Frédéric Silvestre and the rest of the LEAP team for the assistance and feed-back during my time in the lab.

## 7. Bibliography

- Avise, J. C., & Tatarenkov, A. (2015). Population genetics and evolution of the mangrove rivulus *Kryptolebias marmoratus*, the world's only self-fertilizing hermaphroditic vertebrate. *Journal of Fish Biology*, 87(3), 519-538.
- Bannister, A. J., & Kouzarides, T. (2011). Regulation of chromatin by histone modifications. *Cell research*, 21(3), 381-395.
- Baroiller, J.F., (1995). Temperature and sex chromosomes govern sex ratios of the mouthbrooding cichlid fish *Oreochromis niloticus*. *J. Exp. Zool.* 273, 216–223.
- Baroiller, J.F., D’Cotta, H., Bezault, E., Wessels, S., Hoerstgen-Schwark, G., (2009a). Tilapia sex determination: where temperature and genetics meet. *Comp. Biochem. Physiol. - A Mol. Integr. Physiol.* 153, 30–38.
- Baroiller, J. F., D’Cotta, H., & Saillant, E. (2009b). Environmental effects on fish sex determination and differentiation. *Sexual development*, 3(2-3), 118-135.
- Bachtrog, D., Mank, J. E., Peichel, C. L., Kirkpatrick, M., Otto, S. P., Ashman, T. L., ... & Tree of Sex Consortium. (2014). Sex determination: why so many ways of doing it?. *PLoS biology*, 12(7), e1001899.
- Berger, S. L., Kouzarides, T., Shiekhattar, R., & Shilatifard, A. (2009). An operational definition of epigenetics. *Genes & development*, 23(7), 781-783.
- Bertho, S., Herpin, A., Branthonne, A., Jouanno, E., Yano, A., Nicol, B., ... & Guiguen, Y. (2018). The unusual rainbow trout sex determination gene hijacked the canonical vertebrate gonadal differentiation pathway. *Proceedings of the National Academy of Sciences*, 115(50), 12781-12786.
- Blázquez, M., & Piferrer, F. (2005). Sea bass (*Dicentrarchus labrax*) androgen receptor: cDNA cloning, tissue-specific expression, and mRNA levels during early development and sex differentiation. *Molecular and cellular endocrinology*, 237(1-2), 37-48.
- Bollati, V., & Baccarelli, A. (2010). Environmental epigenetics. *Heredity*, 105(1), 105-112.
- Breslauer, K. J., Frank, R., Blöcker, H., & Marky, L. A. (1986). Predicting DNA duplex stability from the base sequence. *Proceedings of the National Academy of Sciences*, 83(11), 3746-3750.
- Campos, C., Valente, L. M., & Fernandes, J. M. (2012). Molecular evolution of zebrafish *dnmt3* genes and thermal plasticity of their expression during embryonic development. *Gene*, 500(1), 93-100.
- Capel, B. (2017). Vertebrate sex determination: evolutionary plasticity of a fundamental switch. *Nature Reviews Genetics*, 18(11), 675–689.
- Charlesworth, D. (1984). Androdioecy and the evolution of dioecy. *Biological Journal of the Linnean Society*, 22(4), 333-348
- Chiang, E. F. L., Pai, C. I., Wyatt, M., Yan, Y. L., Postlethwait, J., & Chung, B. C. (2001). Two *sox9* genes on duplicated zebrafish chromosomes: expression of similar transcription activators in distinct sites. *Developmental biology*, 231(1), 149-163.
- Costa, W. J. E. M. (2016). Colouration, taxonomy and geographical distribution of mangrove killifishes, the *Kryptolebias marmoratus* species group, in southern Atlantic coastal plains of Brazil (Cyprinodontiformes: Rivulidae). *Ichthyological Exploration of Freshwaters*, 27(2), 183-192.
- Ellison, A., Jones, J., Inchley, C., & Consuegra, S. (2013). Choosy males could help explain androdioecy in a selfing fish. *The American Naturalist*, 181(6), 855-862.



- Ellison, A., Rodriguez Lopez, C. M., Moran, P., Breen, J., Swain, M., Megias, M., ... & Consuegra, S. (2015). Epigenetic regulation of sex ratios may explain natural variation in self-fertilization rates. *Proceedings of the Royal Society B: Biological Sciences*, 282(1819), 20151900.
- Fellous, A., Labed-Veydert, T., Locrel, M., Voisin, A. S., Earley, R. L., & Silvestre, F. (2018). DNA methylation in adults and during development of the self-fertilizing mangrove rivulus, *Kryptolebias marmoratus*. *Ecology and evolution*, 8(12), 6016-6033.
- Fischer EA, Petersen CW (1986) Contrasts in the behavior of males and hermaphrodites in the hermaphroditic coral reef fish, *Serranus baldwini*. *Ethology* 73:235–246
- Garcia, M. J., Ferro, J. M., Mattox, T., Kopelic, S., Marson, K., Jones, R., ... & Earley, R. L. (2016). Phenotypic differences between the sexes in the sexually plastic mangrove rivulus fish (*Kryptolebias marmoratus*). *Journal of Experimental Biology*, 219(7), 988-997.
- Ge, C., Ye, J., Zhang, H., Zhang, Y., Sun, W., Sang, Y., ... & Qian, G. (2017). Dmrt1 induces the male pathway in a turtle species with temperature-dependent sex determination. *Development*, 144(12), 2222–2233.
- Ge, C., Ye, J., Weber, C., Sun, W., Zhang, H., Zhou, Y., ... & Capel, B. (2018). The histone demethylase KDM6B regulates temperature-dependent sex determination in a turtle species. *Science*, 360(6389), 645-648.
- Godwin, J. (2010). Neuroendocrinology of sexual plasticity in teleost fishes. *Frontiers in neuroendocrinology*, 31(2), 203-216.
- Haag, E. S., & Doty, A. V. (2005). Sex determination across evolution: connecting the dots. *PLoS Biol*, 3(1), e21
- Harrington Jr, R. W. (1961). Oviparous hermaphroditic fish with internal self-fertilization. *Science*, 134(3492), 1749-1750.
- Harrington Jr, R. W. (1967). Environmentally controlled induction of primary male gonochorists from eggs of the self-fertilizing hermaphroditic fish, *Rivulus marmoratus* Poey. *The Biological Bulletin*, 132(2), 174-199.
- Harrington Jr, R. W., & Kallman, K. D. (1968). The homozygosity of clones of the self-fertilizing hermaphroditic fish *Rivulus marmoratus* Poey (Cyprinodontidae, Atheriniformes). *The American Naturalist*, 102(926), 337-343.
- Harrington Jr, R. W. (1968). Delimitation of the thermolabile phenocritical period of sex determination and differentiation in the ontogeny of the normally hermaphroditic fish *Rivulus marmoratus* Poey. *Physiological Zoology*, 41(4), 447-460.
- Hayes, T.B., (1998). Sex determination and primary sex differentiation in amphibians: genetic and developmental mechanisms. *J. Exp. Zool.* 281, 373–399.
- Hu, Y., Wang, B., & Du, H. (2021). A review on sox genes in fish. *Reviews in Aquaculture*, 13(4), 1986-2003.
- Kanamori, A., Sugita, Y., Yuasa, Y., Suzuki, T., Kawamura, K., Uno, Y., ... & Sakakura, Y. (2016). A genetic map for the only self-fertilizing vertebrate. G3: Genes, Genomes, *Genetics*, 6(4), 1095-1106.
- Kelley, J. L., Yee, M. C., Brown, A. P., Richardson, R. R., Tatarenkov, A., Lee, C. C., ... & Earley, R. L. (2016). The genome of the self-fertilizing mangrove rivulus fish, *Kryptolebias marmoratus*: a model for studying phenotypic plasticity and adaptations to extreme environments. *Genome biology and evolution*, 8(7), 2145-2154.
- Kim, J. K., Samaranayake, M., & Pradhan, S. (2009). Epigenetic mechanisms in mammals. *Cellular and molecular life sciences*, 66(4), 596-612.
- Korpelainen, H. (1990). Sex ratios and conditions required for environmental sex determination in animals. *Biological reviews*, 65(2), 147-184.
- Kwon, J. Y., McAndrew, B. J., & Penman, D. J. (2001). Cloning of brain aromatase gene and expression of brain and ovarian aromatase genes during sexual differentiation in genetic male and female Nile tilapia *Oreochromis niloticus*. *Molecular Reproduction and Development: Incorporating Gamete Research*, 59(4), 359-370.

- Lee, J. S., Raisuddin, S., & Schlenk, D. (2008). *Kryptolebias marmoratus* (Poey, 1880): a potential model species for molecular carcinogenesis and ecotoxicogenomics. *Journal of Fish Biology*, 72(8), 1871-1889.
- Li, C. G., Wang, H., Chen, H. J., Zhao, Y., Fu, P. S., & Ji, X. S. (2014). Differential expression analysis of genes involved in high-temperature induced sex differentiation in Nile tilapia. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 177, 36-45.
- Li, X. Y., & Gui, J. F. (2018). Diverse and variable sex determination mechanisms in vertebrates. *Science China Life Sciences*, 61(12), 1503-1514.
- Leonard, J. L. (2018). The evolution of sexual systems in animals. In *Transitions between sexual systems* (pp. 1-58). Springer, Cham.
- Liu, J., Hu, H., Panserat, S., & Marandel, L. (2020). Evolutionary history of DNA methylation related genes in chordates: new insights from multiple whole genome duplications. *Scientific reports*, 10(1), 1-14.
- Loughland, I., Little, A., & Seebacher, F. (2021). DNA methyltransferase 3a mediates developmental thermal plasticity. *BMC biology*, 19(1), 1-11.
- Matsuda, M., Nagahama, Y., Shinomiya, A., Sato, T., Matsuda, C., Kobayashi, T., ... & Sakaizumi, M. (2002). DMY is a Y-specific DM-domain gene required for male development in the medaka fish. *Nature*, 417(6888), 559-563.
- Moore, L. D., Le, T., & Fan, G. (2013). DNA methylation and its basic function. *Neuropsychopharmacology*, 38(1)
- Moreira-Pinto, B., and Pires-daSilva, A. (2018). Comparative Reproductive Biology in Nematodes. In *Encyclopedia of Reproduction* (Second Edition), M.K. Skinner, ed. Academic Press, pp. 508-512.
- Mourabit, S., Edenbrow, M., Croft, D. P., & Kudoh, T. (2011). Embryonic development of the self-fertilizing mangrove killifish *Kryptolebias marmoratus*. *Developmental Dynamics*, 240(7), 1694-1704.
- Nagahama, Y. (2005). Molecular mechanisms of sex determination and gonadal sex differentiation in fish. *Fish physiology and biochemistry*, 31(2), 105-109.
- Nakamura, S., Watakabe, I., Nishimura, T., Toyoda, A., Taniguchi, Y., & Tanaka, M. (2012). Analysis of medaka sox9 orthologue reveals a conserved role in germ cell maintenance. *Plos one*, 7(1), e29982.
- Navarro-Martín, L., Viñas, J., Ribas, L., Díaz, N., Gutiérrez, A., Di Croce, L., & Piferrer, F. (2011). DNA methylation of the gonadal aromatase (cyp19a) promoter is involved in temperature-dependent sex ratio shifts in the European sea bass. *PLoS genetics*, 7(12), e1002447.
- Okano, M., Bell, D. W., Haber, D. A., & Li, E. (1999). DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell*, 99(3), 247-257.
- Ospina-Alvarez, N., & Piferrer, F. (2008). Temperature-dependent sex determination in fish revisited: prevalence, a single sex ratio response pattern, and possible effects of climate change. *PloS one*, 3(7), e2837.
- Pannell, J. R. (2002). The evolution and maintenance of androdioecy. *Annual Review of Ecology and Systematics*, 397-425.
- Penman, D. J., & Piferrer, F. (2008). Fish gonadogenesis. Part I: genetic and environmental mechanisms of sex determination. *Reviews in Fisheries Science*, 16(sup1), 16-34.
- Petersen CW (2006) Sexual selection and reproductive success in hermaphroditic seabasses. *Integr Comp Biol* 46(4):439-448

- Piferrer, F., Ribas, L., Díaz, N., 2012. Genomic approaches to study genetic and environmental influences on fish sex determination and differentiation. *Mar. Biotechnol.* 14, 591–604.
- Pla, S., Benvenuto, C., Capellini, I., & Piferrer, F. (2022). Switches, stability and reversals in the evolutionary history of sexual systems in fish. *Nature Communications*, 13(1), 1-13.
- Ribas, L., Vanezis, K., Imués, M. A., & Piferrer, F. (2017). Treatment with a DNA methyltransferase inhibitor feminizes zebrafish and induces long-term expression changes in the gonads. *Epigenetics & chromatin*, 10(1), 1-16.
- Sandra, G. E., & Norma, M. M. (2010). Sexual determination and differentiation in teleost fish. *Reviews in Fish Biology and Fisheries*, 20(1), 101-121.
- Santi, S., Gennotte, V., Toguyeni, A., Mélard, C., Antoine, N., & Rougeot, C. (2016). Thermosensitivity of the sex differentiation process in the African catfish, *Clarias gariepinus*: Determination of the thermosensitive period. *Aquaculture*, 455, 73-80
- Santos, F., & Dean, W. (2004). Epigenetic reprogramming during early development in mammals. *Reproduction*, 127(6), 643-651.
- Smith, C. A., Roeszler, K. N., Ohnesorg, T., Cummins, D. M., Farlie, P. G., Doran, T. J., & Sinclair, A. H. (2009). The avian Z-linked gene DMRT1 is required for male sex determination in the chicken. *Nature*, 461(7261), 267-271.
- Speijer, D., Lukeš, J., & Eliáš, M. (2015). Sex is a ubiquitous, ancient, and inherent attribute of eukaryotic life. *Proceedings of the National Academy of Sciences*, 112(29), 8827-8834.
- Subramoniam, T. (2017) Sexual Systems. In *Sexual Biology and Reproduction in Crustaceans* (First Edition). 1st ed. Academic Press. pp. 57-103.
- Taylor, D. S. (2000). Biology and ecology of *Rivulus marmoratus*: new insights and a review. *Florida Scientist*, 242-255.
- Tatarenkov, A., Earley, R. L., Perlman, B. M., Scott Taylor, D., Turner, B. J., & Avise, J. C. (2015). Genetic subdivision and variation in selfing rates among Central American populations of the mangrove rivulus, *Kryptolebias marmoratus*. *Journal of Heredity*, 106(3), 276-284.
- Taylor, D. S., Fisher, M. T., & Turner, B. J. (2001). Homozygosity and heterozygosity in three populations of *Rivulus marmoratus*. *Environmental Biology of Fishes*, 61(4), 455-459.
- Taylor, D. S., Turner, B. J., Davis, W. P., & Chapman, B. B. (2008). A novel terrestrial fish habitat inside emergent logs. *The American Naturalist*, 171(2), 263-266.
- Taylor, D. S. (2012). Twenty-four years in the mud: what have we learned about the natural history and ecology of the mangrove rivulus, *Kryptolebias marmoratus*?. *Integrative and Comparative Biology*, 52(6), 724-736.
- Tong, S. K., & Chung, B. C. (2003). Analysis of zebrafish cyp19 promoters. *The Journal of steroid biochemistry and molecular biology*, 86(3-5), 381-386.
- Trant, J. M., Gavasso, S., Ackers, J., Chung, B. C., & Place, A. R. (2001). Developmental expression of cytochrome P450 aromatase genes (CYP19a and CYP19b) in zebrafish fry (*Danio rerio*). *Journal of Experimental Zoology*, 290(5), 475-483.
- Turner, B. J., Elder Jr, J. F., Laughlin, T. F., & Davis, W. P. (1990). Genetic variation in clonal vertebrates detected by simple-sequence DNA fingerprinting. *Proceedings of the National Academy of Sciences*, 87(15), 5653-5657.
- Turner, B. J., Fisher, M. T., Taylor, D. S., Davis, W. P., & Jarrett, B. L. (2006). Evolution of 'maleness' and outcrossing in a population of the self-fertilizing killifish, *Kryptolebias marmoratus*. *Evolutionary Ecology Research*, 8(8), 1475-1486.

- van Nes, S., & Andersen, Ø. (2006). Temperature effects on sex determination and ontogenetic gene expression of the aromatases *cyp19a* and *cyp19b*, and the estrogen receptors *esr1* and *esr2* in Atlantic halibut (*Hippoglossus hippoglossus*). *Molecular Reproduction and Development*, 73(12), 1481-1490
- Wang, Y. Y., Sun, L. X., Zhu, J. J., Zhao, Y., Wang, H., Liu, H. J., & Ji, X. S. (2017). Epigenetic control of *cyp19a1a* expression is critical for high temperature induced Nile tilapia masculinization. *Journal of thermal biology*, 69, 76-84.
- Wang, X., Lai, F., Xiong, J., Zhu, W., Yuan, B., Cheng, H., & Zhou, R. (2020). DNA methylation modification is associated with gonadal differentiation in *Monopterus albus*. *Cell & bioscience*, 10(1), 1-11.
- Wang, F., Qin, Z., Li, Z., Yang, S., Gao, T., Sun, L., & Wang, D. (2021). *Dnmt3aa* but Not *Dnmt3ab* Is Required for Maintenance of Gametogenesis in Nile Tilapia (*Oreochromis niloticus*). *International journal of molecular sciences*, 22(18), 10170.
- Weeks, S. C., Benvenuto, C., & Reed, S. K. (2006). When males and hermaphrodites coexist: a review of androdioecy in animals. *Integrative and comparative biology*, 46(4), 449-464.
- Weeks, S. C. (2012). The role of androdioecy and gynodioecy in mediating evolutionary transitions between dioecy and hermaphroditism in the Animalia. *Evolution: International Journal of Organic Evolution*, 66(12), 3670-3686.
- Wei, L., Li, X., Li, M., Tang, Y., Wei, J., & Wang, D. (2019). *Dmrt1* directly regulates the transcription of the testis-biased *Sox9b* gene in Nile tilapia (*Oreochromis niloticus*). *Gene*, 687, 109-115.
- Xie, F., Xiao, P., Chen, D., Xu, L., and Zhang, B. (2012). miRDeepFinder: a miRNA Analysis Tool for Deep Sequencing of Plant Small RNAs. *Plant Mol. Biol.* 80, 75–84.
- Yaman R, Grandjean V. Timing of entry of meiosis depends on a mark generated by DNA methyltransferase 3a in testis. *Mol Reprod Dev.* 2010; 73(3):390-397.
- Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., & Madden, T. L. (2012). Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC bioinformatics*, 13(1), 1-11.
- Yoshimoto, S., & Ito, M. (2011). A ZZ/ZW-type sex determination in *Xenopus laevis*. *The FEBS journal*, 278(7), 1020-1026.
- Yusa, Y., Takemura, M., Sawada, K., & Yamaguchi, S. (2013). Diverse, continuous, and plastic sexual systems in barnacles. *Integrative and comparative biology*, 53(4), 701-712.

## Appendexes

### Appendix 1 – Primer efficiency

Primers for 4 genes of interest (dnmt3aa, dnmt3ab, cyp19b and sox9a) and 3 housekeeping genes (18S,  $\beta$ -actin and rpl8) presented an efficiency between 95% and 105% and  $R^2 > 0,98$  (Table S1). Based on RefFinder, the housekeeping gene 18S was kept (Table S2).  $\beta$ -actin could not be used because the number of replicates for certain stages was too low.

Table S1 – List of selected primers

Gene	Gene ID	Primers	Efficacity (%)	Standard Deviation	$R^2$
18S	108251219	F: GAACTCACCGACACCAGCA R: ATCATCGACGCTCCTGGA	100,911	0,062	0,996
$\beta$ -actin	108236185	F : CTTGCGGAATCCACGAGACC R : CCAGGGCTGTGATCTCCTTCTG	102,011	0,064	0,095
rpl8	108232351	F : TCGCTGGAGGCGGTTCGTATTGATAAGC R : CCACCGAAGGGATGCTCAACAGGATTCA	103,665	0,091	0,99
dnmt3aa	108233456	F: CGACGCTGAAGCTGTGGAAATCCAGACTCG R: CGCCTTCTCTGCAGAGTTCTTGCGTGGT	101,56	0,056	0,996
dnmt3ab	108239073	F: TCTTTATGTGGTGTCTGGTTCAAC R: TGTGCA TCAAACCTCACTTTGGA	100,149	0,095	0,99
sox9a	108241363	F: AGCATGGGCGAAGTGCACTC R: CAGCTCAAAGTCGATATTGAGCTGC	102,359	0,083	0,992
cyp19b	108230104	F: TGTTCTCAGGGTGCCTCTGAATGAG R: GAGCCATCCGAGCTTGAAGAAAACGT	102,914	0,106	0,987

Table S2 – RefFinder results for best housekeeping gene

	Ranking order		
	1	2	3
$\Delta$ Ct	18S	$\beta$ -actin	rpl8
BestKeeper	18S	$\beta$ -actin	rpl8
Normfinder	18S	$\beta$ -actin	rpl8
Genorm	$\beta$ -actin  18S		rpl8
<b>Recommended comprehensive ranking</b>	<b>18S</b>	<b><math>\beta</math>-actin</b>	<b>rpl8</b>

## Appendix 2 – Outlier elimination

To eliminate any potential outliers, a linear model  $lm(FC \sim Stage * Temperature)$  was done on fold change values for each gene. The  $p$ -values given by the function `outlierTest()` are presented in Table S3. Based on Cook's distance plot, the sample 25.28.19 was found to be an outlier in `dnmt3aa` data and removed. (Figure S1)

Table S3 – Results of the outlier test on the linear model  $lm(FC \sim Stage * Temperature)$  for each gene of interest.

Gene	Sample removed	unadjust p-value	Bonferroni p-value
<code>dnmt3aa</code>	None	0.00035914	0.046688
<code>dnmt3aa</code>	25.28.19	0.0013426	0.1732
<code>dnmt3ab</code>	None	0.00036868	0.050141
<code>sox9a</code>	None	0.0050525	0.26273
<code>cyp19b</code>	None	0.012533	0.70187




Figure S1 – Cook's distance of each sample from `dnmt3aa` dataset. Sample n°115, corresponding to sample ID 25.28.19, was considered an outlier, and removed.

### Appendix 3 – Residuals normality and Homogeneity

To be able to perform a two-way ANOVA, residuals normality and homogeneity hypotheses have been tested for each gene with Shapiro and Levene's tests respectively (Table S4). A qqplot was carried out for each gene to visualize the residuals distribution (Figure S2). Based on the graphs and the p-values, data for dnmt3aa and dnmt3ab were log2 transformed.

Table S4 – Results of the Shapiro and Levene's tests for each gene of interest.

Gene	Data transformation	Test	F	p-value
dnmt3aa	None	Shapiro	0.990	0.437
		Levene	2.79	0.0000467
	Log2	Shapiro	0.982	0.0873
		Levene	1.22	0.227
dnmt3ab	None	Shapiro	0.959	0.000389
		Levene	2.43	0.000306
	Log2	Shapiro	0.985	0.154
		Levene	0.920	0.599
sox9a	None	Shapiro	0.982	0.598
		Levene	0.478	0.906
cyp19b	None	Shapiro	0.986	0.744
		Levene	0.951	0.503




Figure S2 – qqplot of (A) *dnmt3aa*, (B) *dnmt3aa* (log<sub>2</sub> transformed), (C) *dnmt3ab*, (D) *dnmt3ab* (log<sub>2</sub> transformed), (E) *sox9a*, (F) *cyp19b*