

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Reduced matrix elements in semi-classical line shape calculations

Vispoel, Bastien; Gamache, Robert

Published in:

Journal of Physics : Conference Series

DOI:

[10.1088/1742-6596/1289/1/012023](https://doi.org/10.1088/1742-6596/1289/1/012023)

Publication date:

2019

Document Version

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (HARVARD):

Vispoel, B & Gamache, R 2019, 'Reduced matrix elements in semi-classical line shape calculations: Application to H₂O-H₂', *Journal of Physics : Conference Series*, vol. 1289, no. 1, 012023. <https://doi.org/10.1088/1742-6596/1289/1/012023>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

PAPER • OPEN ACCESS

Reduced matrix elements in semi-classical line shape calculations: Application to $\text{H}_2\text{O}-\text{H}_2$

To cite this article: Bastien Vispoel and Robert R. Gamache 2019 *J. Phys.: Conf. Ser.* **1289** 012023

View the [article online](#) for updates and enhancements.

You may also like

- [Application of \$\text{Li}_{1-x}\text{Al}_x\text{Ge}_{2-x}\(\text{PO}_4\)_3\$ Glass Electrolyte to Air Electrode for All-Solid-State \$\text{Li}-\text{O}_2\$ Battery](#)

Hirokazu Kitaura and Haoshen Zhou

- [Development of the kinetic molecular theory of liquids concept inventory: preliminary results on university students' misconceptions](#)

Nataša Erceg, Ivica Aviani, Klara Grlaš et al.

- [Influence of vibrations and rotations of diatomic molecules on their physical properties: II. Refractive index, reactivity and diffusion coefficients](#)

Alexander S Sharipov, Boris I Loukhovitski and Alexander M Starik



The Electrochemical Society

Advancing solid state & electrochemical science & technology

243rd ECS Meeting with SOFC-XVIII

More than 50 symposia are available!

Present your research and accelerate science

Boston, MA • May 28 – June 2, 2023

[Learn more and submit!](#)

Reduced matrix elements in semi-classical line shape calculations: Application to H₂O-H₂

Bastien Vispoel, Robert R. Gamache

Department of Environmental, Earth, and Atmospheric Sciences, University of Massachusetts Lowell, MA 01854, USA

E-mail address: robert_gamache@uml.edu

Abstract. Making reliable semi-classical calculations using the Modified Complex Robert and Bonamy formalism require an intermolecular potential expanded to a high order. Reduced matrix elements (RMEs) need to be computed for each component of the intermolecular potential using the wavefunctions of the states involved in the transition. Calculations for a number of transitions of 13 vibrational bands were performed using the vibrational dependent RMEs and using the ground state RMEs for both upper and lower states. The calculations show that for transitions that obey the sum rules the spectroscopic effect is small.

1. Introduction

It is now possible to make semi-classical calculations for the collisional broadened half-width and line shift that do not utilize scaling factors or arbitrary adjustable parameters. The theory was developed by Robert and Bonamy [1] and later modified by Ma *et al.* [2] and is called the Modified Complex Robert and Bonamy formalism (MCRB). Reliable calculations can be made if the calculations employ the best-known intermolecular potentials, with the atom-atom component expanded to high order as recommended by Ma *et al.* [3], include the imaginary components of the collisional cross-sections, and the velocity integral is explicitly computed; all of these factors have been shown to be necessary to yield accurate results [4-8]. It was shown that the potential should be expanded up to 20th order and rank = 4 to ensure the convergence of the calculations [3, 9]. The resulting intermolecular potential can contain hundreds to thousands of terms.

To make a complete calculation using the MCRB formalism, one must compute the reduced matrix elements (RMEs) that correspond to each term of the intermolecular potential. The RMEs squared give the probability of a collisionally induced transition between a lower and an upper state. When calculations are performed for vibrational bands, the vibrationally dependent reduced matrix elements must be used.

In the calculations, all collisional connections $i' \leftarrow i$ must be included. A convenient check that all connected states have been included can be accomplished using the sum rules developed by Lamouroux *et al.* [10]. These sums rules states that, for a given state i , the sum of the reduced matrix elements squared for all possible connected states i' equals 1. The sums rules are applied for each term in the intermolecular potential (see Ref. [10] for details).

2. Calculation for the H₂O-H₂ system

Calculations were performed on the H₂O-H₂ collision system using the MCRB formalism for the rotational band, 1-4 quanta of exchange for the 3 fundamental bands, as well as the 301-000 band,



allowing a study of the vibrational dependence of the half-width and the line shift (all parameters used in the calculations can be found in Ref. [11]). The calculation used *ab initio* reduced matrix elements from Lamouroux *et al.* [10]. The H₂O energy levels were taken from the MARVEL list [12].

For a number of transitions, the vibrational dependence showed unphysical behavior, with half-widths too low by as much as ~40%. By looking more closely at the problematic transitions, it was found that the sum rules were not obeyed either for the upper or lower states (or both). In their study of vibrational dependence of H₂O, Gamache and Hartmann [13] showed that the spectroscopic effects, resulting from the modification of the wavefunction and the energy levels (giving different energy gaps) when the vibrational band changes are small. Given this information, test calculations were made using the ground state RMEs for both the upper and lower states. Note, the H₂O energy levels were fixed from MARVEL, making the spectroscopic effect even smaller.

Figure 1 presents the percent difference between the calculated half-width using the ground state RMEs and the vibrational dependent RMEs, for some water vapor transitions perturbed by hydrogen. The selected transitions are taken from the ν_1 , $2\nu_1$, ν_2 , $2\nu_2$, $3\nu_2$, ν_3 , $2\nu_3$, $3\nu_3$, $4\nu_3$ and $3\nu_1+\nu_3$ bands. In Figure 1, transitions for which the upper and lower states obey the sum rules are shown as blue solid circles. The transitions for which the sum rules are not obeyed for all states are shown as red solid triangles. This figure demonstrates that using ground state RMEs does not introduce significant error to the calculations. Thus, calculations made in the future will employ the ground state RMEs.




Figure 1. Percent difference between the calculated half-width using the ground state RMEs and the vibrational dependent RMEs for some transitions of the H₂O-H₂. The transitions that obey the sum rules are the blue solid circles, the transitions that does not obey the sum rules are the red solid triangles.

3. Conclusion

Calculations were performed for water vapor transitions broadened by hydrogen using the vibrational dependent RMEs and using the ground state RMEs for both upper and lower states. The percent difference between the calculated half-width using the ground state RMEs and the vibrational RMEs was computed for a large number of transitions. Small differences were observed for the transitions that obey the sum rules, showing a small spectroscopic effect of the reduced matrix elements.

The effects of using vibrationally dependent RMEs is undergoing further investigation. Until the study is finalized ground-state RMEs will be utilized in the MCRB calculations.

Acknowledgements

The authors are pleased to acknowledge support of this research by the National Science Foundation through Grant No. AGS-1622676. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

References

- [1] Robert, D. and J. Bonamy, Short range force effects in semiclassical molecular line broadening calculations 1979 *J. Phys. France* **40** 923
- [2] Ma, Q., R.H. Tipping, and C. Boulet, Modification of the Robert-Bonamy formalism in calculating Lorentzian half-widths and shifts 2007 *J. Quant. Spectrosc. Radiat. Transfer* **103** 588
- [3] Ma, Q., R.H. Tipping, and R.R. Gamache, Uncertainties Associated with Theoretically Calculated N₂-Broadened Half-Widths of H₂O Lines 2010 *Mol. Phys.* **108** 2225
- [4] Lynch, R., R.R. Gamache, and S.P. Neshyba, Fully Complex Implementation of the Robert-Bonamy Formalism: Halfwidths and Line Shifts of H₂O Broadened by N₂ 1996 *J. Chem. Phys.* **105** 5711
- [5] Lamouroux, J., et al., Semiclassical calculations of half-widths and line shifts for transitions in the 30012←00001 and 30013←00001 bands of CO₂ II: collisions with O₂ and Air 2012 *J. Quant. Spectrosc. Radiat. Transfer* **113** 991
- [6] Gamache, R.R. and L. Rosenmann, The Effects of Velocity Averaging in Broadening Coefficient Calculations 1994 *J. Mol. Spectrosc.* **164** 489
- [7] Gamache, R.R., Analytical evaluation of the Maxwell-Boltzmann velocity average in pressure-broadened half-width calculations 2001 *J. Mol. Spectrosc.* **208** 79
- [8] Tran, H., et al., Collisional parameters of H₂O lines: Velocity effects on the line shape 2007 *J. Quant. Spectrosc. Radiat. Transfer* **108** 126
- [9] Gamache, R.R., A.L. Laraia, and J. Lamouroux, Half-widths, their temperature dependence, and line shifts for the HDO-CO₂ system for applications to planetary atmospheres 2011 *Icarus* **213** 720
- [10] Lamouroux, J., R.R. Gamache, and D.W. Schwenke, Determination of the reduced matrix elements using accurate ab initio wavefunctions: Formalism and its application to the vibrational ground state (000) of H₂¹⁶O 2014 *J. Quant. Spectrosc. Radiat. Transfer* **148** 49
- [11] Renaud, C.L., et al., Line shape parameters for the H₂O–H₂ collision system for application to exoplanet and planetary atmospheres 2018 *Icarus* **306** 275
- [12] Tennyson, J., et al., IUPAC critical evaluation of the rotational–vibrational spectra of water vapor, Part III: Energy levels and transition wavenumbers for H₂¹⁶O 2013 *J. Quant. Spectrosc. Radiat. Transfer* **117** 29
- [13] Gamache, R.R. and J.-M. Hartmann, Collisional parameters of H₂O lines: effects of vibration 2004 *J. Quant. Spectrosc. Radiat. Transfer* **83** 119