

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

What Are the Main Risk Factors for Lower Extremity Running-Related Injuries? A **Retrospective Survey Based on 3669 Respondents**

Sanfilippo, Damien; Beaudart, Charlotte; Gaillard, Allan; Bornheim, Stephen; Bruyere, Olivier; Kaux, Jean François

Published in: Orthopaedic Journal of Sports Medicine

DOI: 10.1177/23259671211043444

Publication date: 2021

Document Version Publisher's PDF, also known as Version of record

Link to publication

Citation for pulished version (HARVARD):

Sanfilippo, D, Beaudart, C, Gaillard, A, Bornheim, S, Bruyere, O & Kaux, JF 2021, 'What Are the Main Risk Factors for Lower Extremity Running-Related Injuries? A Retrospective Survey Based on 3669 Respondents', Orthopaedic Journal of Sports Medicine, vol. 9, no. 11. https://doi.org/10.1177/23259671211043444

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

What Are the Main Risk Factors for Lower Extremity Running-Related Injuries?

A Retrospective Survey Based on 3669 Respondents

Damien Sanfilippo,^{*†‡} MD, Charlotte Beaudart,^{†§} PhD, Allan Gaillard,[†] MSc, Stephen Bornheim,[†] PhD, Olivier Bruyere,[§] PhD, and Jean-François Kaux,^{†‡} MD, PhD Investigation performed at University and University Hospital of Liège, Liège, Belgium

Background: Despite the many studies on running-related injuries (RRIs), risk factors for injury remain unclear in the literature. **Purpose:** To investigate the risk factors of RRIs.

Study Design: Case-control study; Level of evidence, 3.

Methods: An online survey was conducted among 3669 injured and noninjured runners. Injury was defined as pain of various kinds, without attention to its consequences on running practice. The survey included 41 questions on 5 main categories—personal characteristics, daily lifestyle, training and running characteristics, practice of other sporting activities, and prevention habits—as well as information about the occurrence of RRI over the previous 12 months. Continuous and qualitative variables were analyzed by Student *t* test and chi-square test, respectively. Sixteen variables were selected for multivariate binary logistic analysis.

Results: Among the 3669 runners, 1852 (50.5%) reported at least 1 injury over the previous 12 months. Overuse injuries were largely represented (60.6%). The variables associated with RRIs that remained significant in the fully adjusted model were previous injury (odds ratio [OR], 1.62; 95% CI, 1.42-1.86), higher weight (OR, 1.006; 95% CI, 1.00-1.012), competitive running (OR, 1.53; 95% CI, 1.19-1.98), running >2 h/wk (OR, 1.28; 95% CI, 1.01-1.62), running >20 km/wk (OR, 1.25; 95% CI, 1.001-1.55), and stretching before running (OR, 1.46; 95% CI, 1.25-1.71).

Conclusion: Previous injury remains the most relevant risk factor for RRIs according to the current study and previous data. Many training characteristics seem to be involved but still have to be confirmed in view of conflicting data in the literature. Further research would help clinicians better understand RRIs and how to prevent them.

Keywords: running-related injuries; prevalent injuries; risk factors; survey

Over the past few decades, there has been increased focus on healthier lifestyles in developed countries. Physical activities in daily life and running activities in particular have become popular because of their ease of implementation, low cost, and reputation of being beneficial for health.^{6,19,23,27}

Endurance running efficiently provides substantial beneficial effects on body mass, body fat, resting heart rate, maximum amount of oxygen, triglycerides, and highdensity lipoprotein cholesterol. It has also been shown that the longer the length of training, the higher the achieved health benefits.⁹ However, endurance running is associated with a high rate of injuries, especially lower extremity injuries. A large recent systematic review⁵ highlighted an injury prevalence in runners of $43\% \pm 19.8\%$. The large variability observed in these results may be explained by differences in the characteristics of the target populations investigated as well as in the definition of running-related injury (RRI) used across studies.⁸

Acute injuries (ankle strain, fracture) can occur, but most of the time, runners experience overuse injuries.¹⁰ These have been defined as an injury to the muscularskeletal system that results from repetitive submaximal loading applied over a prolonged period, causing strain that goes beyond the structure's ability to adapt.²² This can limit running speed, distance, duration, or frequency for at least 1 week.^{10,14} The predominant site of lower extremity overuse injury is the knee,²³ and the most common injury is patellar-femoral pain syndrome. Other common overuse injuries include stress fractures, medial tibial stress syndrome, plantar fasciitis, and patellar and Achilles tendinopathies.¹⁰

Multiple authors have tried to identify the cause and risk factors of RRI to reduce the injury rate. Risk factors are

The Orthopaedic Journal of Sports Medicine, 9(11), 23259671211043444 DOI: 10.1177/23259671211043444 \circledcirc The Author(s) 2021

This open-access article is published and distributed under the Creative Commons Attribution - NonCommercial - No Derivatives License (https://creativecommons.org/ licenses/by-nc-nd/4.0/), which permits the noncommercial use, distribution, and reproduction of the article in any medium, provided the original author and source are credited. You may not alter, transform, or build upon this article without the permission of the Author(s). For article reuse guidelines, please visit SAGE's website at http://www.sagepub.com/journals-permissions.

usually classified into 2 categories: intrinsic (anatomic, biomechanical, and global health) and extrinsic (training characteristic). Most studies^{8,11,20,23} suggest that previous injuries are the most well-established RRI risk factor. However, these injuries are not the result of an isolated variable but rather the consequence of a structural and functional combination of different factors, and this is the main difficulty for clinicians and researchers.⁷ Other risk factors are still not clear in the literature.

The goal of our study was therefore to investigate the risk factors associated with injury in a population of runners. With this survey study in a large population, our aim was to confirm that previous injury is a risk factor for further RRI. Our secondary aim was to investigate a large number of variables as potential risk factors.

METHODS

This study followed the guidelines for reporting observational studies from the STROBE checklist (Strengthening the Reporting of Observational Studies in Epidemiology).²⁵ We conducted an online survey among runners to identify the potential risk factors for lower extremity RRI. As the participants completed the survey anonymously and on a volunteer basis, this study did not require ethical committee approval.

Population

To be included in this study, runners had to be at least 18 years old and have at least 12 months of running experience. After completing the survey, eligible participants were placed into an injured or noninjured group depending on whether they had experienced a lower extremity RRI within the previous 12 months. Injury was considered any kind of pain, without paying attention to its consequences on running practice.

The link to the survey was published on social media platforms. Several running event organizers also agreed to send the link to their large mailing lists. Numerous running magazines published the survey on their websites, and the survey was promoted at multiple running events. Data were collected from April to July 2017.

Development of the Survey

Our survey was developed through the following steps: (1) We investigated the literature to identify categories of

potential risk factors, and (2) we then contacted experts (doctors, physical therapists, running coaches) and asked them to validate our categories or generate additional ones using their expertise. Finally, we developed a survey including all categories identified in steps 1 and 2. The questionnaire was developed by translating items into questions. It was made up of 35 general questions plus 6 questions related to injured runners. The questionnaire was composed of a majority of closed but also semiclosed and open questions.

The survey was divided into 5 parts.

- Personal characteristics: sex, age, weight, height, laterality, previous injury
- Daily lifestyle: professional activity type, weekly work hours, daily sleeping hours, dietary habits, tobacco/ alcohol consumption
- Training and running characteristics: running speed, mileage and hours per week, weekly number of workouts, types of practice, trail versus road and drop, years of practice, goals of practice, annual number of competitions and distance, hydration behavior, forefoot versus rearfoot strike, planning methodology, fasted running sessions or not
- Practice of others sports activities and prevention habits Occurrence of RRI over the last 12 months: 1 question allowed us to categorize athletes as injured or noninjured ("I have suffered from an RRI in the last 12

Statistical Analysis

months").

Normality of continuous variables were investigated using histogram, Q-Q plot, comparison of mean with median, and the Shapiro-Wilk test. As all the continuous variables followed a normal distribution pattern, data were expressed as means and standard deviations. Furthermore, categorical variables were expressed as absolute (number) and relative (percentage) frequencies. To identify possible associations between risk factors and RRI, we initially performed a univariate analysis using the generalized estimating equation for each independent variable, with occurrence of RRI as the dependant variable. Continuous and qualitative variables were analyzed by Student t test and chi-square test, respectively. All variables that had significant associations (P < .05) in the univariate analysis were selected for inclusion in the multivariate binary logistic analysis. The results were expressed as

*Address correspondence to Damien Sanfilippo, MD, Department of Rehabilitation and Sports Sciences, University of Liège, Liège, Belgium (email: d.sanfilippo@chuliege.be).

[§]World Health Organization Collaborating Center for Public Health Aspects of Musculoskeletal Health and Ageing, Department of Public Health, Epidemiology and Health Economics, University of Liège, Liège, Belgium.

Final revision submitted April 20, 2021; accepted May 25, 2021.

Ethical approval was not sought for the present study.

[†]Department of Rehabilitation and Sports Sciences, University of Liège, Liège, Belgium.

[‡]Department of Physical Medicine and Sports Traumatology, Sports², FIFA Medical Centre of Excellence, IOC Research Centre for Prevention of Injury and Protection of Athlete Health, FIMS Collaborative Center of Sports Medicine, University and University Hospital of Liège, Liège, Belgium.

The authors have declared that there are no conflicts of interest in the authorship and publication of this contribution. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

odds ratio (OR) and 95% confidence interval. All data analyses were performed using SPSS Statistics (Version 24.0; IBM).



Figure 1. Flowchart of included population.

RESULTS

Participants

The survey was completed by 4105 runners. However, 436 respondents were excluded: 428 did not provide information about injury, and 8 gave inconsistent answers to open questions related to injury. Therefore, 3669 respondents were included for statistical analysis (Figure 1): 2123 (57.8%) men and 1546 (42.2%) women. The mean age of the population was 36.1 ± 10.9 years, and the mean body mass index (BMI) was 22.9 ± 4.5 kg/m².

A total of 1852 runners (50.5%) reported at least 1 lower extremity RRI over the previous 12 months and were included in the injury group. Injured participants were significantly more often older (P < .001), heavier (P = .01), and male (P < .001). They also presented a higher prevalence of previous injuries beyond the previous 12 months as compared with noninjured participants (P < .001) and indicated a greater importance on diet (P = .03). No difference was found for height, BMI, laterality, tobacco/alcohol consumption, professional activity, or hours of work and sleeping time (Table 1).

Incidence and Typology of RRI

Most injured runners (69.5%) cited only 1 injury (Table 2). A total of 2538 injuries were reported, and 2218 pathologies

	$All \ (N=3669)$	Noninjured $(n = 1817)$	Injured $(n = 1852)$	P Value ^b
Female sex	1546 (42.1)	820 (45.1)	726 (39.2)	<.001
Age, y	36.1 ± 11.1	35.4 ± 11.3	36.8 ± 10.7	<.001
Weight, kg	67.9 ± 11.4	67.3 ± 11.5	68.4 ± 11.3	.01
Height, cm	172.9 ± 8.73	172.7 ± 8.7	173.2 ± 8.8	.09
Body mass index, kg/m ²	22.9 ± 4.60	22.7 ± 4.5	23.0 ± 4.6	.13
Tobacco use				.89
None	3482 (94.9)	1723 (94.8)	1759 (95)	
1-10 cigarettes/d	165(4.5)	82 (4.5)	83 (4.5)	
>10 cigarettes/d	22(0.6)	12(0.7)	10 (0.5)	
Alcohol use				.77
Never	504 (13.7)	256 (14.1)	248 (13.4)	
Once a week	2333 (63.6)	1146 (63.1)	1187 (64.1)	
More than once a week	832 (22.7)	415 (22.8)	417 (22.5)	
Sleep, h				.12
<6	263 (7.2)	116 (6.4)	147 (7.9)	
6-8	2855 (77.8)	1416 (77.9)	1439 (77.7)	
>8	551 (15.0)	285 (15.7)	266 (14.4)	
Importance given to diet				.03
Low	415 (11.3)	227 (12.5)	188 (10.2)	
Moderate	2517 (68.6)	1247 (68.6)	1270 (68.6)	
High	737 (20.1)	343 (18.9)	394 (21.3)	
Laterality: right	3178 (86.7)	1575 (86.7)	1603 (64.3)	.91
History of injury beyond the previous 12 mo	2149 (58.6)	957 (52.7)	1192 (35.6)	<.001

 $\begin{array}{c} {\rm TABLE \ 1} \\ {\rm Characteristics \ of \ the \ Study \ Participants}^a \end{array}$

^{*a*}Data are reported as No. (%) or mean \pm SD.

 ^{b}P value obtained from Student *t* test for continuous variables and from chi-square test for categorical variables. Bold *P* values indicate statistically significant difference between the noninjured and injured groups (*P* < .05).

were identified. The remaining 320 were classified as *other* because of unclear description. Among pathologies, overuse was widely represented (60.6%) and included patellofe-moral pain syndrome, Achilles tendon injury, medial tibial stress syndrome, plantar fasciitis, iliotibial band syndrome, stress fracture, and back pains. Traumatic injuries (fracture and strain) and muscular injuries (stiffness/tear) represented 19.1% and 20.3%, respectively.

TABLE 2 Information on Injuries a

	Injured Group $(n = 1852)$		
No. of injuries			
1	1288 (69.5)		
2	461 (24.9)		
3	88 (4.75)		
4	11 (0.6)		
5	4 (0.2)		
Type of injuries			
Overuse	928 (50.1)		
Traumatic	293 (15.8)		
Muscle	311 (16.9)		
Other	320 (17.3)		

^{*a*}Data are reported as No. (%).

Risk Factors for RRIs

Univariate analyses revealed that some lifestyle factors were significantly associated with the risk of injuries (Table 3). Regarding training characteristics, results showed higher risks for training >2 sessions per week, training >2 h/wk, running >21 km/wk, and running >10 km/h during training (P < .001 for all). We also found significant differences for type of practice (trail runners at higher risk vs road runners; P < .001), practice goals (runners preparing for a competition at higher risk vs runners not preparing; P < .001), number of competitions per year (higher risk if >5; P < .001), running surface (higher risk if on an athletic track [P < .001] or in the woods [P = .01]), vertical drop (higher risk if the usual route includes significant elevation; P < .001), stretching session (higher risk for the ones who stretch before running; P < .001), prevention technique (higher risk for those who performing prevention techniques; P < .001).

Based on the analysis, 16 variables were included in the multivariate binary logistic analysis, and some of these associations were maintained in the multivariate model (Table 4). The following variables increased the risk of injury: heavier weight (OR, 1.006; 95% CI, 1.00-1.012), presence of a previous injury (OR, 1.62; 95% CI, 1.42-1.86), participating in competitions (OR, 1.535; 95% CI, 1.19-1.98). running >2 h/wk (OR, 1.28; 95% CI, 1.01-1.62),

TABLE 3				
Injury Risk Factors: Univariate Analyses ^a				

	All $(N = 3669)$	Noninjured $(n = 1817)$	Injured $(n = 1852)$	P Value ^b
Years of practice	7.47 ± 7.74	7.45 ± 7.5	7.49 ± 7.9	.87
Type of practice				<.001
Trail	694 (18.9)	305 (16.8)	389 (21.0)	
Road	976 (26.6)	556 (30.6)	420 (22.7)	
Both	1999 (54.5)	956 (52.6)	1043 (56.3)	
Competition: yes	3262 (88.9)	1543(84.9)	1719 (92.8)	<.001
No. of races per y: >5	2160 (57.4)	944 (37.3)	1162 (62.7)	<.001
Sessions per wk: >2	2226 (60.7)	997 (54.9)	1229 (66.4)	<.001
Hours run per wk: >2 h	2778 (75.7)	1268 (69.8)	1510 (81.5)	<.001
Distance per wk: >20 km	2482 (67.6)	1108 (61.0)	1374 (74.2)	<.001
Pace: >11 km/h	1856 (50.6)	782 (44.9)	959 (53.0)	<.001
Drop: >100 m/10 km	2566 (69.9)	1166 (67.1)	1270(70.5)	<.001
Type of ground				
Woods	3180 (86.7)	1547 (14.9)	1633 (88.2)	.01
Road	2106 (57.4)	1049 (57.7)	1057 (57.1)	.69
Track	753 (20.5)	331 (18.2)	422 (22.8)	<.001
Other	34 (0.9)	19 (1.0)	15 (0.8)	.46
Stretching before running				<.001
Never	2747 (74.9)	1416 (77.9)	1331 (71.9)	
Yes, most of the time	795 (21.7)	346 (19.0)	449 (24.2)	
Yes, always	127(3.5)	55 (3.0)	72 (3.9)	
Stretching after running				.1
Never	921 (25.1)	481 (26.5)	440 (23.8)	
Yes, most of the time	1943 (53.0)	933 (51.3)	1010 (54.5)	
Yes, always	805 (21.9)	403 (22.2)	402 (21.7)	

^{*a*}Data are reported as mean \pm SD or No. (%).

 ^{b}P value obtained from Student t test for continuous variables and from chi-square test for categorical variables. Bold P values indicate statistically significant difference between the noninjured and injured groups (P < .05).

TABLE 4 Injury Risk Factors: Multivariate Analysis

	В	Odds		
	Coefficient	Ratio	95% CI	P Value ^a
Male sex	-0.11	0.99	0.85-1.15	.888
Age	0.001	1.001	0.99 - 1.008	.795
Weight	0.006	1.006	1.00 - 1.012	.039
Low importance of food	-0.21	0.81	0.65 - 1.002	.052
Previous injury: yes	0.485	1.62	1.42 - 1.86	<.001
Competition: yes	0.428	1.53	1.19 - 1.98	.001
Races per y: >5	0.143	1.15	0.99 - 1.35	.07
Sessions per wk: >2	0.037	1.037	0.86 - 1.25	.70
Hours per wk: >2 h	0.24	1.28	1.01 - 1.62	.041
Distance per wk: >20 km	0.22	1.25	1.001 - 1.55	.049
Pace: >11 km/h	0.139	1.15	0.99 - 1.34	.07
Drop: >100 m/10 km	-0.153	0.86	0.73 - 1.01	.86
Type of ground: woods	0.10	1.11	0.89 - 1.38	.34
Type of ground: track	0.12	1.13	0.95 - 0.34	.17
Stretching before: yes	0.38	1.46	1.25 - 1.71	<.001
Type of practice: trail	0.16	1.17	0.97 - 1.41	.10

^{*a*}Bold *P* values indicate statistical significance (P < .05).

running $>\!20$ km/wk (OR, 1.25; 95% CI, 1.001-1.55), and, surprisingly, stretching before running (OR, 1.46; 95% CI, 1.25-1.71).

DISCUSSION

In the context of the growing interest in running and the numerous injuries resulting from it, the goal of our study was to investigate the risk factors associated with injuries among a large and heterogeneous population of runners. Several risk factors were identified, but only some were confirmed through multivariate regression analyses: presence of previous injury, doing competitions, running >2 h/wk, running >20 km/wk, heavier weight, and, quite surprisingly, stretching before running.

In our sample of 3669 runners, we found a 50.5% prevalence of RRIs. The prevalence of injuries has been reported to range from 19.4% to 79.3% in 1 systematic review published in 2007.⁸ The large range of observed prevalence can be explained by several assessment biases. First, the RRI definition seems often to be poorly described. For example, Buist et al⁴ defined an injury when pain is responsible for a total stop of running for a week, while Hespanhol et al⁹ considered an injury to be a single training session not performed because of pain. In our study, we considered an injury as any kind of pain without paying attention to its consequences on running practice. Despite this very broad definition of injury, the percentage of injured runners remained modest as compared with other studies. Second, we observed large heterogeneity among studies in terms of the definition of runner and one's level of running. Indeed, most of the training factors and personal characteristics could be different when comparing, for example, recreational and elite runners.

In our study, more than half of the injuries were overuse ones (60.6%), which seems in accordance with the available

literature.^{13,18} Our results highlighted that the presence of previous injuries increased the odds of injury by 62%. This is the largest association found in our multivariate model. Studies^{11,23,24} have reported previous injury as one of the most common factors associated with RRI. However, Buist et al⁴ emphasized that studies did not used to pay attention to the nature of these previous injuries (eg, type of injury, connection or not with running practice). Furthermore, it would be very useful to know whether the recurrent injuries are the same or related.

In addition, the results of the current study indicated that runners who took part in competitions were more at risk to develop RRI, with an increased risk of 53%. We found that only 1 study, by Walter,²⁶ considered this risk factor. It was a prospective study with a cohort of 1680 runners monitored during a 12-month follow-up, and the authors noted a similar strength of association (risk ratio, 1.73 [95% CI, 1.21-2.49] for men and 1.93 [95% CI, 0.97-3.89] for women). Competitive runners will usually run at a faster speed, run more kilometers, and run more frequently. However, the association of competition with higher RRI risk seems confirmed in the multivariate analysis that considered these individual parameters. Moreover, when nonprofessional runners sign up for a competition, they often do it rather late, mainly because of their other activities (eg, family, professional activities, other sports). As a result, a significant increase in the volume and intensity of their practice is typically observed within a short period. Workout-planning error (ie, errors in the variation and distribution over time of the volume and intensity of training) may also be a major factor in the occurrence of RRI, but we did not investigate this parameter in our study.

Our survey included several questions on training characteristics. Training errors are considered to be responsible for a large part of RRI.^{2,8,12,26} According to our data about weekly mileage, running >20 km/wk multiplied runners' risk of having an injury by 1.25. In the same vein, we observed that the risk of injury is increased for runners who perform >2 hours of running per week. In 2 prospective studies, Walter²⁶ and Macera¹⁴ provided strong evidence that a weekly distance >64 km increased the risk of RRI. Conversely, Kelsey et al¹² and many other studies found that either there was no connection or longer distances had a protective effect. Bovens et al,² who expressed rate of injuries on 1000 hours of exercise, reported that the number of injuries decreased when weekly mileage during an 18-month period increased from 15 to 37 miles per week. Accordingly, in a recent systematic review that took into account data from 36 studies, Hulme et al¹¹ found no compelling evidence to support whether longer or shorter absolute running distances affect RRI. In this same review, the authors concluded, using 8 prospective and retrospective studies, that there was not enough scientific evidence of any relation between duration and RRI development.

Univariate and multivariate analysis uncovered connections with weight. The heavier runners would be more at risk of RRI. In contrast, no significant differences were found for BMI. Most of the studies did not find any connection with weight,^{11,26} while others reported some conflicting data. Our data are consistent with several studies^{11,15,23} showing no significant connections between BMI and RRI. In contrast, Nielsen et al,¹⁶ who performed a 1-year follow-up of 930 novice runners, stated that runners with a BMI >30 kg/m² were significantly more at risk to injury. Buist and Bredeweg³ noted similar results: increased risk by 31.8% of sustaining an RRI among novice runners with a BMI >25 kg/m² versus <25 kg/m². Faced with the conflicting results, more studies should be performed to assess this point.

We found more injured runners among the ones who are used to stretching before training sessions or at another time. Shrier²¹ conducted a meta-analysis on this variable. Among 17 studies that examined stretching before running, 7 suggested that it was beneficial, 3 indicated that it was detrimental, and 7 noted no difference. When results of all these studies are pooled, the clinical evidence does not support the evidence that stretching before exercise prevents injury. Authors also highlighted that stretching at another time than just before or after the race may have a protective effect. However, only 2 studies isolated the effect of stretching after running or at another time. From our point of view, it is very hard to study the effect of stretching because of numerous variables (number of muscles stretched, tension level, duration, number of repetitions, active or passive stretches, etc). Even if we frequently see in clinical practice that athletes who perform stretching have already been injured and stretch precisely to avoid any new injury, our results did not support this hypothesis. Indeed, results revealed that the contribution of stretching beforehand was independent of history of injury.

One of the particularities of our study was to study RRI incidence between road runners and trail runners. Indeed, we did not find a single study that focused on this aspect. However, once again, these 2 kinds of practices included several variables, and it would be very difficult to extract clear data out of this study. We found significantly more injured runners among trail runners, but the relationship was not confirmed in multivariate analysis. Several points could be discussed regarding this relationship. First, it would be interesting to study injury typology and incidence of traumatic injuries (eg, ankle and knee strain, fracture). Without considering traumatic injuries, this relationship would nevertheless be conserved, explained by the course's topography (eg, more demanding ascents and descents) as well as the ground's topography (importance of the muscles supporting the pelvis and the back when the ground is not flat). Moreover, we found a significant difference regarding vertical drop (ie, more injured runners with higher vertical elevation). Furthermore, the theoretical protective effect of running on a soft surface was discussed in 1 study in which authors measured vertical acceleration of the tibia on 35 runners who ran on different surfaces at different speeds.¹ In contrast, Hulme et al¹¹ gathered results from 19 studies focusing on this point and concluded that topographic features and the compliancy of particular surfaces did not affect RRI risk. Our data on surface uncovered more injured runners among those who perform some athletic track sessions. This result makes sense because track

racers usually do a lot of interval running. In hindsight, it would have been interesting in our questionnaire to consider this type of training as a potential risk factor of RRI.

We consider that the main strength of this study was the large number of respondents and variables studied. Furthermore, to our knowledge, no studies have compared injury rate between trail runners and road runners. There are, however, some limitations to this study. First, it was a retrospective study, and runners had to remember a lot of information about last 12 months, leading to a recall bias. Second, the injured status was self-reported. There was no medical diagnosis allowing for the objectification of the injury. Last, regarding training characteristics, we took into account the mean value for pace, mileage, and frequency over a period of time. For these parameters, we discussed the importance of variation over time to explain injury occurrence.

In light of these last sections, the link between training characteristics and running injuries seems to be very complex. We discussed several methodological discrepancies that can explain some of this complexity. Some other points could be considered. Theoretically, injury occurs when cumulative training load, over 1 or several runs, exceeds the capacity of various tissues to perform adaptive repair.¹⁰ Accordingly, this training load should be evaluated at each training session. Furthermore, this training load will have a different impact on tissues depending on distribution over a period of time. This final point explains why fixed or absolute data, such as mean weekly values, seem to be inappropriate. Nielsen et al¹⁷ conducted a systematic review on this point (training errors) and came up with the hypothesis that a complex relationship involving distance, frequency, duration, and pace should exist and has not been investigated yet. It is probably the reason why data recorded until now have not allowed us to better understand these running-related risk factors.

CONCLUSION

This study confirms that previous injury remains the primary risk factor for RRI. Other main risk factors for RRIs have been highlighted in our survey: being a trail runner, doing competitive running, running >2 h/wk, and stretching before running. Future targeted and prospective studies should be able to improve risk factor knowledge and result in the implementation of effective preventive measures.

ACKNOWLEDGMENT

The authors thank all participants who kindly responded to the survey.

REFERENCES

 Boey H, Aeles J, Schütte K, Vanwanseele B. The effect of three surface conditions, speed and running experience on vertical acceleration of the tibia during running. Sports Biomech. 2017;16(2):166-176.

- Bovens AMP, Janssen GME, Vermeer HGW, Hoeberigs JH, Janssen MPE, Verstappen FTJ. Occurrence of running injuries in adults following a supervised training program. *Int J Sports Med.* 1989;10(suppl 3): S186-S190.
- Buist I, Bredeweg SW. Higher risk of injury in overweight novice runners. Br J Sports Med. 2011;45(4):338-338.
- Buist I, Bredeweg SW, Lemmink KAPM, Van Mechelen W, Diercks RL. Predictors of running-related injuries in novice runners enrolled in a systematic training program: a prospective cohort study. *Am J Sports Med.* 2010;38(2):273-280.
- Francis P, Whatman C, Sheerin K, Hume P, Johnson MI. The proportion of lower limb running injuries by gender, anatomical location and specific pathology: a systematic review. *J Sports Sci Med*. 2019;18(1): 21-31.
- Fredericson M, Misra AK. Epidemiology and aetiology of marathon running injuries. Sports Med. 2007;37(4-5):437-439.
- Gijon-Nogueron G, Fernandez-Villarejo M. Risk factors and protective factors for lower-extremity running injuries a systematic review. J Am Podiatr Med Assoc. 2015;105(6):532-540.
- Hespanhol LC Jr, Pena Costa LO, Lopes AD. Previous injuries and some training characteristics predict running-related injuries in recreational runners: a prospective cohort study. *J Physiother*. 2013;59(4): 263-269.
- Hespanhol LC Jr, Pillay JD, van Mechelen W, Verhagen E. Metaanalyses of the effects of habitual running on indices of health in physically inactive adults. *Sports Med.* 2015;45(10):1455-1468.
- Hreljac A. Etiology, prevention, and early intervention of overuse injuries in runners: a biomechanical perspective. *Phys Med Rehabil Clin N Am.* 2005;16(3):651-667.
- Hulme A, Nielsen RO, Timpka T, Verhagen E, Finch C. Risk and protective factors for middle- and long-distance running-related injury. *Sports Med.* 2017;47(5):869-886.
- Kelsey JL, Bachrach LK, Procter-Gray E, et al. Risk factors for stress fracture among young female cross-country runners. *Med Sci Sports Exerc*. 2007;39(9):1457-1463.
- Lopes AD, Hespanhol LC, Yeung SS, Costa LOP. What are the main running-related musculoskeletal injuries? *Sports Med.* 2012;42(10): 891-905.

- Macera CA. Predicting lower-extremity injuries among habitual runners. Arch Intern Med. 1989;149(11):2565-2568. doi:10.1001/archinte.1989.00390110117026
- Malisoux L, Ramesh J, Mann R, Seil R, Urhausen A, Theisen D. Can parallel use of different running shoes decrease running-related injury risk? *Scand J Med Sci Sports*. 2015;25(1):110-115.
- Nielsen RO, Buist I, Parner ET, et al. Predictors of running-related injuries among 930 novice runners: a 1-year prospective follow-up study. Orthop J Sports Med. 2013;1(1):2325967113487316.
- Nielsen RO, Buist I, Sørensen H, Lind M, Rasmussen S. Training errors and running related injuries: a systematic review. *Int J Sports Phys Ther.* 2012;7(1):58-75.
- Roberts WO. A 12-yr profile of medical injury and illness for the Twin Cities Marathon. *Med Sci Sports Exerc.* 2000;32(9):1549-1555.
- Rothschild CE. Primitive running: a survey analysis of runners' interest, participation, and implementation. *J Strength Cond Res.* 2012; 26(8):2021-2026.
- Saragiotto BT, Yamato TP, Hespanhol Junior LC, Rainbow MJ, Davis IS, Lopes AD. What are the main risk factors for running-related injuries? *Sports Med.* 2014;44(8):1153-1163.
- Shrier I. Does stretching help prevent injuries? In: MacAuley D, Best T, eds. *Evidence-Based Sports Medicine*. 2nd ed. Blackwell Publishing; 2007:36-58.
- 22. Stanish WD. Overuse injuries in athletes: a perspective. *Med Sci Sports Exerc.* 1984;16(1):1-7.
- Van Gent RN, Siem D, Van Middelkoop M, Van Os AG, Bierma-Zeinstra SMA, Koes BW. Incidence and determinants of lower extremity running injuries in long distance runners: a systematic review. *Br J Sports Med*. 2007;41(8):469-480.
- Van Mechelen W. Running injuries: a review of the epidemiological literature. Sports Med. 1992;14(5):320-335.
- Von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *PLoS Med.* 2007;4(10):1623-1627.
- 26. Walter SD. The Ontario cohort study of running-related injuries. *Arch Intern Med.* 1989;149(11):2561.
- Willems TM, Witvrouw E, De Cock A, De Clercq D. Gait-related risk factors for exercise-related lower-leg pain during shod running. *Med Sci Sports Exerc.* 2007;39(2):330-339.